

Williams Lake Timber Supply Area Forest Health Strategy 2021/2022



Ministry of
Forests, Lands, Natural
Resource Operations
and Rural Development

Endorsed By:

Jennifer Bowman
A/District Manager
Cariboo-Chilcotin Natural Resource
District

Prepared by: Nicole Stamer, RPF

Date: Nov. 19, 2021

Foreword

The purpose of the Williams Lake TSA Forest Health Strategy 2021/2022 is to promote cooperative forest management between operational planners, reviewing agencies, and approval authorities of forest health risks, issues, and best management practices. This strategy is very important in developing specific management tactics to be utilized for current and future forest health factors.

Updates of Note 2020/2021:

- Beetle Management Unit Maps with Corresponding Strategies.
 - BMU shapes are found here: [2021-2022 BMU Shapes](#)
- Priority and ranking of *Forest Health Factors in the Williams Lake TSA (Table 2)*.
- The District will be following the original Biodiversity Update Note 7B for 2021/2022 in addition to recently approved Forest Stewardship Plans

Forest Health Factors to note for the coming field season are:

- The Cariboo experienced moderate burn intensity and less area burned by wildfire activity during the 2021 fire season, in comparison to the 2017 wildfires.
- Black Army Cutworm – Defoliation can occur when planting recently burned areas. Defoliation significance ranges from low to severe depending on sequence of events, distribution and host susceptibility, and detection and management.
- Douglas-fir beetle - There are concerns about the continuing impact of the 2017, 2018 and 2021 wildfires on the Douglas-fir beetle population.
 - Decreased IBD population compared to 2020.
- Drought – 2021 was one of the driest summers on record in the District. It is anticipated that more drought stress and mortality will become evident throughout 2022. There is also a potential for secondary insects to be attracted to stressed trees, causing further mortality.
 - Blowdown trees can attract and sustain IBD populations.
 - Continued bark beetle mitigation strategies on all harvest activities is crucial to maintaining IBD populations below epidemic levels.
- Western spruce budworm – Continued high levels of defoliation observed in the Pablo Creek/Sheep Creek area of the Fraser River Valley.
 - Egg mass sampling will be conducted fall 2021 to determine potential population for 2022. A spray program is most likely recommended.
- Western hemlock looper – 2021 spray program around Quesnel Lake was conducted. Effectiveness of the spray is TBD through egg mass sampling fall 2021.

- Flooding – Wetter than normal years throughout 2020/2021 has caused blowdown and windthrow events, particularly in areas associated with selective logging.

For more information on pests and diseases in the Williams Lake TSA review the [2019 Overview of Forest Health Conditions in Southern British Columbia](#)

TABLE OF CONTENTS

	Page
<i>Foreword</i>	2
1. Introduction	5
2. Guiding Principles	5
3. Forest Health Objectives.....	6
4. Roles and Responsibilities.....	6
5. Priority Forest Health Factors in the WLTSA.....	7
6. Forest Health Factors and Management Directions.....	8
Definitions	8
Bark Beetles	10
Defoliators.....	14
Weevils.....	17
Diseases.....	18
Abiotic Injuries	22
7. Bark Beetle Management	24
8. Non-Recoverable Losses	25
9. Reducing the Impacts of Climate Change	26
10. Key TSR Issues	27
11. Web Links for Resources.....	28
12. References	30

Tables

Table 1: Implementation schedule for Forest Health meetings	7
Table 2: Priority of the Forest Health Factors within the WLTSA	8
Table 3: Priority for pine salvage based on stand characteristics and level of beetle kill (Modified from McLennan 2003) (Eng 2004).....	13
Table 4: Summary of estimated non-recoverable losses (unsalvaged) by damaging agent 2009-2019	26

Appendices

Appendix 1: Bark Beetle Population Trends 1999-2019	32
Appendix 2: Beetle Management Units and WLTSA Strategies	35

1. Introduction

This strategy recommends actions to address forest health issues in the Williams Lake Timber Supply Area (WLTSA) also known as the Cariboo-Chilcotin Natural Resource District. It is one of the largest Timber Supply Areas (TSAs) in the province, covering approximately 4.93 million hectares. There are three general landscape types: the Chilcotin Plateau, the central portion of the TSA and the eastern portion of the TSA. The Chilcotin Plateau, located west of the Fraser River, is characterized by a drier climate with extensive lodgepole pine forests and some Douglas-fir. The central portion of the TSA, located both east and west of the Fraser River, has mixed-species forests primarily Douglas-fir and lodgepole pine as well as a large portion of dry-belt Douglas-fir stands. The eastern portion of the TSA increases in elevation and moisture to forests of spruce, pine, western redcedar, western hemlock and subalpine fir.

The intent of this document is to provide guidance and information to operational planners, reviewing agencies and approval authorities on forest health risks, issues, and best management practices. It outlines the current status of forest health factors and key strategies and actions to minimize losses from damaging agents. The objective is to enhance ecosystem health by improving forest resiliency and sustainability. By providing acceptable approaches to forest health management this document attempts to simplify the planning and approval process for forest health treatments.

2. Guiding Principles

The WLTSA strategy and Forest Professionals¹ are to be consistent with all relevant legislation and planning guidance including:

- Chief Forester guidance in the [Williams Lake TSA Rational for Allowable Annual Cut \(AAC\) Determination](#) (February 25, 2015);
- Chief Forester [Post –Natural Disturbance Forest Retention Guidance](#) (2017 Wildfires)
- Regional Manager [Post-Wildfire Salvage guidance to Licensees](#)
- Cariboo Region Delegated Decision Makers [Post-Wildfire Salvage Expectations for Land Use Designations in Cariboo Region](#)
- Chief Forester [Guidance on Landscape- and Stand-level Structural Retention in Large-Scale Mountain Pine Beetle Salvage Operations](#) (December 2005); and
- Cariboo-Chilcotin Land Use Plan (CCLUP), Government Actions Regulation (GAR) Orders pertaining to Ungulate Winter Range (#U-5-001, U-5-002 and U-5-003),

¹ Forest Professionals are Registered Professional Foresters, Registered Forest Technologists, Special Permit and Limited License Holders, Accredited Timber Cruisers, Accredited Timber Evaluators and Silvicultural Accredited Surveyors. Forest professionals ensure that the forests are managed sustainably. (<http://abcfp.ca/>)

and associated strategies;

Order for [Shallow/Moderate Snowpack MDWRs](#)

Order for [Transition/Deep Snowpack MDWRs](#)

- [Mule Deer Winter Range exemptions](#) for wildfire impacted stands
- [CCLUP Regional Biodiversity Conservation Strategy and Updates](#);

3. Forest Health Objectives

- Identify beetle management strategies (Suppression, Salvage, and Monitor) for each Beetle Management Unit (BMU). The 2021 BMU strategy maps are found in Appendix 2.
- Provide updated strategic guidance for the ongoing forest health management in the WLTA.
- Identify treatment strategies for forest health management.
- Facilitate co-operative planning between agencies and licensees.
- Establish short- and long-term harvest guidelines to best address opportunities, given the current pest incidences and infestation levels.
- Facilitate the development of scientifically and ecologically sound operational plans and practices.
- In conjunction with the various licensees and MFLNRORD identify areas of responsibility for beetle management. This includes where there may be opportunities for small tenure holders to assist in salvage/suppression efforts.

4. Roles and Responsibilities

- Licensees and FLNRORD will collaboratively undertake implementation of this strategy primarily through the WLTA Forest Health Committee.
- Meetings of this committee will be held on an as-needed basis to discuss the implementation and effectiveness of this strategy. An outline of meeting topics, dates and actions are shown in Table 1.
- Bark beetle treatment planning (detailed aerial surveys, probing, pheromone placement, single tree removal/disposal, trap tree placement/removal and larger scale treatments such as sanitation harvesting) is an iterative process involving the collaborative effort of licensees and FLNRORD in WLTA Forest Health Committee.

Table 1: Implementation schedule for Forest Health meetings

Topic	Meeting Dates	Actions
Planning	June - October	<ul style="list-style-type: none"> • Aerial overview and detailed flight information • Updated BMU strategies and boundaries • Identify areas of responsibility and planned activities based on infestation levels and capability • FLNRORD report on IBD treatments and monitoring evaluations
Implementation	August - March	<ul style="list-style-type: none"> • Discussion of planned and implemented beetle treatments • Update on western spruce budworm treatments of the previous year (if applicable), and egg mass sampling and treatment strategy • WLTS Forest Health Strategy in draft
Monitoring & Evaluation	January – June	<ul style="list-style-type: none"> • Licensee summary submission of Douglas-fir beetle treatment type and location • Evaluation of treatments and monitoring results with recommendations for future planning • Update on western spruce budworm treatment plans and target areas • Final version of WLTS Forest Health Strategy endorsed and released

5. Priority Forest Health Factors in the WLTS

The priority status of forest health agents is derived from the FLNRORD annual aerial [overview survey](#), from regional forest health specialists, district and branch forest health staff, and reports from licensees.

Within the WLTS pest species are ranked according to:

- distribution of pest and current incidence levels,
- available susceptible host species,
- landscape level hazard and risk information,
- known or suspected impacts on forest resource values,
- availability of operational detection and treatment methods, and

- costs and benefits of applying detailed detection and treatment activities.

Table 2: Priority of the Forest Health Factors within the WLTSA

High	Moderate	Low
Wildfire	Armillaria Root Disease	Forest Tent Caterpillar
Douglas-fir Beetle	Tomentosus Root Disease	Aspen Serpentine Leaf Miner
Western Spruce Budworm	Laminated Root Disease	Black Army Cutworm
Spruce Beetle	Western Gall Rust	Lodgepole Pine Terminal Weevil
Gypsy Moth	Comandra Blister Rust	Warren Root Collar Weevil
Douglas-fir Tussock Moth	Lodgepole Pine Dwarf Mistletoe	Stalactiform Blister Rust
Western Hemlock Looper	Post-wildfire Damage	Atropellis Canker
	Elytroderma Needle Cast	Lophodermella Needle Cast
	White Pine Weevil	Pine Needle Sheathminer
	Two Year Cycle Budworm	Flood Damage
	Drought Damage	Bear Damage
	Windthrow	Winter Desiccation
	Mountain Pine Beetle	Balsam Woolly Adelgid
	Western Balsam Bark Beetle	

6. Forest Health Factors and Management Directions

Definitions

The following definitions, from the [*Provincial Bark Beetle Management Technical Implementation Guidelines, Spring 2003*](#), are used for the purposes of this document regardless of any changes that may be made to the provincial definitions.

Beetle Management Unit (BMU): is a planning and reporting unit for operational beetle management. Its purpose is to facilitate the implementation of beetle management activities.

The following are strategies that may be employed within BMUs. An assigned strategy is based on the level of outbreak in an area and the estimated effectiveness of selected treatments in achieving stated objectives.

Suppression: This is the most aggressive strategy. It is selected when the infestation status is such that aggressive direct control actions are expected to keep an area at endemic population levels. Areas are lightly infested, and resources for direct control or harvesting and milling capacities equal or exceed the amount of the infestation. The intent of the strategy is to reduce or keep the outbreak to a size and distribution that can be handled within “normal resource capability”.

Objective: To reduce populations and maintain them at a relatively low level. Target is to treat 80 to 100% of the known infestation sites. All harvest and treatment is directed at currently attacked trees. The intent is to “control” outbreak in that area and stop spread.

Holding: The intent of this strategy is to maintain an existing outbreak at a relatively static level. It is a delaying strategy until adequate resources are available, or access is created to allow for a more aggressive approach, or to reduce overall loss while waiting for a killing climatic event. This is appropriate in areas with chronic beetle infestations that are too large to deal with using single tree treatments or where access is poorly developed for directed harvesting.

Objective: To maintain the infestation to a relatively static level by treating ~50 – 70% of the known infestation sites. The level of harvest and/or treatment is equal to the rate of infestation expansion. Harvesting should be concentrated in green attacked trees.

Monitor: This strategy is applied to areas where there are low population levels, where management efforts would be ineffective in substantially reducing the beetle population and subsequent levels of damage, or where there is no short term (less than 5 years) possibility of salvaging dead timber. This may be due to management constraints such as wilderness area, Parks or ecological reserve, or because access cannot be put in place before substantial merchantable degradation of the dead material occurs.

Objective: To record the change in attack level with no action required beyond monitoring and recording. This objective is appropriate in parks, ecological reserves and inoperable areas where the outbreak has peaked, salvage is not possible, and there is no current opportunity for any mitigation of further loss.

Salvage: This strategy is applied to areas where management efforts would be ineffective in substantially reducing beetle populations and subsequent levels

of damage. Such areas have extensive outbreaks covering a large proportion of susceptible stands. The objective in this case is to salvage affected stands and minimize value loss.

Objective: To salvage timber previously attacked to minimize value loss.

Indications are that holding the infestation static will fail due to influx of populations from heavily infested BMUs in proximity. Emphasis is more to retrieve values at risk and maximize Crown revenues by directing harvest towards killed stands prior to significant degradation. Treat <50% of the currently infested sites. This objective is appropriate in areas where suppression and holding actions are no longer appropriate or feasible.

Bark Beetles

The goal of bark beetle management is to minimize the spread of bark beetles and the loss of crown timber and to protect non-timber resource values. The [bark beetle hazard maps](#) were updated in 2019 for the WLTA.

Douglas-fir Beetle (IBD) *Dendroctonus pseudotsugae*

Douglas-fir beetle is causing significant damage to forests in the WLTA and is currently at outbreak levels. During outbreaks Douglas-fir beetle can kill large numbers of healthy trees over extensive areas. Outbreaks can have extremely negative impacts on ungulate winter range (MDWR), preservation of Old Growth Management Areas (OGMAs), watershed management and recreation and aesthetic values. Hot beds for beetle activity include along the Fraser River and near Williams Lake, with increases seen in Soda Creek, Mackin Creek, and Chimney Lake.

Detailed helicopter-GPS surveys have been conducted annually for heavily impacted areas of the WLTA that are good candidates for operational treatments. This [bark beetle survey](#) provides accurate GPS coordinates of infestation centres and polygons of spatially continuous infestations. The 2021 detailed flight identified 2300 infestation spots and 3 polygon infestations.

Stand management for Douglas-fir beetle:

- Follow [Best Practices for Managing Douglas-fir Beetle](#):
 - Introduction and Biology
 - Management Strategies and Tactics
 - Landscape and Stand Level Planning
 - Douglas-fir Beetle and Wildfire
- Sanitation harvest activities within constrained areas (OGMAs, RRZs, etc.) must be consistent with the CCLUP and applicable [Update Notes](#). These are ecologically sensitive areas, and it is imperative that any activity in these areas

is consistent with current biodiversity guidance and special care and attention is given to harvesting practices.

WLTSA Strategy: “Suppression” for some Douglas-fir dominated areas (see Appendix 2 for specifics), and “Salvage” for other non-currently infested areas.

Spruce Beetle (IBS) *Dendroctonus rufipennis*

Spruce stands are generally located in moist to wet ecosystems which result in faster decay and a short “shelf-life” of one to three years. However, Spruce beetle may take up to 3-4 years to kill a tree. Many spruce-leading stands have a high site index and if these stands are not harvested and re-forested promptly it could result in significant long-term losses in productivity. High site index stands should be prioritized for salvage of dead trees and suppression of infestation centres with an aim to balance forest health and mid-term timber supply. Wind felled spruce is an excellent habitat to spruce beetles and blowdown centres should be carefully monitored to ensure populations are dealt with before they start increasing and attacking standing healthy spruce.

Spruce beetle is difficult to detect in overview surveys, but they are still a useful tool in identifying general trends in beetle population levels. Attack declined in 2018, with most areas classified as trace or light. The areas to watch are Quesnel Lake and Horsefly Lake in the east, and Big Creek and Churn Creek in the south.

Although infestations appear to be declining, spruce beetle continues to be a species of great concern due to rapid spread, high attack ratios, and short “shelf-life” due to ecotypes where the beetle is located. The main treatments for spruce beetle are trap tree deployment and salvage harvest. Poor access, steep ground, and other economic challenges have led to very little harvesting of infested spruce. Trap trees pre- and post-harvest are important for sanitation harvest and capturing overwintering beetles.

WLTSA Strategy: “Monitor” for the majority of the WLTSA and “Holding” for some areas (see Appendix 2 for specifics).

Mountain Pine Beetle (IBM) *Dendroctonus ponderosae*

In the AAC Rationale for Williams Lake TSA the Chief Forester directs licensees “to continue to focus harvesting as much as possible on mountain pine beetle-impacted pine-leading stands in the Williams Lake TSA; and to harvest no more than their share of the AAC partition attributable to live tree volume.” This strategy should be followed while managing current and future forest values in context of sustainability.

Overall Objective: Salvage affected pine stands in a manner that will:

- maximize the economic value obtained from the killed trees,
- extend the salvage term (i.e. harvest those stands with the most dead wood), and

- expedite the restoration of impacted stands to the harvesting land base.

High Priority for Salvage (Unconstrained Landbase)

- Stands with at least 70 percent pine with high percentages of beetle-infested or killed stems
- Stands located west of the Fraser River
- Low volume stands (<100 cubic metres/ha)
- Pure pine stands with little or no advanced regeneration (especially high site index) to expedite stand recovery. (i.e. Ideal candidate areas for stand rehabilitation)
- Areas where shelf-life is considered short (i.e. wetter BEC zones)

Moderate Priority Salvage (Unconstrained Landbase):

- Areas where shelf-life is considered short
- >50% pine by Volume
- >30% beetle attack (Red green and grey combined)
- High/moderate susceptibility

Low Priority Salvage (Unconstrained and constrained areas)

- Mixed stands (< 50% pine)
- Maximize harvest of infested pine through selective harvest.
- Prescriptions should target pine removal rather than clearcut, where residual stands can be maintained in a wind firm condition to target the maximum volume of infested pine and to encourage natural regeneration of non-pine (climax) species especially where advanced regen exists in understory. Where more than one beetle species has infested mixed stand, then the rational should be explicit.
- OGMA's, MDWR's, riparian and other constrained areas in accordance with higher level plan guidelines, Land Use Orders and GAR orders.
- High amount of advanced regeneration
- Stands with suitable secondary structure (see Appendix 4 for definition)

The following table provides guidance to the placement of salvage areas on the landbase. It serves as guidance for salvage planning, but other values as listed above, should also be considered and rationalized in harvesting proposals.

Table 3: Priority for pine salvage based on stand characteristics and level of beetle kill
(Modified from McLennan 2003) (Eng 2004).

	Percentage of pine killed (Green, Red and Grey attack)				
Percentage of stand volume that is pine	<30%	30-50%	51-70%	>70% West of the Fraser River	>70% East of the Fraser River
<30%	No	No	No	No	No
30-50%	No	No	No	Low	Low
51-70%	Low	Low	Low	High	Moderate
>70%	Low	Moderate	Moderate	High	Moderate

WLTA Strategy: All BMUs within the WLTA have been identified as “salvage” strategy (Appendix 2) for Mountain Pine Beetle.

Western Balsam Bark Beetle (IBB) *Dryocetes confusus*

Stand management for Western Balsam Bark Beetle:

- To prevent high in-stand losses from IBB, manage to younger ages (i.e. harvest at 80 yrs) once >100 yrs very susceptible to attack.
- To some degree, stand edges promote more blowdown of BI than is typical in stands so after harvesting BI leading stands monitor blowdown and any beetle build-up.
- Trap trees have been shown to not be as successful for IBB management. Susceptible, standing trees attracted more beetles in comparison to felled trees (MacLauchlan et al. 2021).
- Stands can be baited prior to harvest to contain as many beetles in the proposed block as possible.
- Stands should be hazard rated and managed (planned for harvest & regeneration) based upon:
 - species (>50% BI)
 - age (older stands higher risk)
 - BEC – ESSF highest hazard in that most BI is found in this ecosystem but BI in the MS & SBS is very susceptible to attack because it is drought stressed more frequently and thus becomes easier to attack.
 - The ESSFdV and ESSFxc are highest hazard, followed by ESSFwc, mw, cv and dc.
 - elevation
 - current and past beetle attack

WLTA Strategy: “Monitor” with no treatments planned.

Defoliators

Defoliators, such as the western spruce budworm and 2-year cycle budworm, impact all age classes of their preferred hosts and can substantially impact growth and productivity and wood quality. Severe defoliation by these insects can result in mortality, especially in younger age classes, and have negative impacts on other stewardship values on the land base (e.g., ungulate ranges).

Western spruce budworm (IDW) *Choristoneura freemani*

Severe defoliation of mature stands typically requires 4-5 years of subsequent infestation to result in mortality, however, defoliation causes stress to the trees thereby reducing wood quality and quantity and predisposes Douglas-fir to attack by Douglas-fir beetle. When a mature stand is infested, understory mortality can be very significant. As a result of this understory mortality and Douglas-fir beetle mortality NSR stands are created.

FLNRO has been conducting aerial treatments of moderate to severe western spruce budworm populations using *Bacillus thuringiensis* var. *kurstaki* (B.t.k.) annually to protect the most valuable stands at risk. Detailed monitoring is conducted each year to determine population dynamics. Egg mass sampling occurs September to October to determine if threshold is met to initiate a spray program. Bud mining surveys are then conducted the following spring to estimate population levels and timing of spray applications.

This pest is of high significance as it not only has the potential to cause mortality and growth loss in the current growing stock, but also stresses the trees, increasing future vulnerability to bark beetle attack and furthering potential losses to mid-term timber supply.

Silviculture strategies:

- Plant higher densities to increase resilience in clearcut stands decreasing the overall susceptibility to severe and sustained western spruce budworm defoliation events.
- Thin dense understories to remove early instar food source

WLTSA Strategy: “Monitor” in conjunction with an aerial spray program. Detailed monitoring at the “moderate” egg mass sites to track larval development and populations levels and tree defoliation will be conducted.

Gypsy moth (IDM) *Lymantria dispar*

Gypsy moth is an invasive non-native species which has not become established in the WLTSA. This insect defoliates deciduous trees and has over 300 known hosts.

Introduction to an area could occur from recreational and other vehicles traveling in or from infested areas carrying pupae or eggs.

WLTSa Strategy: “Monitor” through the use of a trapping program in conjunction with District and Regional staff and the Canadian Forest Service. Early detection of this defoliator is critical for eradication treatments.

Douglas-fir Tussock Moth (IDT) *Orgyia pseudotsugata*

Douglas-fir tussock moth has the potential to cause significant mortality due to defoliation; although, top-kill, growth reduction, and secondary insects and fungi attacks may also occur. All ages of Douglas-fir are susceptible. Outbreaks typically occur every 10- 12 years and last around 4 years.

The outbreak in 2019 was significant as it is the furthest north that IDT has ever been recorded. This outbreak is centered in the Fraser River valley, near the community of Dog Creek.

WLTSa Strategy: “Monitor” in conjunction with an aerial spray program. Detailed monitoring at the identified sites to track larval development and populations levels and tree defoliation will be conducted.

Two Year Cycle Budworm (IDB) *Choristoneura biennis*.

In the interior wet belt two-year cycle budworm defoliates true firs and spruce species during “on” years (in the WLTSa this occurs during even years). This pest has the potential to create significant damage in the eastern portion of the WLTSa, with infestations covering over a million hectares recorded for the Cariboo during previous outbreaks. Potentially, plantations could suffer significant damage from this insect, and severe defoliation may predispose trees to attack by western balsam bark beetle or spruce beetle.

WLTSa Strategy: “Monitor” with no treatments planned.

Western Hemlock Looper (IDL) *Lambdina fiscellaria lugubrosa*

The western hemlock looper is a very destructive insect that defoliates western hemlock, western red cedar, Douglas-fir and spruce species. Untreated, it can lead to high levels of mortality during its two to four-year outbreaks. In severe cases, previously damaged stands may require stand rehabilitation. Generally, this insect has an eight-year cycle between high population levels. Historically outbreaks have been restricted to around Quesnel Lake.

UPDATE: Severe defoliation was observed around Quesnel and Horsefly Lakes in 2020. There also was also large number of moths observed around Williams Lake and Dog Creek/Alkali Lake, but with little to no visible defoliation.

WLTA Strategy: “Monitor” in conjunction with an aerial spray program. Detailed monitoring at the identified sites to track larval development and populations levels and tree defoliation will be conducted.

Forest tent caterpillar (IDF) *Malacosoma disstria*

One or more years of severe defoliation may result in top-kill, branch mortality, reduced radial growth and if the infestation persists, occasional mortality. Many of the infested stands are also infested with aspen serpentine leaf miner.

WLTA Strategy: “Monitor” with no treatments planned.

Aspen Serpentine Leaf Miner (ID6) *Phyllocnistis populiella*

Leaf miners tunnel between the epidermal layer of the leaf, reducing the leaf’s photosynthetic capacity. Outbreaks of this pest are fairly common in western North America. Short (1 or 2 year) outbreaks do not normally have major long-term impacts on tree growth; however, this insect has been active for several years in areas around the Williams Lake TSA. The majority of sites affected in 2018 were classified as moderate to severe.

WLTA Strategy: “Monitor” with no treatments planned.

Black Army Cutworm (IDA) *Actebia fennica*

In the absence of herbaceous plants and shrubs black army cutworm (IDA) will defoliate coniferous seedlings. Monitoring of IDA took place in 2018 due to the large area affected by wildfires in 2017. Populations were found to be low, but planting sites that are devoid of all vegetation should be avoided for 1-2 years after the wildfire.

BAC Management:

- Identify high risk sites
 - Burned openings south or west facing slopes are preferred for egg laying
 - Severe burns decrease natural vegetation which leads to higher risk of plantation defoliation
 - ESSF, MS, SBS, ICH, and IDF
 - Drought-prone sites
- Delay planting for 1 – 2 years
- Avoid spring planting

- Survey for IDA damage on natural vegetation in the spring
- Pheromone monitoring

WLTSA Strategy: “Monitor” with no treatments planned.

Pine Needle Sheathminer (IDI) *Zelleria haimbachi*

The pine needle sheathminer generally attacks juvenile to immature pine stands. Up to 100% of the new growth may be destroyed resulting in growth reduction.

WLTSA Strategy: “Monitor” with no treatments planned.

Balsam Woolly Adelgid (IAB) *Adelges piceae*

Balsam Woolly Adelgid generally attacks all true firs, with amabilis and grand fir attacked most frequently and subalpine fir most easily damaged. Damage includes swelling and distortion at buds and leader resulting in stunted growth, poor timber quality and tree death.

During a survey conducted in 2017 to determine the spread of IAB northward in BC, light infestation of IAB was found along the Horsefly River.

There is no large-scale treatment that can feasibly be employed for mature stands.

WLTSA Strategy: “Monitor” with no treatments planned.

Weevils

White Pine Weevil (IWS) (*Pissodes strobi*): Priority = Moderate

Lodgepole Pine terminal Weevil (IWP) (*Pissodes terminalis*): Priority = Low

Warren’s Root Collar Weevil (IWW) (*Hylobius warreni*): Priority =Low

White pine weevil and lodgepole pine terminal weevil can cause significant forking damage in young stands. Most of the mortality occurs in the first 10 years. The ICH and parts of the SBSdw1 are at high risk for white pine weevil. Warren’s root collar weevil can cause scattered mortality in young stands.

Silviculture strategies:

- Plant alternate species or species mixes.
- Leave non-target species; for example do not brush aspen.
- Plant higher densities.
- Plant genetically resistant stock for terminal weevil.

WLTSa Strategy: “Monitor” individual affected stands.

Diseases

Root Diseases

The old FPC Root Disease Guidebook (1996) has been updated to a new guidance document called “[Managing Root Disease in British Columbia](#)”. The updated and revised [root disease website](#) is also now available.

Armillaria Root Disease (DRA) *Armillaria ostoyae*

Armillaria root disease causes mortality in a wide range of conifers and deciduous trees and increases the susceptibility of infected trees to bark beetles and other pests. Douglas-fir, true firs, and spruce are highly susceptible to Armillaria root disease, western hemlock exhibits medium to high susceptibility, pines are moderately susceptible, and western red cedar and deciduous species have a lower susceptibility. Western larch also appears to be less susceptible to Armillaria after age twenty to forty.

Preharvest assessments for Armillaria root disease should be conducted as part of the site plan in all south facing ICH and SBSdw1 subzones and where root disease has been identified as part of the SP walkthrough in other subzones. South facing low elevation aspects within the SBSdw1 and ICH have the greatest hazard. These root diseases are absent or extremely rare on the Chilcotin Plateau east of Tatlayoko Lake. Some helicopter and ground traversed root rot shapefiles are available on the [RSI ftp site](#).

Management Strategies for Armillaria includes planting a mixture of less susceptible species or stumping where appropriate (refer to [root disease management guidebook](#)). Default stocking standards are not appropriate for root rot areas.

Tomentosus Root Disease (DRT) *Inonotus tomentosus*

Spruce is the primary host for Tomentosus root disease, with lodgepole pine exhibiting medium susceptibility. Infection can occur via spores, or from root to root contact between trees and stumps. Tomentosus root disease often goes undetected and can cause considerable growth loss in mature spruce stands. Tomentosus is high hazard in the drier SBS subzones and ICH. The presence of Tomentosus is easily determined by looking for advanced honeycomb decay in blowdown prior to harvest. Where spruce is prescribed for planting in stands to be harvested with a significant spruce component, a post-harvest stump assessment is recommended to identify the extent and location of Tomentosus within the stand.

Management strategies include avoidance planting and regenerating with a species other than spruce.

Laminated Root Rot (DRL) *Coniferiporia sulphurascens*

Laminated root rot is primarily a disease of Douglas-fir. It is most prevalent on warm south facing slopes in the ICH and SBSdw1. It forms discrete root rot centres or openings with associated windthrown and dead standing trees. Windthrown trees generally fall in random directions, have characteristic root balls, and laminar decay. Conduct preharvest assessments for laminated and armillaria root disease as part of the silviculture prescription in all ICH and SBS subzones and where root disease identified as part of the SP walkthrough in other subzones. Some helicopter and ground traversed root rot shapefiles are available on the RSM ftp site. South facing low elevation aspects within the SBSdw1 and ICH have the greatest hazard. These root diseases are absent or extremely rare on the Chilcotin Plateau east of Tatlayoko Lake.

Management Strategies for Laminated include planting a mixture of less susceptible species (pines and cedar) or stumping where appropriate (refer to root disease management guidebook).

Root Rot silviculture Strategy:

- Free growing assessments should be conducted after age 12 in high hazard root rot BEC zones. Root diseases are difficult to detect before age 12.
- Plant tolerant or resistant species
- Root Rot centres are more easily identified and mapped before harvest

Stem Rusts of Pine

Comandra blister rust (DSC) *Cronartium comandrae*

Stalactiform blister rust (DSS) *Cronartium coleosporioides*

Western gall rust (DSG) *Endocronartium harknessii*

DSS, DSC, and DSG all occur within the WLTSa. These pests are particularly important in young pine stands (<25 years of age). Comandra and stalactiform rust can only infect lodgepole pine from spores produced on the alternate host plant. These spores are typically only transmitted over short distances. The frequency of alternate plant hosts present on a site is likely an important risk factor for the *Cronartium* stem rusts. Western gall rust is the most common forest health factor affecting young lodgepole pine stands and does not require an alternate host.

WLTSa Strategy: “Monitor” young stands (esp. pure pine stands).

Silviculture strategy:

- Where pine is a preferred species, minimum stocking should be increased to 2000sp/ha to offset mortality due to rust.
- Plant alternate species where possible to minimize the risk of losses.

- Before considering spacing it is important to conduct pest surveys and to consider the impact of forest pests on final stocking levels. Delay free growing assessment to after 15 years in high hazard stands. Contact your regional forest specialist for hazard maps based on RESULTS data.

Atropellis Canker (DSA) (*Atropellis piniphila*)

Produces cankers with abundant sap on the main stem. Unlike dwarf mistletoe or stem rusts these cankers are not associated with squirrel feeding. Most infections occur on bark at 14-40 years of age. Infections much more common in high density pine stands and on the north side of the stem. When spacing pine stands older than 14 years old, space from north to south to aid in selection of disease-free trees.

Dwarf Mistletoe

Lodgepole Pine Dwarf Mistletoe (DMP) *Arceuthobium americanum*

Priority = Moderate

There are high incidences of lodgepole pine dwarf mistletoe in the SBPSxc where lodgepole pine is the only preferred species. The disease is caused by a parasitic plant and spread occurs from infected residual trees. Removal of all infected residual stems over 2m is the best way of reducing future infection. Preliminary modelling suggests that sanitation down to 0.5m may not be necessary if stands are being managed on short rotations. Lodgepole pine dwarf mistletoe hazard within the SBPSxc and IDFdk3 is very high. Refer to new Land Management Handbook for additional management information. Pre harvest stratification of blocks to identify post-harvest sanitation treatment areas is recommended.

WLTSa Strategy: "Monitor" young pine stands.

Harvest systems:

- Minimize the amount of dwarf mistletoe infected edges when laying out cut blocks and make use of natural barriers whenever possible (e.g. roads, non lodgepole pine timber type, water bodies, etc.).
- Avoid leaving wildlife tree patches and leave strips infected with dwarf mistletoe within or adjacent to the block whenever possible especially along road right of ways.
- Stands or portions of stands which are infested should be treated following harvest by removal of all residual lodgepole pine over 2m in height. Residuals will generally have dwarf mistletoe plants unless they were shaded prior to harvesting. On average it takes four years between infection and the first appearance of plants.

Needle Casts

Elytroderma Needle Cast (DFE) *Elytroderma deformans*

Elytroderma needle cast is easily recognized and well documented on Ponderosa pine but often goes misdiagnosed on Lodgepole pine. It is unique from other foliage disease in its ability to infect and persist in shoots where it can cause severe stunting and brooming. Elytroderma displays similar symptoms to Lophodermella but it can be distinguished by the fruiting bodies and stunting that it causes. Elytroderma is widespread in the IDF and SBPS, with wet low lying areas adjacent to swamps or lakes generally having a higher incidence. Mean Impact on stand volume in the IDFdk3 has been estimated to be 2% by age 40. Elytroderma causes irregular stem growth and large upturned branches that result in large elongated knots and reduced lumber quality even in trees with low disease severity. Conversion of mixed Fd/PI stands to PI plantations is not recommended within the IDFdk because of the impacts of this disease. Where clearcutting is practiced in mixed IDF forest types, maintain a component of Douglas-fir residuals and include Douglas-fir in the planting mix. An assessment for Elytroderma is recommended before spacing in lodgepole pine stands (especially in repressed pine stands).

Silviculture Strategy:

- Increase regeneration densities to 2000sph offset potential losses and damage from Elytroderma
- Manage for alternate species where appropriate for the site especially in low lying areas adjacent to riparian areas
- Dwarf mistletoe sanitation treatments can effectively remove infected advanced regen.
- Avoid using planting stock from higher elevations or drier locations than the area to be planted.

Lophodermella Needle Cast (DFL) *Lophodermella concolor*

When conditions are optimal (wet springs and summers), needle cast can infect large tracts of pine forest, particularly plantations. The long-term impact of this foliar disease is not completely understood, although permanent sample plots in affected plantations show significant growth reduction and mortality. This agent should be closely monitored, mapped, and the impact on plantations carefully recorded (esp. in the IDF & SBPS). Conditions were optimal in 2018 for high intensity infections in 2020.

WLTA Strategy: plant a diversity of species at the stand and landscape level

Dothistroma Needle Blight (DFS) *Dothistroma septosporum*

Dothistroma as a needle blight of pines that affects all ages of needles, particularly in the lower crown. It causes crown lift and can result in significant growth loss and even mortality when infection occurs in multiple years. Dothistroma is high hazard on lodgepole pine in the ICH, particularly in low lying areas adjacent to riparian areas. and in Limit the planting of lodgepole pine in the ICH except where needed to reduce the spread of root disease.

WLTSa Strategy: avoid planting lodgepole pine in the ICH except where needed to reduce the spread of root disease

Abiotic Injuries

Wildfire Damage and Post-Wildfire Mortality

The 2017 wildfires were the most extensive on record in BC's history with the Cariboo Region being the most severely impacted. These large wildfires affectsocial, economic and environmental factors which all must be considered when determining a treatment for an area.

Post-wildfire mortality due to the 2017 wildfires should have peaked in 2018.

WLTSa Strategy: Follow

- Chief Forester [Post –Natural Disturbance Forest Retention Guidance](#) (2017 Wildfires)
- Regional Manager [Post-Wildfire Salvage guidance to Licensees](#)
- Cariboo Region Delegated Decision Makers [Post-Wildfire Salvage Expectations for Land Use Designations in Cariboo Region](#)
- [Best Practices for Managing Douglas-fir Beetle](#) - Douglas-fir Beetle and Wildfire
- [Mule Deer Winter Range exemptions](#) for wildfire impacted stands

Drought Damage

We are experiencing more frequent and unpredictable summers in terms of dry, hot conditions. The 2021 heat spell resulted in some of the hottest days ever recorded in the District. It is anticipated that more drought stress and mortality will become evident throughout 2022. Major heat stress has been observed through discolouration of needles, and vast dropping of needles throughout the late summer and fall months in mature trees. Seedling mortality was evident after the major heat, as well as stress in younger stands.

There is also a potential for secondary insects to be attracted to these stressed trees and cause further mortality. Sites mapped for drought should be monitored closely when they are detected because the trees become stressed making them more susceptible to bark beetle infestations causing further mortality. Areas of drought should therefore be subject to ground checks if they are in suppression zones.

WLTA Strategy: Monitor closely for Douglas-fir beetle in impacted mature Douglas-fir stands. Plant drought resistant species. Minimize soil compaction resulting from harvesting activities and avoid harvesting areas with shallow soils. Avoid redirecting historical water flows as a result of road or ditch construction.

Windthrow

The area of windthrow recorded during the overview survey was 35 ha. The occurrence of windthrow can contribute to the maintenance and build-up of local spruce and Douglas-fir beetle populations.

Windthrow patches of Douglas-fir and spruce should be addressed promptly to minimize the expansion of beetle populations. It may be appropriate in some areas to leave blown-down trees on the ground until after the beetle flight and utilize them as trap trees. Windthrown trees should be removed prior to the beetle flight of the following year so that attacked blowdown does not contribute to increases in beetle attack.

WLTA Strategy: Allow windthrow to act as trap trees through summer and harvest before April 1st to prevent bark beetle population build-up. Alternative strategies are to use MCH, trap trees and/or fall and burn.

Flooding Damage

Flooding was extensive in 2019, 2020 and 2021. Sites mapped for flooding should be monitored closely when they are detected because the trees become stressed making them more susceptible to bark beetle infestations. Areas of flooding should therefore be subject to ground checks if they are located in suppression zones.

WLTA Strategy: “Monitor” with no treatments planned. Strategies are to use MCH, trap trees or fall and burn to combat associated IBD infestations.

Bear Damage

This damage has been occurring in lodgepole pine plantations in the eastern Cariboo and has been ongoing for several years.

WLTA Strategy: “Monitor” with no treatments planned.

7. Bark Beetle Management

The Aerial Overview Survey (AOS) is conducted annually over 80-90% of the province provides a way to track changes in various visible forest health agents. The graphs in Appendix 1 are based on AOS data from 1999-2015 showing the total area impacted by each bark beetle.

Follow guidance provided by [Best Practices for Managing Douglas-fir Beetle](#):

- Introduction and Biology
- Management Strategies and Tactics
- Landscape and Stand Level Planning
- Douglas-fir Beetle and Wildfire

[Bark beetle hazard maps](#) identify stands that are highly susceptible to bark beetle because of their stand characteristics.

[Detailed](#) and [overview](#) aerial survey data for 2018 is the most up to date information for infestation location.

Mule Deer Winter Range (MDWR) and Bark Beetles

Follow Government Actions Regulation (GAR) orders pertaining to ungulate winter range (#U-5-001, U-5-002, and U-5003):

- [Ungulate Winter Ranges Cariboo Chilcotin Land Use Plan, Shallow and Moderate Snowpack](#)
- [Ungulate Winter Ranges Cariboo Chilcotin Land Use Plan, Transition and Deep Snowpack](#)
 - Use existing access only.
 - Trees selected for trap trees should be ones that would be harvested in a regular MDWR entry
 - Trap trees should not be located on ridge lines or topographic breaks.
 - Trap trees should be mark-to-cut and fallers should be instructed clearly as to what the goals are and should be supervised appropriately.

- Trap tree sites must be accurately mapped with trees removed prior to next beetle flight without damage to the surrounding stand.
- Trap tree sites in MDWRs would be followed up with a second treatment (eg., trap trees, anti-aggregation pheromone baiting (MCH))
- All infested Douglas-fir trees, stumps, and debris associated with sanitation harvest must be treated and/or removed from the site prior to beetle flight.

Old Growth Management Areas (OGMAs) and Bark Beetles

In many areas, OGMAs contain the only representation of old forests in a landscape/BEC zone. The qualities present in the OGMAs (large, old trees) are also attractive for bark beetles. To minimize the impact on OGMAs the following precautions will be taken:

- Be consistent with the [Land Use Objectives for the Cariboo-Chilcotin Land Use Plan](#) and [Biodiversity Updates](#).
- Identify OGMAs of major concern for annual bark beetle monitoring
- Susceptible stands outside OGMAs will aid in prioritizing OGMA treatments
- Where infestations occur inside OGMAs, if appropriate, locate an intensive trap tree program outside the boundary to draw beetles out of the OGMA
- Consider using MCH on Douglas-fir where it could be of benefit to protect healthy stands inside OGMAs which may be provided and/or placed by FLNRORD
- All infested Douglas-fir trees, stumps, and debris associated with sanitation harvest must be treated and/or removed from the site prior to beetle flight.

8. Non-Recoverable Losses

Non-recoverable losses account for the average volume lost each year due to natural causes, such as pests, fire and wind, that are not recovered or salvaged. The WLTSR Rationale for AAC Determination assumed a total of 149,553 cubic metres per year. Since this time a new tool for estimating NRL has been developed using the provincial aerial overview survey data. The salvage areas are netted out of the mapped mortality resulting in a rough estimate of unsalvaged volumes. This spreadsheet is posted on the FTP site at [https://www.for.gov.bc.ca/ftp/HFP/external/!publish/Forest Health/NRLs/](https://www.for.gov.bc.ca/ftp/HFP/external/!publish/Forest_Health/NRLs/) The most recent NRL estimates for the WLTSR are provided in Table 4.

Table 4: Summary of estimated non-recoverable losses (unsalvaged) by damaging agent 2009-2019

Year	Douglas-fir beetle (m3/yr)	Drought (m3/yr)	Fire (m3/yr)	Flood (m3/yr)	Mountain pine beetle (m3/yr)	Spruce beetle (m3/yr)	Western balsam bark beetle (m3/yr)	Total (m3/yr)
2009	60,160	0	514,717	0	1,506,064	87,598	208	2,168,747
2010	4,416	0	557,308	124	135,053	122,556	9,74	819,540
2011	5,799	0	944	297	35,835	10,933	2,299	56,107
2012	25,238	0	6,888	6,136	30,346	85,204	9,813	163,625
2013	2,602	0	1,283	4,593	8,368	17,536	2,116	36,498
2014	34,531	36	7,266	6,812	5,746	1,950	3,426	59,768
2015	36,200	21,123	39,082	1,490	7,462	4,274	615	110,246
2016	121,996	7,915	1,332	1,821	20,615	18,884	5,647	178,210
2017	169,823	0	2,866,697	388	18,761	36,488	2,609	3,094,766
2018	91,476	1,076	173,832	3,698	19,483	1,447	8,151	299,163
2019	36,268	6,410	0	2,060	12,686	3,492	1,819	62,735

9. Reducing the Impacts of Climate Change

There is a growing body of evidence that global climate change has contributed to increased incidence and severity of several forest health factors (Woods et al., 2010). Therefore, adapting natural resource management to climate change is necessary to ensure forests are productive and ecosystems are resilient.

The [Cariboo Region Extension Note on Adapting Natural Resource Management to Climate Change](#) includes climate change projections; projected impacts of climate change to ecosystems; and adaptation strategies for natural resource management.

Website for Climate Change and Adaptation in B.C. Forests:

<https://www2.gov.bc.ca/gov/content/environment/natural-resource-stewardship/natural-resources-climate-change>

The choice of silviculture strategies and/or harvesting systems can be an important consideration in establishing a healthy future stand. These strategies need to be considered in the context of integrated resource management.

Some silvicultural strategies and harvesting systems are listed below to help minimize the impact from forest pests and mitigate the risks associated with climate change:

- Treat the site prior to planting. For obligate parasites that have short distance spread (dwarf mistletoes) or site specific pests (some root rots), site treatment such as sanitation spacing or stumping can be an effective method of reducing losses from disease.
- Planting within two years of slash burning may increase the risk of black army cut worm or Rhizina root disease in certain areas.
- Before considering spacing it is important to conduct pest surveys and to consider the impact of forest pests on final stocking levels. In some instances, it is preferable to delay spacing until the final impact of forest pests can be more accurately assessed.
- The Ministry of Forests Lands, Natural Resource Operations and Rural Development has modified their seed transfer standards to [climate based seed transfer](#) to mitigate the impacts of climate change.
- Partial harvest systems are generally not recommended in stands with root disease or dwarf mistletoe but may be beneficial for regenerating well stocked stands with greater structural and species diversity and may provide some mitigation against growing season frosts and other weather events.

10. Key TSR Issues

In February 2015 a new Annual Allowable Cut Determination for the WL TSA was published. In this document FLNRO staff, other agencies and licensees (as appropriate) are directed by the Chief Forester to undertake or support specific tasks and studies. The key tasks and studies that relate to this strategy are noted below. These projects are important to help reduce the risk and uncertainty associated with factors that affect the timber supply in the Williams Lake TSA.

Short-term considerations:

1. Licensees are to continue to focus harvesting as much as possible on mountain pine beetle-impacted pine-leading stands in the Williams Lake TSA; and to harvest no more than their share of the AAC partition attributable to live tree volume.
2. District staff are to monitor the following and report semi-annually to the chief forester:

- a. Harvest performance within MPB-killed pine-leading stands and the volume attributable to live trees within those stands; and
- b. Harvest contribution from non-pine leading stands.

Mid-term timber supply considerations:

1. *Low productivity sites*: Staff are to explore opportunities (eg., rehabilitation and bioenergy) for the use of some of the low productivity sites before the next determination
2. *Mule deer winter range*: Staff are to address MDWR issues including meeting minimum basal area targets relative to current or projected stand conditions; incentives for non-clearcut harvesting approaches given higher administrative and planning costs; and addressing poor forest health conditions that may be exacerbated by lack of management before the next determination so that this factor can be better considered. There is also a need for a standard Cariboo region-wide approach to modelling winter range requirements stemming from an order under the Government Actions Regulation.
3. *Douglas-fir stands*: Staff are to explore innovative ways (such as use of LiDAR) to improve the inventory, including growth and yield, of Interior Douglas-fir stands given their contribution to mid-term timber supply.

Other considerations:

1. *Climate change*: Staff are to try and understand projected climate change impacts in the TSA so that this important consideration can be factored into the next determination.

11. Web Links for Resources

Cariboo Region Forest Health Strategies and Data

Guide to identifying Douglas-fir beetle attack from ground assessments

https://www.for.gov.bc.ca/ftp/RSI/external/!publish/Forest%20Health/Forest%20Health%20Presentations/Douglas-fir_beetle_signs.pdf

2015 updated Seral Analysis. This analysis is based on the 2015 VRI data.

Report:

ftp://ftp.geobc.gov.bc.ca/publish/Regional/WilliamsLake/forest/seral/seral_2015/report/

Maps:

ftp://ftp.geobc.gov.bc.ca/publish/Regional/WilliamsLake/forest/seral/seral_2015/maps/

Root Rot shape files

<https://www.for.gov.bc.ca/ftp/RSI/external/!publish/Forest%20Health/Root%20Rot%20Shapefiles/>

Biodiversity Committee Updates

Cariboo-Chilcotin Land Use Plan. 2006. An Integrated Strategy for Management of Biodiversity and Bark Beetles in Douglas-fir and Spruce Stands. Update Note #7b.
https://www2.gov.bc.ca/assets/gov/farming-natural-resources-and-industry/natural-resource-use/land-water-use/crown-land/land-use-plans-and-objectives/cariboo-region/cariboochilcotin-rlup/update_note_7b.pdf

Land Use Order

Objectives for wildlife tree retention, OGMAs, Critical Fish Habitat, Lakes, Riparian Reserve Zones, Grasslands, High Value Moose Wetlands, and Grizzly Bear.

[ftp://ftp.geobc.gov.bc.ca/publish/Regional/WilliamsLake/Cariboo-Chilcotin LUOR Order/legal order document/CaribooChilcotinLUO_May2011.pdf](ftp://ftp.geobc.gov.bc.ca/publish/Regional/WilliamsLake/Cariboo-Chilcotin_LUOR_Order/legal_order_document/CaribooChilcotinLUO_May2011.pdf)

MDWR Management Strategies

[Management Strategy for Mule Deer Winter Ranges in the Cariboo-Chilcotin Part 1a:](#)

Management Plan for Shallow and Moderate Snowpack Zones. 2007. R.J. Dawson, H.M. Armleder, B. Bings, and D. Peel.

[Management Strategy for Mule Deer Winter Ranges in the Cariboo-Chilcotin Part 1b:](#)

Management Plan for Transition and Deep Snowpack Zones. 2006. R.J. Dawson, H.M. Armleder, B. Bings, and D. Peel.

General Wildlife Measures

Order for Shallow/Moderate Snowpack MDWRs:

http://www.env.gov.bc.ca/wld/documents/wha/Amendment_ShallowModerate_Feb07_Ord.pdf

Order for Transition/Deep Snowpack MDWRs:

http://www.env.gov.bc.ca/wld/documents/wha/Amendment_TransDeep_Feb07_ord.pdf

Individual MDWR Long-term Objectives Maps, Overview TSA MDWR maps, Snowpack Zones map:

<http://www.env.gov.bc.ca/esd/distdata/ecosystems/mdwr/>

Information Notes:

Mule Deer Winter Range Strategy Committee. 2014. Regional mule deer winter range strategy. Information Note #1. Guidance for fire damaged stands.

https://www2.gov.bc.ca/assets/gov/farming-natural-resources-and-industry/natural-resource-use/land-water-use/crown-land/land-use-plans-and-objectives/cariboo-region/cariboochilcotin-rlup/fire_damaged_stands.pdf

Mule Deer Winter Range Strategy Committee. 2014. Regional mule deer winter range strategy. Information Note #2. Guidance for MDWR General Wildlife Measure Exemption Requests for Salvage of Insect-killed Douglas-fir.

https://www2.gov.bc.ca/assets/gov/farming-natural-resources-and-industry/natural-resource-use/land-water-use/crown-land/land-use-plans-and-objectives/cariboo-region/cariboochilcotin-rlup/mule_deer_winter_range_wildfire_exemption_request.pdf

Frost Hazard Guide and Extension Note

Steen, O.A.S., R. Stathers, and R. Coupé. 1990. Identification and management of summer frost prone sites in the Cariboo Forest Region. For. Can. and B.C. Min. For., Victoria, B.C. FRDA Rep. 157. 23 pp.

<http://www.for.gov.bc.ca/hfd/pubs/Docs/Frr/Frr157.htm>

<https://www.for.gov.bc.ca/rsi/research/cextnotes/extnot05.pdf>

Ecora Resource Group Ltd. 2012. Drought risk and frost hazard mapping for the Williams Lake TSA. And Frost Hazard Workbook (Excel). (maps available upon request).

<https://www.for.gov.bc.ca/ftp/DCC/external/!publish/Frost%20Hazard/Report%20and%20tools/>

General Forest Health Information

Provincial Aerial Overview Survey

https://www.for.gov.bc.ca/ftp/HFP/external/!publish/Aerial_Overview/

RESULTS Incidence Maps

https://www.for.gov.bc.ca/ftp/HFP/external/!publish/Forest_Health/RESULTS%20incidence%20maps/

12. References

Eng, M. 2004 Forest Stewardship in the Context of Large-scale Salvage Operations: An Interpretation Paper, B.C. Ministry of Forests Forest Science Program Technical Report 019

Heineman, J.L. et al. 2010 Investigating the influence of climate, site, location, and treatment factors on damage to young lodgepole pine in southern British Columbia. Can J. For. Res. 40: 1109-1127

Maclauchlan, Lorraine E. & Brooks, Julie E. 2021. Comparison of Two Treatment Regimes for Managing Western Balsam Bark Beetle. *Journal of Ecosystems and Management* 21(2):1–11.


McLennan, D. 2003. Ecosystem restoration recommendations in areas affected by major salvage harvesting. British Columbia Ministry of Water, Land and Air Protection. Terrestrial Ecosystem Restoration Program, Victoria, B.C. Review draft.

Westfall J. & Brooks, J. 2001 Cariboo Forest Region Pests of Young Pine Stands: Incidence and Impact 1996-1999. B.C. Min. of For.


Woods, A.J. et al. 2010. Forest health and climate change: a British Columbia perspective. *Forestry Chronicle* 86(4): 412-417.

Appendix 1: Bark Beetle Population Trends 1999-2019



Area affected (ha) by Douglas-fir Beetle: 1999 – 2019




Area (ha) affected by Spruce Beetle: 1999 – 2019



Area (ha) affected by Mountain Pine Beetle: 1999 – 2019



Area (ha) affected by Western Balsam Bark Beetle: 1999 – 2019



Appendix 2:

**Beetle Management Units
and WLTSa Strategies**

Beetle Management Unit (BMU) Priority Areas

1. Talla/Little Eagle
2. Talla/Little Eagle
3. Westbranch
4. Eniyud Community Forest
5. Middle Lake
6. Westbranch
7. Cheshi Stikean

8. Clisboko
9. Policy
10. AFRF-Gavin Lake
11. Eniyud Community Forest - East
12. Big Lake
13. BCTS-Gavin Lake West
14. Proposed FNWL Anaham 2
15. BCTS-Tiran West
16. WLCF-Fat Rock East
17. Proposed FNWL Anaham 3
18. BCTS-Medium
19. BCTS-Tiran East
20. BCTS-Medium
21. BCTS-Mackin 1

22. ARM - Woodlot W0501
23. Chimney
24. Aikali East
25. ARM - Abbot
26. ARM - Johnny Stoks
27. ARM - Mayfield
28. Rate
29. ARM - Jimmy Dave's
30. ARM - Froggy Swamp
31. Aikali CF
32. Aikali CF
33. Aikali CF
34. ARM - Little Dog
35. Enterprise
36. RMC
37. BCTS-Bambrick-Big Creek-Gaspard
38. ARM - Gaspard
39. BCTS-Gaspard
40. BCTS-Bambrick-Big Creek-Gaspard
41. BCTS-Dash

42. Williams Lake East
43. WLCF-Fat Rock West
44. ARM - North Chimney
45. ARM - Pabo Creek
46. Chimney
47. ARM - Pelier
48. ARM - South Chimney
49. ARM - Tuckers
50. Aikali CF
51. ARM - Mitchell's Meadow
52. ARM - Desous Mountain
53. ARM - Pabo Lake
54. Aikali CF
55. ARM - Emerald Lake
56. ARM - Expansion
57. ARM - Sandy Harry

58. BCTS-Queenel Lake
59. Beaver Valley
60. WLCF-Potato Mountain
61. BCTS-Monimbo
62. AFRF - Knife Creek
63. ARM - Knife Creek
64. Chimney

65. BCTS-Queenel Lake
66. Beaver Valley
67. WLCF-Potato Mountain
68. BCTS-Monimbo
69. AFRF - Knife Creek
70. ARM - Knife Creek
71. Chimney

72. Williams Lake East
73. WLCF-Fat Rock West
74. ARM - North Chimney
75. ARM - Pabo Creek
76. Chimney
77. ARM - Pelier
78. ARM - South Chimney
79. ARM - Tuckers
80. Aikali CF
81. ARM - Mitchell's Meadow
82. ARM - Desous Mountain
83. ARM - Pabo Lake
84. Aikali CF
85. ARM - Emerald Lake
86. ARM - Expansion
87. ARM - Sandy Harry

88. BCTS-Queenel Lake
89. Beaver Valley
90. WLCF-Potato Mountain
91. BCTS-Monimbo
92. AFRF - Knife Creek
93. ARM - Knife Creek
94. Chimney

95. Williams Lake East
96. WLCF-Fat Rock West
97. ARM - North Chimney
98. ARM - Pabo Creek
99. Chimney
100. ARM - Pelier
101. ARM - South Chimney
102. ARM - Tuckers
103. Aikali CF
104. ARM - Mitchell's Meadow
105. ARM - Desous Mountain
106. ARM - Pabo Lake
107. Aikali CF
108. ARM - Emerald Lake
109. ARM - Expansion
110. ARM - Sandy Harry

111. BCTS-Queenel Lake
112. Beaver Valley
113. WLCF-Potato Mountain
114. BCTS-Monimbo
115. AFRF - Knife Creek
116. ARM - Knife Creek
117. Chimney

118. Williams Lake East
119. WLCF-Fat Rock West
120. ARM - North Chimney
121. ARM - Pabo Creek
122. Chimney
123. ARM - Pelier
124. ARM - South Chimney
125. ARM - Tuckers
126. Aikali CF
127. ARM - Mitchell's Meadow
128. ARM - Desous Mountain
129. ARM - Pabo Lake
130. Aikali CF
131. ARM - Emerald Lake
132. ARM - Expansion
133. ARM - Sandy Harry

134. BCTS-Queenel Lake
135. Beaver Valley
136. WLCF-Potato Mountain
137. BCTS-Monimbo
138. AFRF - Knife Creek
139. ARM - Knife Creek
140. Chimney

141. Williams Lake East
142. WLCF-Fat Rock West
143. ARM - North Chimney
144. ARM - Pabo Creek
145. Chimney
146. ARM - Pelier
147. ARM - South Chimney
148. ARM - Tuckers
149. Aikali CF
150. ARM - Mitchell's Meadow
151. ARM - Desous Mountain
152. ARM - Pabo Lake
153. Aikali CF
154. ARM - Emerald Lake
155. ARM - Expansion
156. ARM - Sandy Harry

157. BCTS-Queenel Lake
158. Beaver Valley
159. WLCF-Potato Mountain
160. BCTS-Monimbo
161. AFRF - Knife Creek
162. ARM - Knife Creek
163. Chimney

164. Williams Lake East
165. WLCF-Fat Rock West
166. ARM - North Chimney
167. ARM - Pabo Creek
168. Chimney
169. ARM - Pelier
170. ARM - South Chimney
171. ARM - Tuckers
172. Aikali CF
173. ARM - Mitchell's Meadow
174. ARM - Desous Mountain
175. ARM - Pabo Lake
176. Aikali CF
177. ARM - Emerald Lake
178. ARM - Expansion
179. ARM - Sandy Harry

180. BCTS-Queenel Lake
181. Beaver Valley
182. WLCF-Potato Mountain
183. BCTS-Monimbo
184. AFRF - Knife Creek
185. ARM - Knife Creek
186. Chimney

187. Williams Lake East
188. WLCF-Fat Rock West
189. ARM - North Chimney
190. ARM - Pabo Creek
191. Chimney
192. ARM - Pelier
193. ARM - South Chimney
194. ARM - Tuckers
195. Aikali CF
196. ARM - Mitchell's Meadow
197. ARM - Desous Mountain
198. ARM - Pabo Lake
199. Aikali CF
200. ARM - Emerald Lake
201. ARM - Expansion
202. ARM - Sandy Harry

203. BCTS-Queenel Lake
204. Beaver Valley
205. WLCF-Potato Mountain
206. BCTS-Monimbo
207. AFRF - Knife Creek
208. ARM - Knife Creek
209. Chimney

210. Williams Lake East
211. WLCF-Fat Rock West
212. ARM - North Chimney
213. ARM - Pabo Creek
214. Chimney
215. ARM - Pelier
216. ARM - South Chimney
217. ARM - Tuckers
218. Aikali CF
219. ARM - Mitchell's Meadow
220. ARM - Desous Mountain
221. ARM - Pabo Lake
222. Aikali CF
223. ARM - Emerald Lake
224. ARM - Expansion
225. ARM - Sandy Harry

226. BCTS-Queenel Lake
227. Beaver Valley
228. WLCF-Potato Mountain
229. BCTS-Monimbo
230. AFRF - Knife Creek
231. ARM - Knife Creek
232. Chimney

233. Williams Lake East
234. WLCF-Fat Rock West
235. ARM - North Chimney
236. ARM - Pabo Creek
237. Chimney
238. ARM - Pelier
239. ARM - South Chimney
240. ARM - Tuckers
241. Aikali CF
242. ARM - Mitchell's Meadow
243. ARM - Desous Mountain
244. ARM - Pabo Lake
245. Aikali CF
246. ARM - Emerald Lake
247. ARM - Expansion
248. ARM - Sandy Harry

249. BCTS-Queenel Lake
250. Beaver Valley
251. WLCF-Potato Mountain
252. BCTS-Monimbo
253. AFRF - Knife Creek
254. ARM - Knife Creek
255. Chimney

256. Williams Lake East
257. WLCF-Fat Rock West
258. ARM - North Chimney
259. ARM - Pabo Creek
260. Chimney
261. ARM - Pelier
262. ARM - South Chimney
263. ARM - Tuckers
264. Aikali CF
265. ARM - Mitchell's Meadow
266. ARM - Desous Mountain
267. ARM - Pabo Lake
268. Aikali CF
269. ARM - Emerald Lake
270. ARM - Expansion
271. ARM - Sandy Harry

272. BCTS-Queenel Lake
273. Beaver Valley
274. WLCF-Potato Mountain
275. BCTS-Monimbo
276. AFRF - Knife Creek
277. ARM - Knife Creek
278. Chimney

279. Williams Lake East
280. WLCF-Fat Rock West
281. ARM - North Chimney
282. ARM - Pabo Creek
283. Chimney
284. ARM - Pelier
285. ARM - South Chimney
286. ARM - Tuckers
287. Aikali CF
288. ARM - Mitchell's Meadow
289. ARM - Desous Mountain
290. ARM - Pabo Lake
291. Aikali CF
292. ARM - Emerald Lake
293. ARM - Expansion
294. ARM - Sandy Harry

295. BCTS-Queenel Lake
296. Beaver Valley
297. WLCF-Potato Mountain
298. BCTS-Monimbo
299. AFRF - Knife Creek
300. ARM - Knife Creek
301. Chimney

302. Williams Lake East
303. WLCF-Fat Rock West
304. ARM - North Chimney
305. ARM - Pabo Creek
306. Chimney
307. ARM - Pelier
308. ARM - South Chimney
309. ARM - Tuckers
310. Aikali CF
311. ARM - Mitchell's Meadow
312. ARM - Desous Mountain
313. ARM - Pabo Lake
314. Aikali CF
315. ARM - Emerald Lake
316. ARM - Expansion
317. ARM - Sandy Harry

318. BCTS-Queenel Lake
319. Beaver Valley
320. WLCF-Potato Mountain
321. BCTS-Monimbo
322. AFRF - Knife Creek
323. ARM - Knife Creek
324. Chimney

325. Williams Lake East
326. WLCF-Fat Rock West
327. ARM - North Chimney
328. ARM - Pabo Creek
329. Chimney
330. ARM - Pelier
331. ARM - South Chimney
332. ARM - Tuckers
333. Aikali CF
334. ARM - Mitchell's Meadow
335. ARM - Desous Mountain
336. ARM - Pabo Lake
337. Aikali CF
338. ARM - Emerald Lake
339. ARM - Expansion
340. ARM - Sandy Harry

341. BCTS-Queenel Lake
342. Beaver Valley
343. WLCF-Potato Mountain
344. BCTS-Monimbo
345. AFRF - Knife Creek
346. ARM - Knife Creek
347. Chimney

348. Williams Lake East
349. WLCF-Fat Rock West
350. ARM - North Chimney

Last updated: 2021-11-04
Updated by: prodger

36

Beetle Management Unit (BMU) Priority Areas




Mountain Pine Beetle (IBM) Strategy November 2021

Beetle Management Unit Strategy (IBM)

NOV2020IBM

Salvage



Ministry of
Forests, Lands, Natural
Resource Operations
and Rural Development

Last updated: 2021-11-04
Updated by: prodger

Document Path: W:\for\RS\IDCC\Projects\2021\IDCC_21_041_BMU_update\MXD\IDCC_BMU_update.aprx

The map displays the following parcels:

- 1. Tantalus Eagle Westborough
- 2. Enlud Community Forest - East
- 3. Cheshi Stikekan
- 4. Tantalus Eagle Westborough
- 5. Tantalus Eagle Westborough
- 6. Tantalus Eagle Westborough
- 7. Tantalus Eagle Westborough
- 8. Tantalus Eagle Westborough
- 9. Tantalus Eagle Westborough
- 10. Tantalus Eagle Westborough
- 11. Tantalus Eagle Westborough
- 12. Tantalus Eagle Westborough
- 13. Tantalus Eagle Westborough
- 14. ARM - Place Lake
- 15. Rieke
- 16. Aikil East
- 17. ARM - Pecos Mountain
- 18. ARM - Alton
- 19. ARM - Johnny Sticks
- 20. ARM - Mayfield
- 21. ARM - Jimmy Dave's
- 22. ARM - Snoddy Swamp
- 23. Aikil Churnam
- 24. ARM - Little Dog
- 25. Enlud
- 26. ARM - Gaspar
- 27. Proposed FNWL Anaham 3
- 28. BCTS - Gaspar
- 29. BCTS - Barilbrook-Big Creek-Gaspar
- 30. BCTS - Dash
- 31. BCTS - Dash
- 32. BCTS - Dash
- 33. BCTS - Dash
- 34. BCTS - Dash
- 35. BCTS - Dash
- 36. BCTS - Dash
- 37. BCTS - Dash
- 38. BCTS - Dash
- 39. BCTS - Dash
- 40. BCTS - Dash
- 41. BCTS - Dash
- 42. BCTS - Dash
- 43. BCTS - Dash
- 44. BCTS - Dash
- 45. BCTS - Dash
- 46. BCTS - Dash
- 47. BCTS - Dash
- 48. BCTS - Dash
- 49. BCTS - Dash
- 50. BCTS - Dash
- 51. BCTS - Dash
- 52. BCTS - Dash
- 53. BCTS - Dash
- 54. BCTS - Dash
- 55. BCTS - Dash
- 56. BCTS - Dash
- 57. BCTS - Dash
- 58. BCTS - Dash
- 59. BCTS - Dash
- 60. BCTS - Dash
- 61. BCTS - Dash
- 62. BCTS - Dash
- 63. BCTS - Dash
- 64. BCTS - Dash
- 65. BCTS - Dash
- 66. BCTS - Dash
- 67. BCTS - Dash
- 68. BCTS - Dash
- 69. BCTS - Dash
- 70. BCTS - Dash
- 71. BCTS - Dash
- 72. BCTS - Dash
- 73. BCTS - Dash
- 74. BCTS - Dash
- 75. BCTS - Dash
- 76. BCTS - Dash
- 77. BCTS - Dash
- 78. BCTS - Dash
- 79. BCTS - Dash
- 80. BCTS - Dash
- 81. BCTS - Dash
- 82. BCTS - Dash
- 83. BCTS - Dash
- 84. BCTS - Dash
- 85. BCTS - Dash
- 86. BCTS - Dash
- 87. BCTS - Dash
- 88. BCTS - Dash
- 89. BCTS - Dash
- 90. BCTS - Dash
- 91. BCTS - Dash
- 92. BCTS - Dash
- 93. BCTS - Dash
- 94. BCTS - Dash
- 95. BCTS - Dash
- 96. BCTS - Dash
- 97. BCTS - Dash
- 98. BCTS - Dash
- 99. BCTS - Dash
- 100. BCTS - Dash

Last updated: 2021-11-04
Updated by: prodger