

Climate-Based Seed Transfer

Greg O'Neill and Nicholas Ukrainetz

Research Branch, BC Ministry of Forests and Range
Kalamalka Forestry Centre, Vernon BC

October 2008




Overview


- Rationale
- Delineation methods
- Seed transfer systems
- Assisted migration

Rationale


- The genecology revolution (Climate models, GIS, new statistical methods) allows for improvements to seed transfer systems.




Geographic-based deployment zone for Sx PG Low




Geographic-based deployment zone for Sx PG Low



Geographic and climate-based deployment zone for PG Sx Low



Climate-based deployment zone for Sx PG Low




- **Rationale**



- The genecology revolution (Climate models, GIS, new statistical methods) allows for improvements to seed transfer systems.
- Assisted migration requires a climate-based seed transfer system.



Overview


- Rationale
- Delineation methods
- Seed transfer systems
- Assisted migration















Mean Annual Temp
 Mean Annual Precip
 Mean Cold Month Temp
 Continentality
 Degree Days > 5 °C
 Mean Summer Precip
 Annual Drought Index
 Summer Drought Index








Evaluation



Climatic transfer distance


Deployability





Climatic transfer distance


Deployability


MAT zone	Area (km2)
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
Average	












Overview


- Rationale
- Delineation methods
- **Seed transfer systems**
- Assisted migration


Seed Transfer Systems





Overview


- Rationale
- Delineation methods
- Seed transfer systems
- Assisted migration
















		Climate zone Plantation											
		1	2	3	4	5	6	7	8	.	.	.	56
Climate zone Seed source	1	√	√	√									
	2	√	√	√	√								
	3	√	√	√	√	√							
	4		√	√	√	√	√						
	5			√	√	√	√	√					
	6				√	√	√	√	√				
	7					√	√	√	√				
	8						√	√	√				
	.												
	.												
	.												
	56												√

√ = acceptable deployment


Progressive Seed Zones - 2 climate variables





Seed transfer variables

- Species
- SPZ
- SPU
- Class A/B/B+
- BEC zone
- Latitude
- Longitude
- Elevation
- Overlap zones
- QCI (Ss, Hw)
- > 1100 m (Yc)
- North of 56 (Pli)
- North of 52 and EK SPZ (Fdi)

- 
- Species
 - BEC variant
 - Latitude

- Conclusions regarding CBST
 - Climate transfer distance halved
 - Deployability doubled
 - Low cost
 - Rapid implementation
 - Progressive ST system combines merits of fixed and focal ST
 - Assisted migration easily accommodated
 - Operationally simple
- Next
 - BEC vs climate-based ST system
 - Number of zones to optimize adaptation vs deployability

Acknowledgements



Photo: Nick Wheeler

- **Data**
 - Doug Ashbee
 - Andreas Hamann
 - Barry Jaquish
- **Analysis**
 - Frank van Thienen
- **Discussions**
 - Sally Aitken
 - Andreas Hamann
 - Jerry Rehfeldt
 - Tongli Wang
 - Jack Woods
 - Alvin Yanchuk