



# Adapting forest and range management to climate change in the Northeast Region:

## Considerations for planners and practitioners

### 1. About this Series

Climate change will fundamentally change ecosystems in British Columbia. Therefore, adapting forest and range management to climate change is necessary to foster resilient<sup>1</sup> ecosystems that continue to provide the services, products and benefits society relies on.

This extension note is part of a series that uses current climate change research<sup>2</sup> to summarize, for each region, projected climate changes, impacts to ecosystems, and adaptation strategies. Where regional information is limited, information is drawn from provincial-scope research.

This extension note is intended to help planners, practitioners and decision-makers with planning under the *Forest and Range Practices Act*, assessing cumulative effects, developing climate change action plans, and monitoring the effectiveness of adaptation practices. Endnotes provide references and further sources of information.

### 2. Provincial Overview<sup>3</sup>

**Climate:** BC has become warmer and wetter over the last century. Winter has warmed the most. Extreme rainfall and dry conditions have increased and snowpacks have decreased. These warming trends will continue. By the end of this century, mean annual temperature in BC could be at least 1.7 to 4.6°C warmer than it was in the last few decades<sup>4</sup>. More winter precipitation will likely fall as rain rather than snow, resulting in lower snowpacks, earlier and more rapid snowmelt, and longer fire seasons.

**Regional differences:** Northern and southern regions of BC are expected to warm more than coastal BC and parts of central BC. Winter precipitation is expected to increase in all regions, but summer precipitation is expected to increase in northern BC and decrease in southern and coastal BC.

**Impacts: Ecosystems** will likely undergo both predictable and unpredictable ecological shifts. Climate envelopes (the climate associated with an ecosystem today) for subalpine and alpine areas will diminish in most locations while those for grasslands, shrub-steppe and dry forested ecosystems are expected to expand. In response, ecological communities will disassemble and reassemble—sometimes into novel combinations—as populations decline, move or adapt. Many species, including trees, will not be able to migrate quickly enough to keep pace with shifting climate. During this transition, ecosystems will be strongly influenced by disturbances and invasive plants. **Natural disturbance** dynamics will change: likely changes include increased fire and drought in southern and coastal BC, increased storms and windthrow on the coast, and more frequent and extensive mortality due to bark beetles, defoliators and

diseases across BC. Invasive species will increase. **Hydrological regimes** will shift due to increased evaporation, altered vegetation communities, increased storm frequency and magnitude, decreased snow accumulation, seasonal changes to precipitation, and accelerated ice melt followed by diminished glacier extent.

**Adaptation:** Many adaptation strategies are similar across BC. With the exception of assisted migration, most strategies are not novel, but are elements of ecosystem management that require broader application. Strategies to reduce risks to forest ecosystems include **promoting resilience** by maintaining or increasing diversity at all scales, **guiding ecological transformation** by maintaining landscape connectivity and assisting migration, **combating detrimental change** by controlling invasive plants and excessive disturbance, and **limiting cumulative effects** of multiple land-use activities. Strategies to reduce risks to forestry-dependent communities include **increasing monitoring** of change, **increasing capacity** to respond to change (e.g., by economic diversification), **strategically harvesting** at-risk forests, managing fire in wildland-urban **interfaces**, and increasing capacity of **infrastructure** to withstand extreme events.

### 3. Description of Region

The Northeast Region includes the vast Northeastern Plains edged by the Rocky Mountains to the west. The mountains act as a barrier to Pacific influence. Unlike the rest of BC, the Northeast is dominated by continental air masses (both polar and arctic), and therefore experiences drier conditions and a large spread in temperatures. Winters are long and cold, and the growing season is relatively short. Precipitation is low except at high elevations in the southwest of the region. The boundary between mountains and plains represents an enduring feature that will shape ecosystems in any climate regime. A less obvious and perhaps less enduring boundary divides the northern and southern portions of the region. For more information on weather systems for BC, visit [BC Weather](#)<sup>5</sup>.

Current ecoprovinces, Taiga Plains, Boreal Plains, Northern Boreal Mountains, and Sub-boreal Interior, represent a reasonable division of the Northeast region into four broad climatically-relevant portions (Figure 1). At this broad scale, current forested biogeoclimatic (BEC) zones<sup>6</sup> match the sections well: the **plains** sub-regions are dominated by BWBS with extensive muskeg, the **sub-boreal mountains** sub-region by ESSF, and the **northern boreal mountains** sub-region by SWB with BWBS in valleys. Alpine ecosystems are common in the mountains. Fires are extensive; hence, many forests are seral stands dominated by aspen. Non-forested scrub and boreal grassland occurs throughout the region, particularly in the north. For more information on BEC zones in this region, visit [BEC WEB](#).




**Figure 1.** Climate zones, based on ecoprovinces, within the Northeast Region. Spatial data from [DataBC](#).

## 4. Climate Change Projections

Our understanding of climate change is adjusting continually as models are improved with new research and methods. Projections in this document are derived from the Pacific Climate Impacts Consortium's regional climate summary for the [Northeast Region](#) and its [Plan2Adapt](#) tool for projecting future climate conditions.

The climate in the Northeast Region has changed more rapidly over the past century than the rest of the province and is expected to continue to change. Averaged across the region, over 2°C of warming has occurred during the 20<sup>th</sup> century, with most warming in winter; by the 2050s, winter is projected to warm an additional 2.4°C (0.7 – 3.6°C). Projections suggest the region may warm, on average, by about 1.8 to 4.6°C by the end of this century, similar to moving from Fort St. John to Williams Lake or Fort Nelson to Prince George (2.2° and 4.4°C warmer respectively).

Over the Northeast Region, annual precipitation has generally increased over the past century, but has decreased in winter from 1951 – 2009. Currently, most precipitation (about 210 mm) falls in summer, the majority as convective showers or thunderstorms. Changes in precipitation are projected to be modest relative to historic variability, with slight increases in all seasons but little change in summer. Snowfall may increase in winter. The large percentage change in spring (March, April, May) snowfall is only about 10% of the annual snowfall and approximately 5% of total annual precipitation.

## Summary of climate projections for the Northeast for the 2050s\*

| Climate variable           | Change in Northeast                                                                         | Variation within region                |
|----------------------------|---------------------------------------------------------------------------------------------|----------------------------------------|
| <b>Temperature</b>         |                                                                                             |                                        |
| Mean (°C)                  | +1.8 (1.5 to 2.8)                                                                           | Fairly consistent                      |
| Summer (°C)                | +1.4 (1 to 2.6)                                                                             | Fairly consistent                      |
| Winter (°C)                | +2.2 (0.7 to 3.4)                                                                           | Slightly higher on plains and in north |
| <b>Precipitation (%)</b>   |                                                                                             |                                        |
| Annual                     | +6 (0 to 16)                                                                                | Slightly higher in northern mountains  |
| Summer                     | +4 (-6 to 13)                                                                               | Higher in north                        |
| Winter                     | +11 (-6 to 22)                                                                              | Fairly consistent                      |
| <b>Snowfall (%)</b>        |                                                                                             |                                        |
| Winter                     | +7 (-7 to 19)                                                                               | Increase more in north                 |
| Spring                     | -57 (-69 to -23)                                                                            | Least decrease in northern mountains   |
| <b>Snowpack</b>            | Little change                                                                               | Little change in north <sup>7</sup>    |
| <b>Frost-free days</b>     | +16 (9 to 23)                                                                               | Greater increase in mountains          |
| <b>Growing Degree Days</b> | +226 (148 to 392)                                                                           | Greater increase in plains             |
| <b>Extreme weather</b>     | More heat waves, summer drought, wildfires, heavy precipitation and windstorms <sup>8</sup> |                                        |

\* Based on 1961-1990 baseline. Projected changes in temperature continue to increase past 2050. Source: PCIC [Plan2Adapt](#) tool. Projections are based on a combination of A2 and B1 emissions scenarios, where A2 represents roughly business as usual and B1 represents a more optimistic scenario with about ½ of emissions of business as usual. Median of 30 projections with range (in brackets) showing the 10<sup>th</sup> to 90<sup>th</sup> percentile of projected changes.<sup>9</sup>

**Increased Variability and Extremes:** Focusing on mean changes in temperature and precipitation can obscure important changes in climate variability and extremes. Climate projections are based on mean temperature and precipitation per season and do not reflect potentially large changes in variability. Even with constant variability, the frequency of extreme events will increase much more than a small change in mean temperature or precipitation would suggest.

## Climate Variability and Extreme Events

Seemingly small increases in mean values of climate variables can substantially increase the probability of an extreme event. For example, increasing the mean by one standard deviation can lead to a more than 10-fold increase in extremes so that a 1 in 100 year event (e.g., flood) can become a 1 in 10 year event.<sup>10</sup>

## 5. Impacts to Ecosystems

### Ecosystem Climate Envelopes

Climate envelopes describe the climatic conditions associated with currently mapped biogeoclimatic (BEC) subzone/variants<sup>11</sup>. These envelopes help ecologists and foresters integrate climate variables and visualise the potential extent and implications of climate change, but they **do not** predict what future ecosystems will look like for several reasons. First, ecosystems do not move as a unit; second, current climate projections are based on average climate values, ignoring the extreme events that can shape ecosystem structure and composition; third, climate envelopes do not capture site-scale shifts well. Nonetheless, projections can help estimate the relative stress that climate change poses to an ecosystem and its potential to recover to a new functional state.

Climate envelopes are projected to shift upslope and northward across BC, resulting in expansion of some existing lower valley and plateau ecosystems, decline of higher elevation ecosystems, and creation of novel ecosystem assemblages. By the 2050s, climate envelopes for current Northeast biogeoclimatic zones are predicted to shift about 60 – 160 m upward in elevation and 10 – 175 km northward across their range (Figure 2)<sup>12</sup>. High elevation BEC zones across the Northeast will likely experience the highest stress, with ESSF, SWB and BAFA all projected to lose about two-thirds of their current area by the 2050s. The ESSF envelope is projected to move northward into current SWB, and so may be less stressed than SWB. The BWBS climate envelope is projected to lose little current area and to expand slightly inland and upslope. An ICH climate envelope is projected to move into southern valleys of the region.




**Figure 2.** Current (BEC version 8) and projected future (2041-2070) BEC zone locations. Future map shows consensus of multiple projections. Maps retrieved from [ClimateBC](https://climatebc.ca/).

At a smaller site-level scale, the wettest and driest ecosystems may experience regime shifts. For example, some dry sites may shift from forest to scrub or grasslands. An increased frequency of large fire events, particularly late in the growing season when water tables are low and peats have dried, make organic substrate wetlands susceptible to changes in hydrology, plant community composition, and degradation of permafrost after a severe fire.

## Natural Disturbance

The most profound changes to BC's forests in this century will likely be the result of more frequent and severe natural disturbances<sup>13</sup>. Changes in **mean climatic conditions** support range expansion of forest pests that are currently limited by climate. Northeast ecosystems are likely to undergo shifts due to the combination of droughts and enhanced pest incidence. Stressed aspen would be more susceptible to mortality from agents such as aspen leaf-miner and forest tent caterpillar. A combination of increased windthrow occurrence and more frequent dry summers could lead to more frequent outbreaks of the spruce bark beetle under future climates in the northeast. The limited number of tree species in the region means ecosystems may be more sensitive to pest outbreaks than more diverse ecosystems.

Changes in **extreme weather events** lead to increased abiotic and biotic disturbances. Although the Northeast is less likely to experience severe heat waves than southern portions of the province, and although precipitation is projected to increase on average, periods of relative drought projected for some sub-regions will influence fire hazard and drought stress, making trees more susceptible to insects and disease. Fewer extreme cold events will occur to inhibit insect outbreaks.

The increased prevalence of disturbance will vary by climatic sub-region, elevation and ecosystem type.

| Disturbance                                 | Projected changes                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|---------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Fire and drought                            | <ul style="list-style-type: none"> <li>• <b>Changes in fire regime</b> are <b>uncertain</b> in this region. Depending on interactions between precipitation, temperature and growing season length, projections range from minor decrease to substantial increase.</li> <li>• <b>Reduced spring moisture</b> availability and <b>increased</b> occurrence of late season <b>drought</b> events will potentially <b>increase</b> in <b>length of fire season</b>.</li> <li>• Periods of <b>drought</b> may increase, leading to <b>reduced water supplies</b> for residents and resource based industries, and <b>decreased</b> forest <b>regeneration success</b> and <b>increased tree mortality</b> due to moisture stress.</li> </ul> |
| Wind and mechanical damage                  | <ul style="list-style-type: none"> <li>• <b>Damage</b> to trees from <b>ice and snow</b> may proliferate with increased storms. Uncertainty is high.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Hydrogeomorphic (flooding and mass wasting) | <ul style="list-style-type: none"> <li>• <b>Increased peak flows</b> and sediment delivery may <b>affect aquatic ecosystems</b> and fish habitat as well as <b>damage infrastructure</b>. Uncertainty is high.</li> <li>• <b>Increased</b> seasonal <b>soil moisture</b> due to snowmelt and rainfall may <b>increase mass movement</b> in susceptible terrain.</li> <li>• <b>Mountain permafrost degradation</b> may lead to <b>more</b> frequent <b>landslides</b>.</li> <li>• <b>Loss of snow and ice</b> at high elevation may <b>increase mass wasting</b> from increasing freeze-thaw of exposed rock in mountainous sub-regions.</li> </ul>                                                                                       |
| Insects and disease                         | <ul style="list-style-type: none"> <li>• <b>Spruce beetles</b> may <b>increase</b> and move from a two-year to one-year cycle, increasing the probability of an outbreak.</li> <li>• <b>Broadleaf defoliators</b> may <b>increase</b> and, combined with drought, increase the probability of regime shifts from deciduous parkland to grassland in BWBS ecosystems with limited moisture storage capacity.</li> <li>• <b>Willow</b> may <b>decline</b> as the introduced willow stem borer spreads northward.</li> </ul>                                                                                                                                                                                                                |

## Hydrology

The Northeast Region is expected to follow projected provincial trends in hydrology<sup>14</sup>. Increased winter temperatures, increased precipitation, and reduced snowfall in spring (i.e., more rain) will likely shift the hydrological regime from snowmelt-driven to hybrid rain/snow-driven, leading to more frequent rain-on-snow events and smaller spring snowpacks. These changes will affect peak flows, sediment loads, and channel stability, reduce summer low flows, and increase the duration of the low-flow period, with consequent increased risk to fish and infrastructure.

Competition for limited water resources may increase as seasonal flow decreases; water allocation challenges may increase with climate change.

Loss of vegetation through natural or anthropogenic disturbance, combined with climate change, creates cumulative effects. Cumulative effects may decrease the capacity of the landscape to buffer rainfall, increase streamflow flashiness, and increase sediment delivery and channel instability.

| Feature                 | Projected changes                                                                                                                                                                                                                                                                                                                                                                                        |
|-------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Hydrological regime     | <ul style="list-style-type: none"><li>• <b>Glaciers and permafrost will thaw.</b></li><li>• <b>Shift</b> from <b>snowmelt-driven</b> to <b>hybrid rain/snow-driven</b>.</li><li>• Rain-on-snow events may give <b>greater spring peak flows</b>.</li><li>• <b>Altered timing</b> of peak and low flows.</li></ul>                                                                                        |
| Peak flows              | <ul style="list-style-type: none"><li>• <b>Timing</b> of <b>spring peak discharge</b> will <b>change</b>.</li><li>• Shift to hybrid regime could lead to <b>more frequent high flows</b> during fall and spring.</li><li>• These flows could <b>disturb</b> the <b>streambed and spawning habitat</b>.</li></ul>                                                                                         |
| Spring recession        | <ul style="list-style-type: none"><li>• <b>Loss of snow-driven regimes</b> could <b>reduce</b> the <b>spring recession</b> (a relatively stable period of moderate flow and temperature), affecting the sorting and deposition of sediment and profoundly impacting aquatic habitat.</li></ul>                                                                                                           |
| Low flows <sup>15</sup> | <ul style="list-style-type: none"><li>• <b>Smaller spring snowpack</b> and <b>earlier spring freshet</b> will <b>lower</b> summer <b>low flows</b>, <b>lengthen</b> the low-flow <b>period</b>, and <b>decrease groundwater storage</b>.</li><li>• <b>Temperature</b> may <b>increase</b> in some streams and lakes, posing <b>risk to temperature sensitive fish</b> such as Arctic grayling.</li></ul> |
| Variability             | <ul style="list-style-type: none"><li>• <b>Variability</b> in <b>peak flow</b> will likely increase, leading to <b>unstable stream morphology</b>.</li></ul>                                                                                                                                                                                                                                             |

### Flood Return Intervals

Small changes in mean climate can cause large changes in flooding frequency and magnitude. For example, current 50-year floods may become 5-year events and future 50-year floods might be what we now consider a 1,000-year flood.<sup>16</sup>

## Biodiversity

Climate shapes species distributions and ecological communities<sup>17</sup>. Populations faced with a changed environment can die out, move, or adapt to the new conditions. Many species that are adapted to projected future climates currently live several hundred kilometres distant from that area; only the fastest dispersers will be able to keep up with the pace of change. Most invasive plants and generalist weedy species are well-adapted for broad movement. For some ecosystems, potentially irreversible regime shifts may follow intense disturbances, particularly if invasive plants colonise and block historical successional paths.

Mountain ranges are particularly important for conservation of biodiversity. Relative to gentle terrain, mountains accommodate more climatic zones within close proximity; thus, as the climate changes, populations in lower elevation zones may find suitable climatic conditions by migrating upwards.

| Feature                | Projected changes                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cumulative effects     | <ul style="list-style-type: none"> <li>Past human activities, particularly development for shale gas, have altered, degraded and <b>fragmented habitat</b> in large areas of the northeast, making <b>dispersal</b> in response to climate-related disturbance more <b>difficult</b> for <b>specialist species</b>.</li> <li>Human response to increased disturbance (e.g. extensive <b>salvage harvesting</b>) can <b>exacerbate impacts</b> of climate change.</li> <li><b>Old forest</b> will <b>decline</b> due to disturbance and harvesting, threatening associated species.</li> <li><b>Over-grazing</b> and <b>trampling impacts forest understory</b> by changing the plant community, destroying the biological crust and opening soil to invasive plants.</li> </ul>                                                                                                                        |
| Ecological communities | <ul style="list-style-type: none"> <li><b>Communities</b> will <b>reassemble</b>, often into new combinations, as some established species decline or disappear, new species colonise and interactions change.</li> <li>Some ecosystems may undergo <b>regime shifts</b> (e.g. from <b>aspen parkland to scrub or grassland</b>).</li> <li><b>Below-ground communities</b> that support terrestrial vegetation may <b>change</b>.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Interactions           | <ul style="list-style-type: none"> <li><b>Ecological processes</b> and <b>relationships</b> among species (e.g. predation, pollination) may <b>uncouple</b> as the timing of events changes and becomes more variable (e.g., if migration depends on day length, but prey abundance depends on temperature, nesting time may be uncoupled).</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Invasive species       | <ul style="list-style-type: none"> <li><b>Invasive species</b> (plants and other organisms) are expected to <b>increase</b> as temperatures and disturbance increase.</li> <li>Current <b>indigenous species</b> may be <b>less competitive</b> in the new climate and disturbance conditions, facilitating invasive plant population expansion.</li> <li>Some locations in the Northeast already have invasive plant species capable of displacing native plant communities<sup>18</sup>.</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                  |
| Wildlife <sup>19</sup> | <ul style="list-style-type: none"> <li>Wildlife and trophic interactions (e.g., <b>predation</b>) will be particularly <b>affected by</b> changes in <b>snowpack</b> and <b>freeze-thaw regimes</b>. Impacts vary by species.</li> <li><b>Biting insects</b> and <b>disease</b> may <b>increase</b> in response to milder weather; for example, <b>winter ticks</b> may increase impacts to <b>moose</b>.</li> <li>In the northern sub-regions, <b>changed green-up timing</b> could <b>increase mortality</b> of young <b>sheep</b> and <b>goats</b>.</li> <li><b>Increased fire</b> could affect <b>caribou</b> habitat <b>negatively</b> and <b>moose</b> habitat <b>positively</b>.</li> <li><b>Greater plant productivity</b> in shallow boreal <b>lakes</b> will increase oxygen demand during the ice cover period, which may result in more frequent <b>large-scale fish kills</b>.</li> </ul> |

## Ecological Surprises

Current vulnerability modelling does not include ecological surprises or complex climate-ecological relationships. Mountain lions have established permanent, successfully reproducing populations in the Peace River area. Prior to 1990, only transient individuals were recorded. The combination of milder winters and high ungulate populations likely enabled this change. Simplistic predictions in complex systems cannot replace long-term interdisciplinary research and monitoring.

## Trees

Tree species distributions will shift gradually in response to climate change due to physiological tolerances, natural disturbance, and competition<sup>20</sup>. There are few tree species in the Northeast, hence low ecological redundancy. Most tree species will be unable to migrate quickly enough to follow the climate envelopes to which they are adapted. Uncertainty about climate projections leads to uncertainty about which trees may be best-suited to changing conditions. For example, warmer, wetter projections favour different species than warmer, drier projections<sup>21</sup>. Suitable trees at any point in time may be maladapted by rotation age, creating additional uncertainty and complexity for management.

| Response class          | Projected changes                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Physiological tolerance | <ul style="list-style-type: none"><li>• Most tree species in the Northeast seem physiologically resilient to projected conditions for the next 30 years but, by 2080, many <b>species within the BWBS</b> will be under moderate to high <b>drought risk</b><sup>22</sup>.</li><li>• <b>Interior spruce on dry sites</b> may be vulnerable to <b>moisture stress</b>.</li></ul>                                                                                                                                                                                        |
| Productivity            | <ul style="list-style-type: none"><li>• <b>Tree growth</b> will likely <b>increase</b> in some areas due to elevated CO<sub>2</sub> coupled with warmer temperatures.</li><li>• <b>Growth potential</b>, however, may <b>not be realised</b> because of <b>limited moisture</b> or nutrients, because populations are not adapted to changed seasonality and increased <b>extreme events</b>, or because <b>maladaptation</b> increases susceptibility to insects and disease. Provenances are adapted to even narrower climatic conditions than species.</li></ul>    |
| Natural disturbance     | <ul style="list-style-type: none"><li>• Episodic <b>die back</b> and <b>mortality</b> of trees will occur after <b>drought</b> events.</li><li>• <b>Stressed trees</b> are more <b>susceptible to insects and pathogens</b>.</li><li>• Several <b>deciduous</b> and <b>coniferous</b> species, including the aspen, willow and interior spruce that have defined ecosystems of the Northeast, may <b>have increased mortality</b> and highly <b>variable regeneration</b> due to a variety of factors including physiological stress, pathogens and insects.</li></ul> |
| Competition             | <ul style="list-style-type: none"><li>• Competitive interactions among trees and other plants will change in unexpected ways, leading to <b>shifts in species dominance</b>; for example, after disturbance on steeper warm slopes, <b>shrubs and grass</b> may be <b>favoured</b> over tree regeneration.</li></ul>                                                                                                                                                                                                                                                   |

## Range

Rangeland ecosystems will also shift in response to climate change due to higher temperatures and increased occurrence of droughts and floods. Earlier spring plant growth and a longer growing season may translate into a longer grazing season for livestock. Degradation of rangeland is most likely to occur during extreme events (e.g., soil erosion during intense rainstorms). Invasive species are most likely to establish in locations that were overgrazed during droughts. The most rapid effects of climate change

are likely to occur at high elevations and northern latitudes, which have a limited historical record from which to identify trends<sup>23</sup>.

### **Cumulative Effects**

Cumulative effects are changes to an ecosystem over time caused by a combination of **human activities, natural variability** and **climate change**. Assessment of cumulative effects integrates the effects of past, present and foreseeable future events and processes. FLNR has a [framework in place to guide assessment of the cumulative effects of resource management and climate change](#). Such a framework provides the context for informed decisions about which management actions are most likely to succeed.

## **6. Adaptation—modifying management to account for climate change**

It is necessary to modify management activities – planning, practices, and monitoring - to address the impacts of climate change on ecosystems. Adaptation strategies will vary depending on the ecosystem, the direction of anticipated climate change, the degree of certainty in projected changes, the urgency (risk and vulnerability), and the likelihood of adaptation practices achieving desired outcomes. Hence, management activities under a changing climate will need to be flexible and proactive.

This section includes adaptation strategies to address the current and anticipated impacts to ecosystems described above. These strategies reflect regionally-important ‘best available information’ drawn from research and the input of regional specialists. Practitioners and decision-makers should consider these adaptation strategies as voluntary guidance to inform operational planning and practices.

Many of these adaptation strategies are existing elements of good forest management that require broader application. As with current policy guidance, some adaptation strategies may conflict with each other (e.g., maintaining downed wood to sustain biodiversity may conflict with minimizing forest fuels to reduce catastrophic wildfire risks), requiring decisions that trade-off resource values depending on the priorities for the area in question. Trade-offs will likely become more common and challenging over time.

Although some of these adaptation strategies may incur incremental costs or land base constraints, the long-term economic benefits of adaptation to the productivity of timber, forage and other resource values are predicted to outweigh short-term costs. For example, studies in the Alex Fraser Research Forest, and in the Merritt and Kamloops TSAs, indicate the economic viability of diversifying managed forests to reduce forest health risks and increase resilience<sup>24</sup>. And, designing and maintaining roads and bridges to a higher standard will likely minimize repair and compensation costs after flood events. Some adaptation strategies are also potential climate change mitigation or carbon storage strategies (e.g., networks of reserves, retaining downed wood).

Adaptation strategies in this section reference related policy guidance and policy or knowledge needs.

### **Planning**

Climate change poses at least three broad challenges for planners:

- Existing management **objectives** may be inappropriate because they were developed without considering climate change and do not generally include objectives for mitigation or resilience.
- Existing management **strategies** are unlikely to achieve existing objectives under a changing climate.
- **Uncertainty** about the effectiveness of management strategies will increase.

In addition, regulatory or administrative change may be too slow to prompt timely management responses to changing conditions. At a broader scale, market forces may pose barriers.

## Uncertainty<sup>25</sup>

The impacts of climate change are already present on the landscape. However, projecting the impacts of climate change into the future is fraught with uncertainty due to incomplete ecological and climate models, and to alternative plausible emissions scenarios. Ecological processes that reflect multiple interactions (e.g. shifting species distributions) are more uncertain than processes that correlate strongly with a single variable (e.g. fire hazard and temperature).

Managing in the face of uncertainty requires:

- Recognition of uncertainty (known and unknown sources)
- Information gathering (via monitoring) to reduce uncertainty where possible
- Recognition that uncertainty increases with time span considered
- Acceptance that uncertainty will remain and a decision to either use precaution to maintain a desired value or to put a value at risk

Adaptation requires planning that includes new objectives, new strategies and increased consideration of uncertainty. For example, objectives to maintain biodiversity or timber could be modified to maintain ecological resilience, and strategies to mitigate flood impacts should take into account the higher probability of flooding associated with climate change plus climate oscillations. Planners may need more flexibility to handle regime shifts (e.g., if forested ecosystems lose viability). Best management practices for ecosystem management <sup>26</sup> provide an excellent resource.

## Practices

### Hydrology

To protect aquatic ecosystems and infrastructure near watercourses, adaptation consists of managing water removal to ensure sufficient summer flow in areas with drought, limiting sediment input (from surface erosion, streambank collapse and landslides), limiting increases in peak flows, and limiting increases in stream temperature.

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | Existing policy guidance and policy or knowledge needs                                                                                                                                                                                                                                                                                                                                                                                     |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Decreased summer flow</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <ul style="list-style-type: none"> <li>• Manage water allocations and use to maintain water supply during low flow periods</li> <li>• Increase storage reservoir capacity to meet demand in low flow periods</li> <li>• Evaluate aquifers and other new potential water sources</li> <li>• Retain sufficient riparian cover to maintain stream flow</li> <li>• Manage level of livestock grazing</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                        | <ul style="list-style-type: none"> <li>• Drought guidance<sup>27</sup></li> <li>• <b>Climate-based policy guidance for water allocation/use</b></li> <li>• <b>Updated riparian retention guidance for small streams</b></li> <li>• <a href="#">Water quality and livestock grazing BMPs</a></li> </ul>                                                                                                                                     |
| <b>Increased risk of landslides and surface erosion (that affect streams or infrastructure)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <ul style="list-style-type: none"> <li>• Avoid locating roads and cutblocks on or above unstable terrain</li> <li>• Design roads and drainage structures to accommodate increased peak flow and sediment transport in areas likely to become wetter: e.g., improve surface on high hazard roads; seed erodible cut slopes; build adequate ditches; replace selected culverts with bridges; limit road density in erosion-prone areas</li> <li>• Manage for permafrost; constructing on permafrost may lead to thawing and subsequent instability or erosion</li> <li>• Manage grazing to maintain functional riparian ecosystems; maintain sufficient riparian vegetation to control grazing</li> </ul>                                                                                                                            | <ul style="list-style-type: none"> <li>• <a href="#">Interior watershed assessment</a></li> <li>• Water quality evaluation<sup>28</sup></li> <li>• <b>Inventory “flashy” watersheds; monitor precipitation and runoff</b></li> <li>• <a href="#">Forest road engineering guidebook</a></li> <li>• <a href="#">Guidelines for managing terrain stability</a></li> <li>• <a href="#">Water quality and livestock grazing BMPs</a></li> </ul> |
| <b>Increased peak flows</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <ul style="list-style-type: none"> <li>• Limit Equivalent Clearcut Area (ECA) to 30 to 50% of THLB in sensitive watersheds</li> <li>• Anticipate increased natural disturbance and manage harvest to stay within ECA limits <ul style="list-style-type: none"> <li>○ Evaluate hydrological implications of salvage harvest</li> <li>○ Account for increased runoff from burned sites in ECA calculations</li> </ul> </li> <li>• Assess flood risk and increase design criteria for infrastructure</li> <li>• Protect stream sources from timber harvesting and compaction by livestock and machinery</li> <li>• Leave live standing vegetation and downed wood barriers next to all riparian areas on rangeland.</li> <li>• Limit development on known floodplains</li> <li>• Manage roads to minimize impacts to flows</li> </ul> | <ul style="list-style-type: none"> <li>• <a href="#">Interior watershed assessment</a></li> <li>• Post disturbance watershed effects<sup>29</sup></li> <li>• <a href="#">Practice guidelines for flood assessment</a></li> <li>• <b>Road and bridge design guidance to address increased peak flows</b></li> <li>• <a href="#">Water quality and livestock grazing BMPs</a></li> </ul>                                                     |
| <b>Increased stream temperature</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <ul style="list-style-type: none"> <li>• Retain adequate riparian vegetation next to streams and wetlands <ul style="list-style-type: none"> <li>○ Particularly important in temperature sensitive watersheds and along headwater areas</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     | <ul style="list-style-type: none"> <li>• <b>Inventory and monitor temperature sensitive watersheds<sup>30</sup></b></li> </ul>                                                                                                                                                                                                                                                                                                             |

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                   | Existing policy guidance and policy or knowledge needs                                                                 |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>Maintain ditches and culverts and deactivate roads to restore natural drainage as soon as possible <ul style="list-style-type: none"> <li>Important to prevent water warming on sites that feed streams, particularly in temperature sensitive watersheds</li> </ul> </li> </ul> | <ul style="list-style-type: none"> <li><b>Monitor road drainage</b></li> </ul>                                         |
| <ul style="list-style-type: none"> <li>Avoid harvesting sites with high water tables <ul style="list-style-type: none"> <li>Important for sites with high water tables that feed streams, particularly in temperature sensitive watersheds</li> </ul> </li> </ul>                                                       | <ul style="list-style-type: none"> <li><b>Inventory potential warm water sources (e.g., standing water)</b></li> </ul> |

## Biodiversity

Adaptation strategies for biodiversity are designed to achieve two objectives: (1) reduce the existing anthropogenic pressures that compound the negative effects of climate change on biodiversity (e.g., reduce harvesting and road access where sensitive values exist), and (2) promote resilient ecosystems at stand and landscape scales.

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                                                                                                                                                                     | Existing policy guidance and policy or knowledge needs                                                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Loss of old forest habitat and connectivity, due to increased tree mortality</b>                                                                                                                                                                                                                                                                                                                                                                                       |                                                                                                                                                                                                                  |
| <ul style="list-style-type: none"> <li>Create a network of reserves and corridors at multiple scales <ul style="list-style-type: none"> <li>Include riparian areas, wildlife tree patches, and old growth management areas in reserves</li> <li>Include corridors crossing elevation gradients</li> <li>Include habitat for specialized species and communities at risk</li> </ul> </li> </ul>                                                                            | <ul style="list-style-type: none"> <li><a href="#">Biodiversity Guidebook</a></li> <li><b>Updated land use plans</b></li> <li><b>Climate-based timber supply analysis</b></li> </ul>                             |
| <ul style="list-style-type: none"> <li>Limit salvage in reserve network (e.g., partial cut or avoid harvest) <ul style="list-style-type: none"> <li>Particularly important where stands buffer microclimate or provide large structure</li> </ul> </li> </ul>                                                                                                                                                                                                             | <ul style="list-style-type: none"> <li><a href="#">Chief Forester's retention guidance</a></li> <li>Post-disturbance biodiversity management<sup>31</sup></li> </ul>                                             |
| <b>Loss of suitable microclimate and soil conditions to re-establish historic ecosystems following harvest (e.g., potential regime shift from forest to grassland)</b>                                                                                                                                                                                                                                                                                                    |                                                                                                                                                                                                                  |
| <ul style="list-style-type: none"> <li>Avoid harvesting low elevation, very well drained upland sites prone to greater moisture stress <ul style="list-style-type: none"> <li>Use silviculture practices that conserve snow pack and soil moisture</li> </ul> </li> <li>Partially-cut stands on dry sites (i.e., retain partial overstory shelter)</li> <li>Plant spruce under mature aspen to establish advanced regeneration 20 years prior to aspen harvest</li> </ul> | <ul style="list-style-type: none"> <li><a href="#">Drought risk assessment tool</a></li> <li><a href="#">Enhancing biodiversity through partial cutting</a></li> <li><b>Inventory sensitive sites</b></li> </ul> |
| <ul style="list-style-type: none"> <li>Retain large downed wood <ul style="list-style-type: none"> <li>Particularly important on drier sites and in riparian ecosystems</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                            | <ul style="list-style-type: none"> <li><a href="#">Wildlife trees and coarse woody debris policies</a></li> <li><a href="#">FREP CWD backgrounder</a></li> <li><a href="#">CWD management</a></li> </ul>         |
| <ul style="list-style-type: none"> <li>Promote rapid site recovery to appropriate species (e.g., reforestation of dry sites; retain deciduous trees on moist sites) <ul style="list-style-type: none"> <li>Particularly important on drier sites</li> </ul> </li> </ul>                                                                                                                                                                                                   | <ul style="list-style-type: none"> <li>See Trees section</li> </ul>                                                                                                                                              |

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                                                                                                                                                                                                                    | Existing policy guidance and <i>policy or knowledge needs</i>                                                                                                                                                                                                                                                                                       |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Loss of vigour and diversity in young and maturing forests due to maladaptation from changing climate</b>                                                                                                                                                                                                                                                                                                                                                                                                             |                                                                                                                                                                                                                                                                                                                                                     |
| <ul style="list-style-type: none"> <li>Plant climatically-suited species and genotypes (i.e., facilitate migration)</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                           | <ul style="list-style-type: none"> <li>See Trees section</li> </ul>                                                                                                                                                                                                                                                                                 |
| <ul style="list-style-type: none"> <li>Retain naturally-occurring and regenerating species (including deciduous trees and shrubs) and plant a diverse species mix</li> </ul>                                                                                                                                                                                                                                                                                                                                             | <ul style="list-style-type: none"> <li><a href="#">Climate change stocking standards</a><sup>32</sup></li> </ul>                                                                                                                                                                                                                                    |
| <ul style="list-style-type: none"> <li>Use stand tending to influence successional pathways</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                   | <ul style="list-style-type: none"> <li>See Trees section</li> </ul>                                                                                                                                                                                                                                                                                 |
| <b>Increased spread of invasive plants following disturbance</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                                                                                                                                                                                                                                                                                                                                                     |
| <ul style="list-style-type: none"> <li>Minimize roads <ul style="list-style-type: none"> <li>Especially important in currently unroaded areas and susceptible ecosystems</li> </ul> </li> <li>Minimize road use (e.g., gates, deactivation)</li> <li>Establish competitive vegetation in ditches, on side slopes and other disturbed soil as soon as possible</li> <li>Wash equipment to remove seeds and plants prior to moving into new areas</li> <li>Follow best management practices for invasive plants</li> </ul> | <ul style="list-style-type: none"> <li><a href="#">Invasive plant management practices</a></li> <li><a href="#">Invasive species council of BC</a></li> <li><a href="#">Invasive species working group; IAPP Map, E-Flora BC</a></li> <li><b>Access management policy</b></li> <li><b>Increase monitoring and controlled experiments</b></li> </ul> |
| <ul style="list-style-type: none"> <li>Manage grazing to maintain vigorous native plant communities <ul style="list-style-type: none"> <li>Adjust stocking rates and distribution</li> <li>Integrate invasive plant management with overall rangeland resource management</li> </ul> </li> </ul>                                                                                                                                                                                                                         | <ul style="list-style-type: none"> <li><a href="#">Managing rangeland invasive plants</a></li> <li><b>Updated range policy</b></li> </ul>                                                                                                                                                                                                           |
| <ul style="list-style-type: none"> <li>Minimize site disturbance, especially multiple disturbances <ul style="list-style-type: none"> <li>Particularly important on susceptible sites</li> <li>Ensure revegetation seed mixtures are free of invasive plants</li> </ul> </li> <li>Account for invasive plants in site plans</li> </ul>                                                                                                                                                                                   | <ul style="list-style-type: none"> <li><b>Map distribution of invasive plants</b></li> </ul>                                                                                                                                                                                                                                                        |

## ***Trees***

Adaptation strategies for trees are designed to increase establishment success, survival and growth potential, and to reduce the negative impacts of natural disturbance resulting from climate change. Adaptation strategies have the potential to shift overall climate-induced impacts on timber supply from negative to positive or neutral.

Plantings of pine and spruce from genetic stocks which are better suited for predicted future climates are likely to have the most widespread potential application in the northeast over the next 30 years. Interior Douglas-fir may have application on warmer aspects under suitable silviculture regimes. Planting of white spruce under mature aspen to establish spruce advance regeneration at least 20 years prior to aspen harvest would retain boreal mixed wood ecosystems on the landscape.

Adaptation may have limited success in reducing fire disturbance because effective fire control in average-weather years can be negated by large fires in very dry years. Landscape fire management planning is aimed at reducing these losses as much as possible.

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | Existing policy guidance and <i>policy or knowledge needs</i>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Increased tree growth potential on sites with sufficient moisture</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <ul style="list-style-type: none"> <li>Plant climatically-suited species and genetic stock <ul style="list-style-type: none"> <li>Especially on dry sites or sites facing drought</li> </ul> </li> <li>Establish operational trials to test survival and growth of interior Douglas fir and western larch</li> <li>Plant white spruce under maturing aspen at least 20 years prior to aspen harvest to establish spruce advance regeneration</li> </ul>                                                                                                                                                            | <ul style="list-style-type: none"> <li><a href="#">Tree species selection tool</a></li> <li><a href="#">FFT assisted species migration guidance</a></li> <li><a href="#">Chief Forester standards for seed use</a><sup>33</sup></li> <li><a href="#">Climate-based seed transfer interim policy measures</a></li> <li><a href="#">Seed zone maps</a></li> <li><b>Measures to address cost and risk</b></li> </ul>                                                                                                                                                                                 |
| <ul style="list-style-type: none"> <li>Partially cut stands on dry sites to retain shelter and soil moisture, to conserve snow pack and to increase fire resiliency</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                     | <ul style="list-style-type: none"> <li><a href="#">Drought risk assessment tool</a></li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Increased disease-related mortality (mainly younger stands)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <ul style="list-style-type: none"> <li>Plant climatically suited species and genetic stock <ul style="list-style-type: none"> <li>Especially on sites facing drought and areas shifting to wetter climate</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                                                                                                                                   | <ul style="list-style-type: none"> <li><a href="#">Forest health and species selection</a></li> <li><a href="#">Regional &amp; TSA forest health strategies</a></li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <ul style="list-style-type: none"> <li>Increase within-stand genetic diversity by planting several genotypes for a given species</li> <li>Increase stand-scale species diversity (e.g., retain and plant a variety of species, including broadleaf; expand breadth of “acceptable” species in young stands)</li> <li>Actively manage for mixed species stands by applying knowledge of mixed wood stand development and future desired forest products</li> <li>Increase landscape-scale species diversity by planning retention and reforestation at the landscape level; vary species mix and density</li> </ul> | <ul style="list-style-type: none"> <li><a href="#">Long-term forest health and stocking standards</a></li> <li><a href="#">Guidance on species composition</a></li> <li><a href="#">Guidance on broadleaves</a></li> <li><a href="#">Guidance for FSP stocking standards</a></li> <li><a href="#">Mixed species options for FFT</a></li> <li>Successional responses<sup>34</sup></li> <li><a href="#">Stocking standards reference guide</a></li> <li><a href="#">Climate change stocking standards</a></li> <li><b>Free-growing policy that allows variation for local conditions</b></li> </ul> |
| <ul style="list-style-type: none"> <li>Plant higher initial densities to account for losses to biotic (and abiotic) damage; i.e. build in redundancy</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                    | <ul style="list-style-type: none"> <li><a href="#">Regional &amp; TSA forest health strategies</a></li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <ul style="list-style-type: none"> <li>Minimize mechanical damage from wind, snow and ice</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | <ul style="list-style-type: none"> <li><a href="#">BCTS windthrow manual</a></li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Increased beetle-related mortality (mainly mature/old stands)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| <ul style="list-style-type: none"> <li>Plant climatically-suited species and stock</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | <ul style="list-style-type: none"> <li>See above</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <ul style="list-style-type: none"> <li>Increase stand-scale diversity</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   | <ul style="list-style-type: none"> <li>See above</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <ul style="list-style-type: none"> <li>Shorten rotations <ul style="list-style-type: none"> <li>Especially for relatively productive sites most susceptible to disturbance</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                  | <ul style="list-style-type: none"> <li><b>Benefit/cost analysis that considers timber supply</b></li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <ul style="list-style-type: none"> <li>Monitor and control beetle population sources (e.g., sanitation harvesting) <ul style="list-style-type: none"> <li>Focus on stands where benefit of control outweighs cost to non-timber values</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                                                                                                      | <ul style="list-style-type: none"> <li><a href="#">Regional &amp; TSA forest health strategies</a></li> <li><a href="#">Provincial bark beetle management strategy</a></li> <li><a href="#">Chief Forester’s retention guidance</a></li> </ul>                                                                                                                                                                                                                                                                                                                                                    |
| <b>Increased fire hazard (all stand ages)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |

| Projected ecosystem changes and adaptation strategies                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      | Existing policy guidance and <i>policy or knowledge needs</i>                                                                                                                                                                                                                           |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• Increase fire resilience at the landscape level by creating strategic fuel breaks, prescribing fire, and allowing ecologically appropriate fires in suitable locations to burn under appropriate conditions <ul style="list-style-type: none"> <li>○ Work with range managers to create fire breaks</li> </ul> </li> </ul>                                                                                                                                                                                                                                                                                                                                        | <ul style="list-style-type: none"> <li>• <a href="#">Landscape fire management planning</a></li> </ul>                                                                                                                                                                                  |
| <ul style="list-style-type: none"> <li>• Assess fire hazard</li> <li>• Increase fire resilience at the stand level by managing surface fuels, species composition, density, crown base height, crown bulk density and age-class of forest stands</li> <li>• Reduce post-harvest fuels as necessary (e.g., biomass recovery, broadcast burning, pile and burn, mulching, chipping) <ul style="list-style-type: none"> <li>○ Choose appropriate season and weather for fuel-reduction</li> <li>○ Consider using grazing to manage fine fuels</li> </ul> </li> <li>• Do not reforest areas where climate change creates conditions where there is a low probability of producing commercial timber</li> </ul> | <ul style="list-style-type: none"> <li>• <a href="#">Landscape fire management planning</a></li> <li>• Fire management stocking standards<sup>35</sup></li> <li>• Fire and fuel management guidelines<sup>36</sup></li> </ul>                                                           |
| <ul style="list-style-type: none"> <li>• Reduce human-caused fires</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              | <ul style="list-style-type: none"> <li>• <a href="#">Wildfire Management Branch prevention strategy</a></li> </ul>                                                                                                                                                                      |
| <ul style="list-style-type: none"> <li>• Manage fire hazard around communities</li> <li>• Reduce risk to structures in interface areas</li> </ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          | <ul style="list-style-type: none"> <li>• <a href="#">Strategic wildfire prevention initiative</a><sup>37</sup></li> <li>• <a href="#">Fuel hazard assessment and abatement</a></li> <li>• <a href="#">FireSmart program</a></li> <li>• <a href="#">FireSmart communities</a></li> </ul> |

## Assisted Migration

When trees are harvested 60-120 years after they are planted, the climate could be 3-5 degrees warmer, exposing the trees to maladaptation and health risks. Moving populations of trees today (assisting migration) from their current provenance is one potential solution; growth and health are better when seeds are transferred to follow the climate in which they evolved. However trees have complex symbiotic relationships with many ectomycorrhizal fungal species in the soil and in some cases these bonds are tightly linked to local nutrient and climate conditions<sup>38</sup>. Improved understanding of these interactions in specific ecosystems may increase success. Government is leading a large, long-term [Assisted Migration Adaptation Trial](#) to understand tree species' climate tolerances. Findings are helping inform [Climate Based Seed Transfer policy](#) and tree species selection guidance.

## Range

Adaptation strategies for rangelands<sup>39</sup> aim to capitalize on opportunities (e.g., longer grazing season) and to reduce the negative impacts (e.g., increased invasive plants and drought stress) of climate change. Steps towards adaptation include revising expectation for rangelands to include climate dynamics and expecting higher variability in productivity between years. Adaptation strategies include

applying best range management practices such as: developing water storage and distribution systems to optimise water use while protecting riparian areas; building fences and locating salt where they will promote well-distributed use of the forage resource; and, enhancing monitoring to enable adjustments to stocking rates to reflect changing productivity.

## Monitoring

To develop adaptation strategies that are more likely to achieve management objectives, practitioners and decision-makers need to understand the direction and magnitude of changes in climate and key ecological responses at relevant spatial scales. In the Northeast, trend monitoring should include:

- Climate: temperature, precipitation, snowpack, permafrost, wind and extreme weather.
- Hydrology: stream flow by watershed, channel stability, forest cover, water temperature, erosion, suspended sediment, groundwater and aquifers.
- Disturbance: mass earth movements, insect and disease prevalence by BEC site series, site factors and seral stage, soil moisture, fire weather index and wildfire<sup>40</sup>.
- Biodiversity: regime shifts, seral stage, habitat supply, species health, invasive plant species and distribution shifts.
- Range: plant communities and condition, drought conditions.
- Tree growth and health.

Some of these data are already collected, but are not analysed regularly. A climate network that covers sub-regional variability with sufficient weather monitoring stations will be important.

## Moving Forward

Successful regional adaptation will require innovation and collaboration. Shared learning among practitioners, decision-makers and communities has the best potential for developing suitable adaptation strategies for the Northeast Region that foster resilient ecosystems and sustain natural resources into the future. This document could be a helpful catalyst in collaborative efforts.

## Contact for More Information

1. Regional specialist – Richard Kabzems, Research Silviculturist, Northeast Region, [Richard.kabzems@gov.bc.ca](mailto:Richard.kabzems@gov.bc.ca) (250-784-1256).
2. Provincial specialist – Kristine Weese, Sustainable Forest Management Specialist, Resource Practices Branch, [Kristine.weese@gov.bc.ca](mailto:Kristine.weese@gov.bc.ca) (250-558-1760).

---

<sup>1</sup> Ecological resilience is the capacity of an ecosystem to absorb, recover from and adapt to disturbance or stress caused by agents of change (such as climate change and natural resource management)

<sup>2</sup> Current research outcomes are primarily derived from projects under the [Future Forest Ecosystems Scientific Council](#) (FFESC) research program, but also including related regionally-relevant research

<sup>3</sup> See the report [A Climate Change Vulnerability Assessment for British Columbia's Managed Forests](#) (Morgan and Daust et al, 2013) for more insight into how climate change is expected to impact BC's forest ecosystems

- 
- <sup>4</sup> Projections are based on a combination of A2 and B1 emissions scenarios, where A2 represents roughly business as usual and B1 represents a more optimistic scenario with about ½ of emissions of business as usual ( Trevor Murdock, Pacific Climate Impacts Consortium)
- <sup>5</sup> Moore et al, [Chapter 3 \(Weather and Climate\)](#), Compendium of Forest Hydrology and Geomorphology in British Columbia (Pike et al, editors)
- <sup>6</sup> BEC zone acronyms: BWBS = Boreal White and Black Spruce; ESSF = Engelmann Spruce Subalpine Fir; SWB = Spruce Willow Birch
- <sup>7</sup> Rodenuis et al, [Hydro-climatology and future climate impacts in British Columbia](#) (2009)
- <sup>8</sup> Based on trends for all of BC
- <sup>9</sup> Details of the ensemble PCIC30 are given in Murdock and Spittlehouse, [Selecting and using climate change scenarios for British Columbia](#) (2011)
- <sup>10</sup> Wigley, The effect of changing climate on the frequency of absolute extreme events (2009) (Climatic Change 97:67-76; DOI 10.1007/s10584-009-9654-7) gives a theoretical analysis; Kharin et al, Changes in temperature and precipitation extremes in the CMIP5 ensemble (2013) (Climatic Change 119:345-357; DOI10.1007/s10584-013-0705-8) gives an analysis based on global climate models
- <sup>11</sup> BECWeb includes information on [BEC and climate change](#)
- <sup>12</sup> Wang et al, [Projecting future distributions of ecosystem climate niches: uncertainties and management implications](#) (2012)
- <sup>13</sup> For more information, see [Chapter 2c \(Natural Disturbance\)](#) of *A Climate Change Vulnerability Assessment for British Columbia's Managed Forests*
- <sup>14</sup> For more information, see summary in [Chapter 2b \(Hydrology and Aquatic Ecosystems\)](#) of *A Climate Change Vulnerability Assessment for British Columbia's Managed Forests*, and [Chapter 19 \(Climate Change Effects on Watershed Processes in BC\)](#) in the Compendium of Forest Hydrology and Geomorphology; for the Peace watershed, see Schnorbus et al, [Hydrologic impacts of climate change in the Peace, Campbell and Columbia Watersheds](#) (2011)
- <sup>15</sup> An analysis of streamflow trends in the Northeast is underway (contact Scott Jackson at [scott.jackson@gov.bc.ca](mailto:scott.jackson@gov.bc.ca) for more information)
- <sup>16</sup> See Footnote 10
- <sup>17</sup> For more information, see [Chapter 2e \(Forested Ecosystems\)](#) of *A Climate Change Vulnerability Assessment for British Columbia's Managed Forests*
- <sup>18</sup> See Peace River Regional District's [Invasive Plant Program 2014 Annual Report](#)
- <sup>19</sup> For more information, see [Chapter 2f \(Wildlife\)](#) of *A Climate Change Vulnerability Assessment for British Columbia's Managed Forests*
- <sup>20</sup> For more information, see Hamann and Wang, [Potential effects of climate change on ecosystem and tree species distribution in British Columbia](#) (2006)
- <sup>21</sup> Price and Daust, [Development of a Climate Change Index of Stress Using Future Projected BEC: Proof of Concept for the Nadina TSA](#) (2013)
- <sup>22</sup> Delong et al, [Stand level drought risk assessment tool](#) and [Assessing the risk of drought in BC forests using a stand-level water balance approach](#) (2011)
- <sup>23</sup> For more information, see [Climate Change and BC Range](#)
- <sup>24</sup> Dymond et al, [Diversifying managed forests to increase resilience](#) (2014)
- <sup>25</sup> Fletcher, *Towards a framework to support working with uncertainty in natural resource management (a discussion paper)* (2015); [link to be added prior to publication](#)
- <sup>26</sup> For example, see Leech et al, [Ecosystem management: A practitioners' guide](#) (2009)
- <sup>27</sup> [BC Drought Response Plan 2010](#) (being updated) and [Dealing with Drought \(a Handbook for Water Suppliers in BC\)](#) (2009)
- <sup>28</sup> [FREP water quality effectiveness evaluation indicators and protocols](#); [stream quality crossing index](#)
- <sup>29</sup> Redding et al, [Natural disturbance and post-disturbance management effects on selected watershed values](#) (2012)
-

- 
- <sup>30</sup> Wilford and Lalonde, [A framework for effective watershed monitoring](#) (2004)
- <sup>31</sup> Gayton and Almuedo, [Post-disturbance management of biodiversity in BC forests](#) (2012)
- <sup>32</sup> Updates to the Chief Forester's reference guide for FDP stocking standards based on climate change projections (2014)
- <sup>33</sup> Refer to Section 8, Page 15
- <sup>34</sup> Swift & Ran, [Successional Responses to Natural Disturbance, Forest Management, and Climate Change in British Columbia's Forests](#) (2012)
- <sup>35</sup> Guidance for designing fire management stocking standards is anticipated for release in June 2015
- <sup>36</sup> [Silvicultural regimes for fuel management](#); [Interim guidelines for fire and fuel management \(ABCFP, 2013\)](#)
- <sup>37</sup> This web site includes guidance for Community Wildfire Protection Plans, Fuels Management Prescriptions, and Operation Fuel Treatments
- <sup>38</sup> Recent research shows effects in coastal ecosystems. Kranabetter, Stoehr, and O'Neill, [Ectomycorrhizal fungal maladaptation and growth reductions associated with assisted migration of Douglas-fir](#) (2015)
- <sup>39</sup> For more information on managing the effects of climate change on BC rangelands, refer to: (i) Newman et al, [Managing for the ecological and socioeconomic effects of climate change on BC rangelands: developing strategic Range Use Plans, Range Stewardship Plans, and range management strategic documents](#) (2013); and, (ii) Range Branch's *Range Management Responses to Climate Change* (to be published in summer 2015)
- <sup>40</sup> [Provincial fire research and monitoring needs](#)