

DRAFT CENTRAL COAST BASE CASE FOR CURRENT ENVIRONMENTAL, SOCIAL, CULTURAL AND ECONOMIC CONDITIONS AND TRENDS

Draft Number 2

30 May 2012

DIRECTIONS FOR REPORT USE

This is intended for internal MaPP distribution only.



Marine Planning Partnership for the North Pacific Coast

DRAFT CENTRAL COAST BASE CASE FOR CURRENT ENVIRONMENTAL, SOCIAL, CULTURAL AND ECONOMIC CONDITIONS AND TRENDS

EXECUTIVE SUMMARY

- A base case is a description, or profile, of current biophysical and socioeconomic conditions within a planning area relevant to a plan's goals. As it pertains to the Marine Planning Partnership for the North Pacific Coast (MaPP), four sub-regional base cases are under development to provide snapshots of the current conditions and trends within the MaPP planning area. The MaPP sub-regional base cases follow the Ecosystem-based Management (EBM) framework and address the three EBM Elements: Ecological Integrity, Human Wellbeing and Governance.
- The features selected for the base case provide essential information that support several MaPP outputs including characterise and analyse existing conditions and trends (regional and sub-regional MaPP outputs), analyses that support EBM outputs such as marine spatial plans (regional and sub-regional MaPP outputs) and cumulative effects (regional and sub-regional MaPP outputs).
- Draft 1 of the Sub-regional base case (30 May 2012) follows the draft Base Case Table of Contents (Appendix 2) and includes eight features that address EBM Ecological Integrity and EBM Human Wellbeing: human communities, eelgrass beds, deepwater corals and sponges, herring, eulachon, Federal Marine Protected Areas (MPAs), Provincial MPAs and shellfish aquaculture.

TABLE OF CONTENTS

Executive Summary	2
Introduction and Purpose.....	4
Methods	5
Acronym Definitions	7
MaPP Planning Area	8
Central Coast Sub-regional Planning Area	9
Human Communities	10
Eelgrass Beds	13
Deep Water Corals and Sponges	16
Herring.....	19
Eulachon	22
Federal Marine Protected Areas.....	25
Provincial Marine Protected Areas.....	28
Shellfish Aquaculture.....	31
Acknowledgements	34
References	34
Appendix 1: Sub-regional Base Case Table of Contents (Draft - May 2012).....	42
Appendix 2: Central Coast Base Case Overview (Draft - May 2012).....	44

INTRODUCTION AND PURPOSE

The Marine Planning Partnership for the North Pacific Coast (MaPP) uses an ecosystem-based management, or EBM, framework to develop marine plans that address ecosystem integrity, human wellbeing and governance goals for the North Pacific Coast. Characterising and identifying current conditions within the MaPP planning area is critical for evaluating the successful implementation of the MaPP plans in 2014 and beyond.

Four sub-regional base cases are under development to summarise environmental, social, cultural and economic conditions and trends. Each base case will advance sub-regional plan development, including marine spatial planning, cumulative effects and climate change adaptation strategies. A base case is a description, or profile, of current biophysical and socioeconomic conditions within a planning area relevant to a plan's goals. A base case provides a snapshot of conditions and trends within a planning area, including a determination of what information is important relative to the goals and intended outputs or products of a planning process. A base case can be used to identify important data gaps. It supports completion of other components of an EBM framework for marine planning such as resolving spatial conflicts, cumulative effects assessments and strategies for climate change adaptation. Finally, a base case's ultimate function is to assess a plan's success after implementation.

Each Sub-regional base case will:

- Use both qualitative and quantitative information;
- Use best available data from public, published or peer-reviewed data sources;
- Characterise existing current conditions and trends relevant to MaPP outputs;
- Facilitate the creation of relevant and achievable planning objectives by providing benchmarks for ecological, socio-economic, cultural and governance/management features;
- Provide the data to assess the effectiveness of regional and sub-regional planning outputs; and
- Be a living document. Base cases are typically revised throughout the planning process as additional knowledge becomes available and planning needs defined.

METHODS

The Central Coast's (CC) base case was developed by reviewing existing marine, coastal and terrestrial plans, consulting Provincial and First Nation's governments and technical staff, reviewing MaPP outputs, and consulting international examples. Existing plans included BC Land and Resource Management Plans, terrestrial ecosystem-based management plans, and coastal plans such as the North Island Straits Coastal Plan. International examples consulted included the Massachusetts Ocean Management Plan and Baltic Sea Plan.

This draft base case was developed using environmental, social, cultural and economic features identified and reviewed by the Marine Coordination Team, Sub-regional Marine Technical Teams and Science Coordinator (Appendix 1). The draft Table of Contents for the base case contains at least 30 features and may be revised depending on data availability and MaPP discussions.

This report includes eight features from the draft Table of Contents with best available data from a variety of sources, including:

1. Public, published and peer-reviewed datasets;
2. Primary academic literature (i.e., peer-reviewed literature);
3. Discussions with taxonomic experts; and
4. Information provided by the Sub-regional Technical Teams.

Features for the base case were chosen using EBM principles that link human wellbeing, ecological integrity and governance. Each base case report begins with a brief description of the human communities within each sub-region. Marine uses and activities in the base case were included based on previous work of the Pacific North Coast Integrated Management Area (PNCIMA) Initiative to identify key marine uses and activities in the region. Ecological features were chosen based on an evaluation of the key habitats features, species and ecosystem structure and function for the sub-regions. Technical Teams from all sub-regions were given an opportunity to review and comment on the choice of features to be included within the base case.

All spatial data were compiled in ArcGIS 10 ArcEditor by Coastal Resource Mapping Ltd (CRMLTD). Spatial data were obtained from the same primary data sources that were used to develop the PNCIMA Atlas (PNCIMA 2011) and where additional data sources were used, they are noted on the relevant Table or Figure. Descriptions of the spatial data sources, spatial data layers, spatial statistics and spatial analyses

using ArcGIS are available from MaPP. Data quality assurance and quality control, or QA/QC, were done for all products developed by CRMLTD for the base case, including data layers previously peer-reviewed for the PNCIMA Atlas. If issues arose during QA/QC, the attribute table was examined for outliers or illogical values, for example, eelgrass beds located in depths greater than 100 metres. The outlier or illogical data were examined relative to ecological knowledge of the marine feature and where necessary, were removed and not spatially displayed or used in data summary tables. For example, data discrepancies resulted in removing less than one percent of the data points for deepwater corals and sponges and six percent of the occurrence data for eelgrass beds. Additional ecological, human wellbeing and governance features will be added to the base case, and available summer 2012 (Appendix 2)

ACRONYM DEFINITIONS

BC	British Columbia
BCMCA	British Columbia Marine Conservation Analysis
BCMEC	British Columbia Marine Ecological Classification
COSEWIC	Committee on the Status of Endangered Wildlife in Canada
DFO	Department of Fisheries and Oceans
EBM	Ecosystem-based Management
e.g.	for example
EI	Employment assistance
ha	hectare
i.e.	that is
IA	Important Area
IR	Indian Reserve
km	kilometre
LHA	Local Health Authority
MaPP	Marine Planning Partnership for the North Pacific Coast
MPA	Marine Protected Area
PNCIMA	Pacific North Coast Integrated Management Area
Rockfish Conservation Area	RCA
SARA	Species at Risk Act
VMEF	Valued Marine Environmental Features

MaPP PLANNING AREA




Figure 1: Overview of MaPP planning region including community names and select undersea and geographic features. Sub-regional boundaries are not finalised. All maps and data are approximate.

CENTRAL COAST SUB-REGIONAL PLANNING AREA




Figure 2: Overview of Central Coast sub-region planning area including community names and select undersea and geographic features. Sub-regional boundaries are not finalised. All maps and data are approximate.

HUMAN COMMUNITIES




Figure 3: Central Coast sub-region human communities. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Province of British Columbia 2011, PNCIMA Atlas 2011.

Description	Marine Uses and Activities
<p>The central coast is characterised by coastal islands peninsulas, deep fjords and glaciated river valleys. The only public road access to the area is via Highway 20 from Williams Lake to Bella Coola. A summer-only ferry service links Port Hardy with Bella Bella, Klemtu, Shearwater and Bella Coola. A year-round weekly ferry services Port Hardy, Klemtu, Bella Bella and Prince Rupert.</p> <p>The Central Coast Regional District encompasses most of the sub-region. It is 24,556 km² in area.</p>	<p>Central coast communities are closely tied to the marine environment. Marine resources underpin cultural, social, sustenance and economic activities. These include: sustenance and commercial fishing; aquaculture; marine tourism; sport fishing; and marine transportation.</p>
Current Population and Trends	

The population of the Central Coast Regional District was 3,206 people in 2011 (Table 1). The population density of the regional district is about 1/40th of the provincial average of 4.8 persons/km².

A small change in population can have a considerable impact in small communities. Between 1991 and 2011 population in the central coast area dropped by 8 percent—276 people. Over the same time period British Columbia's (BC) total population increased by 34 percent—1,117,996 people (Figure 4).




Figure 4: Population change over time. Sub-regional boundaries are not finalised. All maps and data are approximate. Source: British Columbia 2012

Table 1: Central coast populations (2011). Sub-regional boundaries are not finalised. All maps and data are approximate.

Community or Nation	Population
Walisla (Bella Bella IR)	1,095
Hagensborg, Firvale, Stuie (Electoral Areas C and D)	972
Ocean Falls, Shearwater (Electoral Area A)	129
Bella Coola IR	852
Bella Coola Townsite	95
Katit IR (Rivers Inlet)	63
Kitasoo/Xai'Xais	490
Heiltsuk	2,128
Nuxalk	1,477
Wuiknuxv	257

Education

Slightly over one third of the area's adult population ages 25-64 do not have formal education to high school or the equivalent level. This is almost twice the provincial average of adults who do not hold a certificate, diploma or degree.

Employment

Fishing is the largest goods producing sector of the area's economy—accounting for 7.3 percent of the Central Coast Regional District labour force in 2006, more than 70 times the provincial average. In many communities, fisheries is the most significant sector of employment. For example, the fisheries sector employs 41 percent of the labour force of the Kitasoo/Xai'Xais Nation living on reserves.

The largest proportion of the labour force in all of the reported First Nations communities in the central coast is employed in occupations related to business, finance and administration (Table 2).

Table 2: Workforce employed in occupations related to business, finance and administration (2008). Sub-regional boundaries are not finalised. All maps and data are approximate.

Nation	% Employed
Kitasoo/Xai'Xais	34%
Heiltsuk	41%
Nuxalk	36%
Wuiknuxv	39%

The unemployment rates in central coast communities are three to five times higher than the provincial average.

Income

Median income for males and females in the Central Coast Regional District in 2007 was \$14,595 and \$11,115, respectively, less than half of the provincial median incomes. The proportion of the labour force receiving Employment Income (EI) assistance based on 2006 census figures ranged from 20-23percent—about two times the provincial average of 11percent.

Health and Wellbeing

BC Stats prepares and index of socio-economic indicators for all regions of BC. The index includes indicators related to economic hardship, crime, health problems, education concerns, children at risk and youth at risk.

Out of 78 Local Health Authorities (LHAs) in BC, the Bella Coola Valley LHA is the second lowest ranked area in the province. Detailed ranking and figures for the central coast LHA are not released by BC Stats due to statistical and confidentiality concerns.

A limited compilation of community wellbeing indicators prepared by the Aboriginal Affairs and Northern Development Canada incorporates indicators related to education, labour force, income and housing. Census divisions identified as Indian Reserves in the central coast show a range of 0.57 to 0.64 under this index. The provincial average is 0.70 for First Nation communities and 0.85 for non-First Nation communities.

EELGRASS BEDS




Figure 5: Distribution of eelgrass beds in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Province of British Columbia 2008, BCMCA Atlas 2011, PNCIMA Atlas 2011.

Importance for MaPP	Social, Cultural, Economic Value																
<p>Eelgrass (<i>Zostera marina</i>) beds are natural nurseries and breeding areas for many species of commercial, cultural and social value such as crab, salmon, and herring. Eelgrass beds also act as natural stabilisers for nearshore sediments. Increasing protection for contiguous eelgrass beds is a possible MaPP strategy to manage for climate change effects and support populations of commercially and culturally valuable species.</p>	<p>The total global economic value of seagrass beds is enormous, estimated at US \$3,801 x 10⁹ per year. Absolute values have not been estimated for eelgrass beds in BC however the same ecosystem services are provided that resulted in the total global economic value namely estuarine filters, shoreline protection from storm effects, wildlife corridors, especially for salmon, and a carbon sink. The ability of eelgrass to sequester atmospheric carbon may provide future revenue through the carbon credits.</p>																
Description																	
<p>Eelgrass is a marine plant, an angiosperm, although it is often confused to be a seaweed or marine algae; it has roots, flowers and disperses seeds.</p> <p>Eelgrass beds are found in all four sub-regions. Eelgrass is most often found along sheltered sediment shores, as compared to surfgrasses which prefer exposed areas with strong wave action.</p> <p>The native species is <i>Zostera marina</i>. Currently, the invasive species <i>Z. japonica</i> is restricted to the Salish Sea region, south of the MaPP region.</p>	<p>Historically, eelgrass rhizomes were cooked in pits to help flavour meats such as deer and seal. They were also formed into cakes to be dried for winter. Eelgrass leaves were used to collect herring eggs.</p>																
Current Conditions and Trends	Current Management Actions																
<p><i>Table 3: Maximum, minimum and average depths and area covered of eelgrass beds within the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate.</i></p> <table> <tr> <th>Description</th><th>Depth</th></tr> <tr> <td>Minimum Depth</td><td>intertidal</td></tr> <tr> <td>Maximum Depth</td><td>9.9 m</td></tr> <tr> <td>Average Depth</td><td>0.7 m</td></tr> <tr> <td>Area covered</td><td>2040.3 km</td></tr> </table> <p>Eelgrass can grow in a variety of substrates. In the Central Coast, and generally those areas north of Campbell River, it is most commonly found growing in rocky substrates rather than soft substrates such as sand or mud as in other sub-regions.</p> <p>Few long-term research studies have been done on eelgrass in this sub-region and most of the mapping effort has focused on beds south of Campbell River. As a result, it is difficult to discern</p>	Description	Depth	Minimum Depth	intertidal	Maximum Depth	9.9 m	Average Depth	0.7 m	Area covered	2040.3 km	<p><i>Table 4: Percent of eelgrass beds distribution within the Central Coast sub-region captured by existing marine protection. Sub-regional boundaries are not finalised. All maps and data are approximate.</i></p> <table> <tr> <th>Type of Protection</th><th>Percent Coverage (%)</th></tr> <tr> <td>Federal</td><td>5.3</td></tr> <tr> <td>Provincial</td><td>1.1</td></tr> </table> <p>Eelgrass beds are currently protected under Canada's <i>Fisheries Act</i> since 1986 as essential fish habitat.</p>	Type of Protection	Percent Coverage (%)	Federal	5.3	Provincial	1.1
Description	Depth																
Minimum Depth	intertidal																
Maximum Depth	9.9 m																
Average Depth	0.7 m																
Area covered	2040.3 km																
Type of Protection	Percent Coverage (%)																
Federal	5.3																
Provincial	1.1																

long-term trends for the extent of eelgrass beds in the Central Coast. To increase what is known about eelgrass beds in the Central Coast, community groups have begun mapping eelgrass.

Climate Change Impacts

The long-term impacts of global warming on eelgrass are still relatively unknown. Climate change may also increase the intensity of storms, which could affect the amount of sediment entering rivers. An increase in sedimentation from terrestrial run off could cause higher turbidity, which could affect the ability of eelgrass to thrive.

Opportunities and Constraints

Eelgrass beds are vulnerable to human disturbance. Mapping of eelgrass beds to identify their locations and marine planning to reduce the overlap between eelgrass and marine uses and activities will minimise disturbance to rhizomes, flowers and leaves and increase the value to ecosystem services. For example, coastal logging can reduce the amount of light reaching eelgrass if sediment inputs increases to nearshore habitats. Log handling areas can cause scouring of eelgrass beds. Boat propeller wash can cause scouring and destroy roots of the eelgrass. Float homes and aquaculture sites can cause shading of the beds. Anchoring can physically uproot eelgrass plants.

Upland activities can influence eelgrass beds with increased run off and nutrient loading. Coastal eutrophication results in decreased light availability, which leads to decreased health of eelgrass beds. As a result upland activities also need to be considered when examining potential impacts on eelgrass beds.

DEEP WATER CORALS AND SPONGES




Figure 6: Distribution of coral and sponge communities and sponge reefs in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Fisheries and Oceans Canada 2007, Natural Resources Canada 2008, BCMCA Atlas 2011, PNCIMA Atlas 2011.

Importance for MaPP	Social, Cultural, Economic Value															
Deep water corals and sponges are found in all four sub-regions. Glass sponge reefs are formed by Hexactinellidae sponges and are only found in the North and Central Coast sub-regions within the MaPP planning area. They are long-lived, extremely fragile and easily destroyed by physical disturbance.	As with eelgrass beds, corals provide habits for a number of species including commercially and culturally valuable fish including rockfish and sablefish. As well, these biogenic structures provide habitat for up to 1300 species, many of which can act as prey species for mid-water species such as swordfish and turbot.															
Description																
Deep water coral and sponges are found in all four sub-regions. Glass sponge reefs are formed by Hexactinellidae sponges and are only found in the North and Central Coast sub-regions. They are long-lived, extremely fragile and easily destroyed by physical disturbance.																
In BC, the glass sponge reefs are globally unique biogenic habitat. They were first mapped in 1987-88 in Queen Charlotte Sound and Hecate Strait. There are more than 250 species of sponges in the Canadian Pacific.																
Current Conditions and Trends	Current Management Actions															
Corals have been known to fishermen for many years and science began actively studying their distribution in the early 1980s. Due to increased awareness and interested, there have been several academic-lead dedicated surveys for corals. Currently most of the location data are obtained from Department of Fisheries and Oceans (DFO) groundfish bottom trawls.	Fisheries and Oceans Canada has developed fishery management closures to protect all currently known locations of sponge reefs, primarily from bottom contact fishing gear. Currently coral and sponges and sponge reefs are protected by Federally designated areas. There are no provincially designated protected areas that are designated for coral or sponge protection.															
There are currently no coral or sponge species listed under <i>Species at Risk Act</i> (SARA) legislation.																
<i>Table 5: Maximum, minimum and average depths and area covered of coral and sponge communities and sponge reefs within the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate.</i>																
<table><tr><th>Description</th><th>Coral and sponges</th><th>Sponge Reefs</th></tr><tr><td>Min Depth (m)</td><td>167.6</td><td>170.9</td></tr><tr><td>Max Depth (m)</td><td>192.8</td><td>240.0</td></tr><tr><td>Average Depth (m)</td><td>180.2</td><td>207.1</td></tr><tr><td>Total area (km)</td><td>2222.5</td><td>201.3</td></tr></table>		Description	Coral and sponges	Sponge Reefs	Min Depth (m)	167.6	170.9	Max Depth (m)	192.8	240.0	Average Depth (m)	180.2	207.1	Total area (km)	2222.5	201.3
Description	Coral and sponges	Sponge Reefs														
Min Depth (m)	167.6	170.9														
Max Depth (m)	192.8	240.0														
Average Depth (m)	180.2	207.1														
Total area (km)	2222.5	201.3														
<i>Table 6: Percent of coral and sponge communities and sponge reefs within the Central Coast sub-region captured by existing marine protection. Sub-regional boundaries are not finalised. All maps and data are approximate.</i>																
<table><tr><th>Description</th><th>Coral and sponges (%)</th><th>Glass Sponge Reefs (%)</th></tr><tr><td>Federal</td><td>35.4</td><td>100</td></tr><tr><td>Provincial</td><td>0</td><td>0</td></tr></table>		Description	Coral and sponges (%)	Glass Sponge Reefs (%)	Federal	35.4	100	Provincial	0	0						
Description	Coral and sponges (%)	Glass Sponge Reefs (%)														
Federal	35.4	100														
Provincial	0	0														

Climate Change Impacts	Opportunities and Constraints
<p>Ocean acidification will be the most influential impact of climate change to affect corals and sponges as it will affect the ability of corals and sponges to produce and maintain their calcite or silicate structures.</p> <p>Other deepwater or oceanic impacts that corals and sponges may be sensitive to include: hypoxia due to the upwelling of deeper waters, increases in ocean temperatures, reduced oxygen levels, and increases in sedimentation</p>	<p>The primary source of mortality is from physical damage, such as contact with demersal trawl gear used in groundfish and shrimp fisheries. The recent fisheries closure of known locations of corals and sponge reefs is aimed at eliminating the physical disturbance to some of the largest aggregations in BC.</p>

HERRING




Figure 7: Herring spawn and Herring Important Areas. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Fisheries and Oceans Canada 2005, BCMCA Atlas 2011, PNCIMA Atlas 2011




Figure 8: Estimates of total biomass at the start of the year (solid line) and post-fishery spawning stock biomass (dashed line) for the Area 27 herring stock. Source: Fisheries and Oceans Canada 2011.

Importance for MaPP

Herring are an important food source for hundreds of marine species and an important cultural, social and economic species for First Nations and coastal communities. Increasing the percentage of Herring Important Areas (IA) in designated MPAs and developing strategies to improve the resilience of herring spawning habitats for the Central Coast Stock are possible outcomes of the Central Coast Marine Plan.

Description

Pacific herring, *Clupea pallasii*, is a pelagic species that inhabits the inshore and offshore waters of the Northeast Pacific Ocean. Ninety percent of herring return to spawn in the same general areas in subsequent years.

Five major and two minor herring stocks are managed in BC. The central coast contains one major stock, the Central Coast stock, and any central coast management recommendations intended to influence herring could target this stock.

Herring are an important food source for a variety of species, including Pacific hake, halibut, Arrowtooth flounder, and dogfish as well as Steller sea lions, harbour seals, harbour porpoises, Pacific white-sided dolphins and humpback whales. Herring annually migrate between feeding and spawning areas.

Social, Cultural, Economic Value

There are four Pacific herring commercial fisheries: roe, spawn-on-kelp, special use, and food and bait. As of 2007, herring fisheries in the MaPP planning area accounted for 30 to 50 percent of the coast-wide catch, and, on an economic basis, were one of the most important fisheries in BC. The landed value of BC's commercial herring fishery in 2010 was \$11.8 million, which accounted for 3.6 percent of the total landed value of the provincial capture fishery for that year. The value and catch level of the Province's commercial herring fishery has fluctuated over time (Figure 9).




Figure 9: Commercial herring landings in BC from 1951-1995 (Black line indicates weight, gray line indicates value). Source: Fisheries and Oceans Canada 2008.

Current Conditions and Trends	Current Management Actions
<p>Dramatic population fluctuations are common in all stocks of herring (Figure 8).</p> <p>In the mid 1960s, BC's herring stocks collapsed in response to poor environmental conditions and overfishing. The herring fishery was essentially closed from 1967-1971 and many stocks rapidly rebuilt.</p> <p>The Heiltsuk, who harvest herring for food, social and ceremonial purposes, were unable to harvest sufficient amounts of herring in 2005 and 2007.</p> <p>Over the last decade, herring have been declining on the central coast and the causes of low productivity are unknown. The fishery has been closed since 2008 and is not expected to open in 2012. Small roe and food and bait fisheries have occurred during closure years.</p> <p>Currently, the Central Coast stock has relatively low abundance (Figure 1). Recruitment for 2012 is anticipated to be poor—11,357 t of mature stock biomass—and is not anticipated to be open in 2012.</p>	<p>Approximately 10.9 percent and 4.1 percent of herring IAs are within federal and provincial protected areas, respectively. If not already in place, specific objectives for herring IAs would be needed within existing management plans to provide eulachon populations significant protection within the existing protected areas. Management efforts could potentially address the level and amount of protection afforded herring in the NVI sub-region.</p>
Climate Change Impacts	Opportunities and Constraints
<p>The coast of BC is experiencing an outburst of pelagic species abundance as current ecosystem conditions in BC seem to favour small pelagics. The synchronous change in a range of pelagic species across ecosystems indicates that a large-scale force is likely involved. Changes in herring spawning patterns coincide with changes in sea-surface temperature and other indices of climate change. Although climate change may be involved in the province's 'pelagic outburst', it is difficult to demonstrate because a range of mechanisms are likely involved.</p>	<p>Overfishing within the commercial fishery has historically led to fishery collapse (see above).</p> <p>The use of best management practices within marine activities that have the potential to degrade sensitive spawning and rearing herring habitat, such as shoreline development, log handling and uses that produce coastal or marine pollution, can reduce or prevent adverse impacts and protect herring resilience.</p> <p>When log handling sites are deactivated, foreshore riprap and other fill can be left in place so as to provide important habitat for herring.</p> <p>As herring produce sounds that may facilitate communication within the species, noise pollution from marine activities such as marine transportation may have consequences for herring behaviour.</p>

EULACHON




Figure 10: Eulachon river sites, run sizes and eulachon Important Areas. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Fisheries and Oceans Canada 2005 and PNCIMA Atlas 2011.

Importance for MaPP	Social, Cultural, Economic Value
<p>Eulachon, <i>Thaleichthys pacificus</i>, are a vital cultural, social and health component of First Nations communities in the central coast. Strategies to increase the percentage of Eulachon IAs in protected areas reduce or eliminate fisheries bycatch or increase catch rates for First Nations are possible measureable outcomes of the Central Coast Marine Plan.</p>	<p>Eulachon play a vital role to First Nations in BC due to their high nutritional value and cultural significance. They are a valued source of fresh protein in the spring and are dried for consumption during the rest of the year. The fish are rendered for their grease, once a highly valued trading item along the coast and in the interior, the so-named grease trails.</p>
Description	
<p>Eulachon are a small, anadromous fish ranging from the southern Bering Sea to northern California. They are an important part of First Nations diet and known as candlefish because of their high oil content. As of 2000, eulachon spawned in 33 rivers along the BC coast, 13 of which were on the central coast (Table 7). Management recommendations intended to influence eulachon could target these rivers and their marine waters they directly enter.</p> <p>After spawning, eulachon often, but not always, die and provide important nutrients to freshwater habitats and to scavenger species.</p> <p>Eulachon are a valuable prey species for terrestrial and marine birds because of their high fat content. Seals, sea lions, fish and whales also prey on eulachon in the ocean and in spawning rivers.</p>	<p>Catch rates of adult eulachon by First Nations vary with season, with relatively higher catches likely taking place in the summer.</p> <p>Eulachon IAs are areas worthy of enhanced protection as determined by eulachon and eulachon habitat uniqueness, aggregation, fitness consequences, naturalness and resilience (see Clarke and Jamieson 2006).</p>

Table 7: Known and probable central coast spawning areas (Spawning indicates whether a spawn is regular and occurs most years (R) or unknown (-); Fishing Effort identifies rivers routinely fished by First Nations (FN). Sub-regional boundaries are not finalised. All maps and data are approximate.

Spawning Area	Spawning	Fishing Effort	Marine Area	Stock
Kainet or Lard Creek	-		Kynoch Inlet – Mathieson Channel	N/A
Bella Coola River	R	FN	Dean Channel North Bentick Arm	Dean Channel
Kimsquit, Dean Rivers	R	FN	Dean Channel	Dean Channel
Noeick River	-	FN	South Bentick Arm	Dean Channel
Taleomy River	-	FN	South Bentick Arm	Dean Channel
Skowquiltz River	-	FN	Dean Channel – west side	Dean Channel
Cascade Inlet	-	FN	Dean Channel	Dean Channel
Kwatna River	-	FN	East side Burke Channel	Dean Channel
Chuckwalla/Kilbella	R	FN	Rivers Inlet – Queen Charlotte Strait	Rivers Inlet
Wannock/Oweekeno	R	FN	Rivers Inlet – Queen Charlotte Strait	Rivers Inlet
Clyak River, Moses Inlet	-		Rivers Inlet – Moses Inlet	Rivers Inlet
Hardy Inlet	-		Rivers Inlet	Rivers Inlet
Nekite River, Smith Inlet	-		Smith Inlet	N/A

Current Conditions and Trends	Current Management Actions
<p>Nearly all eulachon spawning runs have declined from California to southeastern Alaska in the past 30 years.</p> <p>All of the central coast populations are substantially lower than they were in the 1800s and before.</p> <p>The Kingcome and Klinaklini Rivers have had substantial declines but remain modest, while eulachon runs in Kitimat, Kemano, Bella Coola, and Rivers Inlet Rivers are nearly extirpated.</p>	<p>The Province has listed BC's eulachon as a species of 'special concern' (i.e., eulachon are blue listed).</p> <p>Eulachon Important Area (IAs) are areas worthy of enhanced protection as determined by eulachon and eulachon habitat uniqueness, aggregation, fitness consequences, naturalness and resilience (see Clarke and Jamieson 2006).</p> <p>Approximately 0.2 and 0 percent of eulachon Important Areas in the Central Coast sub-region are within existing federal or provincial protected areas, respectively. If not already in place, specific objectives for eulachon IAs would be needed within existing management plans to provide eulachon populations significant protection within the existing protected areas. Management efforts could potentially address the level and amount of protection afforded eulachon in the Central Coast sub-region.</p>
Climate Change Impacts	Opportunities and Constraints
<p>Climate change effects that are likely to negatively affect eulachon are changes to their marine or terrestrial food supplies and changes to their predator populations, for example Pacific hake (<i>Merluccius productus</i>). In addition, changes to sea surface temperatures are weakly associated with eulachon declines.</p>	<p>In the mid-1990s, increased shrimp trawling led to increased eulachon bycatch. In 1997, bycatch was estimated at 25 percent of that year's spawning biomass. Gear modifications and area closures were introduced to reduce bycatch in the 1990s and led to reduced bycatch. However, the fishery is likely still affecting eulachon recovery.</p> <p>Preventing or minimising dredging and using best management practices in forestry proximate to and during spawning will increase survival of in-river eulachon adults and eggs by limiting changes to river volume, flow rate, sediment level and discharge pattern. Logging-related marine activities in estuaries can minimise debris deposition to prevent anoxic water conditions that lead to juvenile and adult mortality.</p> <p>Shrimp and hake catches, seal and sea lion abundance and increased water temperatures are weakly associated with eulachon declines.</p>

FEDERAL MARINE PROTECTED AREAS




Figure 11: Federal MPAs in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: Fisheries and Oceans Canada 2007, Parks Canada 2010, PNCIMA Atlas 2011.

Importance for MaPP

The *National Framework for Canada's Network of Marine Protected Areas* contains 13 ecologically defined bioregions within the nation's oceans and Great Lakes. The MaPP planning area is located within the Northern Shelf bioregion. The Government of Canada recognises a range of benefits associated with well-designed and effectively implemented marine protection strategies, including the extension of the protection afforded by terrestrial protection; enhancement of fish recruitment, including commercially valuable species; achievement of sustainable fisheries management objectives; reduction of the effects of local catastrophes through the protection of multiple ecosystems and regions that provide buffers against local environmental change; support for First Nations traditional activities; provision of social and economic benefits, such as enhanced opportunities for recreation, promotion of cultural heritage and enhanced planning and coordination of ocean activities; and protection of unique, endemic, rare and threatened species.

Description

MPAs are established to protect a portion of the sea's surface, water column and/or seabed. They may be designated to protect specific plants, animals or commercial, recreational, scientific, cultural and/or historic features. The three core elements of Canada's federal MPA network include Fisheries and Oceans Canada's MPAs, Environment Canada's Marine Wildlife Areas and Parks Canada's National Marine Conservation Areas. Other federal MPAs within the network include Environment Canada's migratory bird sanctuaries and national wildlife areas as well as and Parks Canada's national parks.

Social, Cultural, Economic Value

MPAs can provide social, cultural and economic benefits both within and proximate to their boundaries by improving habitat and species wellbeing and abundance. No-take MPAs, which prohibit resource extraction, have been linked to increased density, biomass, size and reproductive potential of individuals as well as increased species richness. Partial protection MPAs that regulate instead of prohibit resource use within their boundaries have also been connected to a range of benefits relative to open access areas, including higher biomass, organism density, species richness and individual organism size. As marine protection can lead to spillover, which is the export of individuals such as adults or larvae from protected areas to adjacent waters, the benefits MPAs often extend past their boundaries.

Current Conditions and Trends

The Central Coast sub-region currently contains two types of Federal MPAs: Rockfish Conservation Areas (RCAs) and Sponge Reef Fishery Closures (Table 8).

Table 8: Type, number and coverage of federal protection areas in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate.

MPA Type	Count	Total Area (km ²)	Percent of Sub-region Covered (%)
Rockfish Conservation Area	8	395	0.8
Sponge Reef Fishery Closure	2	788	1.6

Climate Change Effects

Marine protection contributes to climate change mitigation and adaptation by protecting habitats that capture and store carbon, such as coastal salt marshes, sea grasses and kelp forests; conserving multiple ecosystem types and special ecological features, such as underwater canyons or critical habitats, as insurance against catastrophic events; and protecting coastal barrier ecosystems, such as wetlands, that

buffer ecosystems and human developments against extreme weather events.

Current Management Actions

Federal and provincial MPAs, including provincial conservancies designated to address First Nations interests, cover an area equal to 28.2 percent of the Central Coast sub-region, while federal MPAs alone cover an area equal to 2.5 percent of the Central Coast sub-region. Fisheries and Oceans Canada is currently implementing its *Federal Marine Protected Areas Strategy*. While RCAs and Sponge Reef Fishery Closures serve different purposes, they are both administered by Fisheries and Oceans Canada under the *Fisheries Act* (Table 9).

Table 9: Administration and purposes of federal protected areas. Sub-regional boundaries are not finalised. All maps and data are approximate.

MPA Type	Agency Act	Purpose
Rockfish Conservation Area	Fisheries and Oceans Canada <i>Fisheries Act</i>	To preserve rockfish biomass
Sponge Reef Fishery Closure	Fisheries and Oceans Canada <i>Fisheries Act</i>	To conserve and protect fish and fish habitat; to manage inland fisheries (among other purposes)

Within the federal protected areas on the central coast, a range of activities are permitted, restricted or prohibited. Within RCAs, the permissible recreational fisheries are invertebrates by hand picking or dive, crab by trap, shrimp/prawn by trap and smelt by gillnet, while the permissible commercial fisheries are invertebrates by hand picking or dive, crab or prawn by trap, scallops by trawl, salmon by seine or gillnet, herring by gillnet, seine and spawn-on-kelp, sardine by gillnet, seine and trap, smelt by gillnet, euphausiid (krill) by mid-water trawl, opal squid by seine and groundfish by mid-water trawl. Within the Sponge Reef Fisheries Closure, bottom trawling is prohibited and mid-water trawling, prawn and shrimp fishing by trap and bottom longline fishing are permitted. In federal protected areas, Aboriginal rights for food, social and ceremonial purposes are not impacted by protected areas unless conservation issues arise.

The federal and provincial governments are working together to coordinate marine protection efforts off the coast of BC as demonstrated by the development of the *Draft Canada – British Columbia Marine Protected Area Network Strategy* in 2011.

Opportunities and Constraints

MPAs can influence activities within their boundaries by affecting what uses are permitted, including allowing, restricting or prohibiting activities; how activities can be conducted, such as requiring specific gear types or gear modifications; when activities are acceptable, including seasonal closures; and who may conduct activities, such as commercial or recreational users. While MPAs can affect resource use, they can lead to benefits within their boundaries as well as spillover effects that combine to create net gains for marine resource users.

PROVINCIAL MARINE PROTECTED AREAS




Figure 12: Provincial MPAs in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: PNCIMA Atlas 2011, Province of British Columbia 2011.

Importance for MaPP

The Province of BC recognises that MPAs are a vital part of its commitment to viable coastal communities, sustainable economies and a healthy, diverse marine environment. The Province has committed to protecting and conserving the natural beauty and richness of BC's marine areas, maintaining ecological diversity and preserving the province's marine recreational, natural and cultural features for all time through the creation and maintenance of provincial MPAs.

Description

MPAs provide legally binding protection of the sea surface, water column, seabed and/or associated flora, fauna and recreational, scientific, cultural and historical features within their boundaries. The MPAs within BC include Ecological Reserves, Provincial Parks, Protected Areas, Conservancies, Recreation Areas and Wildlife Management Areas.

Social, Cultural, Economic Value

MPAs have social, cultural and economic value as they provide benefits (see the Federal MPAs section above for further detail) that support a range of marine uses, including, but not limited to, commercial fisheries, recreational activities and the collection of species by First Nations for food, social and ceremonial purposes. As MPAs conserve historic and archeological sites in addition to ecological sites, they contribute to the long-term preservation of the diverse identities of BC's inhabitants.

Current Conditions and Trends

Federal and provincial MPAs, including provincial conservancies designated to address First Nations interests, cover an area equal to 28.2 percent of the Central Coast sub-region, while provincial MPAs alone (not including First Nations-related conservancies) cover an area equal to 7.6 percent of the Central Coast sub-region. The Central Coast sub-region currently contains a provincial Ecological Reserve and eight Provincial Parks (Table 10).

Table 10: Type, number and coverage of provincial MPAs in the Central Coast sub-region. Sub-regional boundaries are not finalised. All maps and data are approximate.

MPA Type	Count	Total Area (km ²)	Percent of Sub-region Covered (%)
Ecological Reserve	1	10	0.02
Provincial Park	8	3,634	7.3

As demonstrated by the creation of the *Draft Canada – British Columbia Marine Protected Area Network Strategy* in 2011, the provincial government is working with the Government of Canada on the creation of a coordinated network of MPAs for BC's coast.

Climate Change Effects

As noted above in the Federal MPAs section, marine protection efforts can contribute to climate change mitigation and adaptation. MPAs can support carbon sequestration by conserving marine habitats, contribute to the ongoing delivery of ecosystem goods and services, such as habitat, food and nutrient cycling, by safeguarding replicate ecosystem types and ecological features, and shelter coastal ecosystems and human settlements during extreme weather events by buffering impacts.

Current Management Actions

The Province is developing planning products to support the establishment of MPAs as well as management in existing MPAs. These products include the British Columbia Marine Ecological Classification (BCMEC); a preliminary list of Valued Marine Environmental Features (VMEF), which are key conservation, recreation and cultural-heritage elements of marine environments such as kelp beds, abandoned canneries and archaeological sites; and a MPA gap analysis based on the BCMEC, VMEFs and identification of biophysical and human use values within existing MPAs. The gap analysis work will be used to identify potential candidate areas for provincial conservation as well as existing provincial protected areas that might benefit from federal fisheries closures.

The MPAs types currently within the Central Coast sub-region serve different purposes, are administered under different provincial Acts and manage marine uses to different degrees (Tables 10 and 11). Aboriginal rights for food, social and ceremonial purposes are not impacted by protected areas unless conservation issues arise.

Table 11: Administration and purposes of BC Provincial protected areas.

MPA Type	Agency Act	Purpose
Ecological Reserve	BC Ministry of Environment <i>Ecological Reserves Act</i>	To reserve Crown land for ecological purposes, including areas suitable for research and education, that are representative of natural ecosystems in BC that are examples of modified environments that can be studied for their recovery, and that are habitats of rare, endangered or unique species
Provincial Park	BC Ministry of Environment <i>Park Act</i>	To protect representative ecosystems and special natural, cultural and recreation features, and to provide outdoor recreation in natural environments for the inspiration, use and enjoyment of the public

Opportunities and Constraints

As noted above in the Federal MPAs section, MPAs can influence activities within their boundaries by affecting what uses are permitted, how activities can be conducted, when activities are acceptable and who may conduct activities. At the same time, MPAs contribute to ecosystem resilience and productivity and can provide spillover effects that lead to overall gains for marine resource users (see above).

SHELLFISH AQUACULTURE




Figure 13: Central Coast sub-region marine aquaculture (shellfish) tenures and applications. Sub-regional boundaries are not finalised. All maps and data are approximate. Sources: PNCIMA Atlas 2011, Province of British Columbia.

Importance for MaPP	Conditions and Trends
Shellfish aquaculture is a significant component of BC's coastal economy with significant and consistent increases in production and value in recent years, as well as potential for increased production. Activities associated with shellfish production and processing have potential to provide employment and income in coastal communities. MaPP provides an opportunity to identify and address related ecosystem integrity, human wellbeing and governance considerations.	As of December 2010, two active shellfish aquaculture licence holders and one application were registered in the Central Coast sub-region. Shellfish species of interest for farming in BC include: <ul style="list-style-type: none"> Pacific oyster (<i>Crassostrea gigas</i>), Manila clam (<i>Venerupis philippinarum</i>) with 89percent of total cultured shellfish harvest in 2010. Japanese scallop (<i>Patinopecten yessoensis</i>), giant rock scallop (<i>Crassadoma gigantea</i>) and blue and Gallo mussel (<i>Mytilus spp.</i>) with 11 percent of 2010 harvest. Giant Californian sea cucumber (<i>Parastichopus californicus</i>), sea urchins (<i>Strongylocentrotus spp.</i>), geoduck clam (<i>Panopea abrupta</i>), spot prawn (<i>Pandalus platyceros</i>). Limited or no cultured production presently. Abalone (<i>Haliotis kamtschatkana</i>). Potential for aquaculture if population recovery, protection and tracking concerns addressed.
Description	
Shellfish aquaculture is the production of farmed marine invertebrates in an aquatic environment or human-made container of water for commercial purposes, including the cultivation of shellfish on, in or under the foreshore or in open water. Types of shellfish farming: <ul style="list-style-type: none"> Off-beach from rafts or buoys, generally higher productivity than on-beach methods. On or near seabed in the intertidal zone, may conflict with wild shellfish stocks. Closed containment, no commercial operations at present. 	
Social, Cultural, Economic Value	

Coastal First Nations have tended clam gardens along the BC coast over many centuries and shellfish are an important component of the seafood diet for central coast populations. Oyster seed was introduced to BC in 1903. Clam farming began in the mid-1980s and in recent years the farmed harvest of clams has exceeded the wild harvest. Recent production and value of shellfish harvest in BC is shown below (Table 12).

Table 12: BC Shellfish Production – Harvest and Landed Value 2008-2010. Sub-regional boundaries are not finalised. All maps and data are approximate.

	BC-HARVEST ('000 Tonnes)				BC-LANDED VALUE (\$ Millions)			
	2008	2009	2010	% Change (09/10)	2008	2009	2010	% Change (09/10)
Clams	1.3	1.3	1.5	+15.4%	7.2	7.2	8.1	+12.5%
Oysters	5.6	5.7	7.4	+29.8%	6.5	7.1	8.8	+23.9%
Scallops & Mussels	0.6	0.7	1.1	+57.1%	2.5	3.0	4.8	+60.0%
TOTAL	7.5	7.7	10.0	+29.9%	16.2	17.3	21.7	+25.4%

Climate Change Impacts	Opportunities and Constraints
<p>Ocean acidification has been linked to the collapse of oyster seed production in Washington and Oregon and is a concern for BC.</p> <p>The water along the Pacific Coast is layered. During certain weather events, deep layers are brought to the surface, bringing old water that has absorbed carbon dioxide and carbonic acid, a process of ocean acidification to describe waters with a lower pH than fresher, oxygen-rich surface waters. The acidic water dissolves the calcium carbonate in the shells of oyster larvae faster than they can produce it, resulting in mortality and thus failure in seed production.</p> <p>Monitoring sea water at existing or proposed shellfish aquaculture hatchery locations for pH and other factors, such as temperature, salinity and bacteria, can support hatchery operators and shellfish farmers in identifying and adapting to periods of low pH.</p>	<p>Ecosystem Integrity: Potential to impact beaches and/or sites supporting wild shellfish stocks or other marine species. Potential for introduction of invasive and/or exotic species. Shellfish can affect food supply of adjacent or nearby species. Vulnerable to contamination from shore or water-based activities. Off-beach farming has lower impacts than on-bottom farming with potential for further reduction of impacts if closed containment used.</p> <p>Human Wellbeing: Capital investment needs, risk of harvest failures and cash flow requirements can limit participation in this sector. Investment in hatchery and/or processing operations can provide additional revenue and employment opportunities. Potential to provide year round and seasonal employment. Wages and return on effort/investment are not assured. Requires training and experience involving business and practical skill sets. Transportation costs and access to processing facilities and/or markets may limit viability.</p>
Current Management Actions	Governance
<p>All commercial marine, freshwater and land-based finfish and shellfish aquaculture operations are licensed under the Pacific Aquaculture Regulation of the federal <i>Fisheries Act</i>. The Province remains responsible for the administration of Crown land for aquaculture and issuance of <i>Fisheries Act</i> aquaculture business licenses.</p> <p>In December 2011 the Department of Fisheries and Oceans released an interim plan for discussion addressing integrated management of shellfish aquaculture.</p> <p>The Coastal First Nations and North Coast Skeena First Nations Stewardship Society have been working with federal and provincial agencies and partners over the past decade in a coordinated effort to establish commercial shellfish aquaculture hatchery and farming operations.</p>	<p>DFO is in the process of developing a new regulatory and management regime. At present the regulatory approval process is lengthy and uncertain.</p> <p>Species-specific considerations</p> <p>Scallops: Grow well in cool, nutrient rich waters. Capital intensive, with limited seed sources. Established demand in BC with potential to increase market to the US and Asia.</p> <p>Oysters: Most are currently grown in southern waters of the province where markets are established and transportation costs are low. Shucking and packing is labour intensive but unshucked oysters have a lower value for weight than shucked. Background cadmium levels may limit exports to China.</p> <p>Clams: On-beach farming competes with wild stocks and other uses. Established markets and low transportation costs.</p> <p>Other Species: Opportunities for high economic and cultural value species such as geoduck clams, abalone and sea cucumbers.</p>

ACKNOWLEDGEMENTS

Report developed by the Marine Planning Partnership for the North Pacific Coast (MaPP) a partnership between the Province of B.C., represented by the Ministry of Forests, Lands and Natural Resource Operations, and the Coastal First Nations, the North Coast – Skeena First Nations Stewardship Society and the Nanwakolas Council, which include 20 member First Nations.

REFERENCES

Human Communities

Notes:

1. Detailed data from the 2011 Canada Census are scheduled for release in 2013. Population estimates are available to 2011 however socio-economic indices, community wellbeing and aboriginal community data rely for the most part on 2006 and preceding Canada Census figures.
 2. Population figures for communities are from BC Stats sources. Population figures for First Nations are from Government of Canada sources and include on-reserve and off-reserve populations. The population numbers include overlaps, as some off-reserve registered aboriginals live in communities that are documented in BC Stats population estimates.
- .British Columbia. 2010. Overall Regional Socio-economic Index. BC Stats.
Links to data and methodology reports available from BC Stats at:
<http://www.bcstats.gov.bc.ca/StatisticsBySubject/SocialStatistics/SocioEconomicProfilesIndices/SocioEconomicIndices/LHAREports.aspx>
- British Columbia. 2011. Census Subdivisions. GeoBC
- British Columbia. 2012. Community Facts. BC Stats.
Links to reports for Regional Districts and communities available from BC Stats at:
<http://www.bcstats.gov.bc.ca/StatisticsBySubject/SocialStatistics/CommunityFacts.aspx>
- British Columbia. 2012. 2011 Sub-Provincial Population Estimates. BC Stats
- Ference Weicker & Company. 2009. Social and Economic Assessment and Analysis of First Nation Communities and Territorial Natural Resources for Integrated Marine Use Planning in the Pacific North Coast Integrated Management Area. Report can be accessed from:
<http://ccira.ca/media/documents/pdf/marine-sector-report-f-w.pdf>
- Government of Canada. 2009. 2006 Aboriginal Community Data Initiative. Statistics Canada.
Links to individual reports by aboriginal community are provided by BC Stats:
www.bcstats.gov.bc.ca/StatisticsBySubject/AboriginalPeoples/AboriginalReports.aspx
- Government of Canada. 2010. First Nation and Inuit Community Well-Being : Describing Historical Trends (1981-2006). Department of Indian Affairs and Northern Development.
- Government of Canada. 2012. Registered Indian Population by Sex and Residence 2010. Department of Indian Affairs and Northern Development.

Sheltair Group. 2009. Ecosystem-based Management Human Wellbeing Indicators 2006 Baseline Report – Final. Prepared for Ecosystem-based Management Working Group for the North and Central Coast. Report can be accessed from:
http://archive.ilmb.gov.bc.ca/slrp/lrmp/nanaimo/cencoast/ebmwg_docs/hw02_baseline_final_report.pdf

Eelgrass Beds

Booth, J., N. Dale, and D. Haggarty. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences.

British Columbia Marine Conservation Analysis. 2011. Marine Atlas of Pacific Canada: a product of the British Columbia Marine Conservation Analysis (BCMCA), British Columbia Marine Conservation Analysis, 2011.

Dunster, K. 2003. Eelgrass Mapping Review: Eelgrass Mapping Initiatives in Coastal British Columbia. Prepared for Canadian Wildlife Service. Delta, BC. Accessed May 2012.
http://www.shim.bc.ca/eelgrass/Eelgrass_Mapping_Inventory_Final_v1.pdf.

Hall, A. 2008. State of the Ocean in the Pacific North Coast Integrated Management Area (PNCIMA). David Suzuki Foundation. 159pp.

Hutchings, J.A., Cote, I.M., Dodson, J.J., Fleming, I.A., Jennings, S., Mantua, N.J., Peterman, R.M., Riddell, B.E., Weaver, A.J., and D.L. VanderZwaag. 2012. Sustaining Canadian marine biodiversity: responding to the challenges pose by climate change, fisheries, and aquaculture. Expert panel report prepared for the Royal Society of Canada, Ottawa.

Lucas, B.G., Johannessen, D. and Lindstrom, S. 2007. Appendix E: Marine plants. In Ecosystem overview: Pacific North Coast Integrated Management Area (PNCIMA). Edited by Lucas, B.G., Verrin, S., and Brown, R. Can. Tech. Rep. Fish. Aquat. Sci. 2667: iv + 23 p.

Okey, T.A., H.M. Alidina, A. Montenegro, V. Lo, S. Agbayani, S. Jessen. 2012. Climate change impacts and Vulnerabilities in Canada's Pacific marine ecosystems. CPAWS BC and WWF Canada, Vancouver, BC.

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Province of British Columbia. 2008. ShoreZone Mapping System. Coastal and Oceans Resources Inc. Available at <http://www.coastalandoceans.com/shorezone.html>

Seagrass Conservation Working Group. 2004. The BC Coastal Eelgrass Stewardship Project. Accessed at: http://www.stewardshipcentre.bc.ca/static/eelgrass/eelgrassfinalreport12_04.pdf

Seagrass Conservation Working Group. 2005. Communities connecting to place: A strategy for eelgrass restoration in British Columbia. Accessed at: <http://www.stewardshipcentre.bc.ca>.

Sloan, N.A., and P.M. Bartier. 2000. Living Marine Legacy of Gwaii Haanas I: Marine Plant Baseline to 1999 and Plant-related Management Issues. Technical Reports in Ecosystem Science; no. 27.

Deepwater Corals and Sponges

- Ardron, J.A. 2005. Protecting British Columbia's Corals and Sponges from Bottom Trawling. A report by Living Oceans Society (v. 1.0). Box 320, Sointula, BC, V0N 3E0.
- British Columbia Marine Conservation Analysis. 2011. Marine Atlas of Pacific Canada: a product of the British Columbia Marine Conservation Analysis (BCMCA). British Columbia Marine Conservation Analysis, 2011.
- Crawford, W.R. and J.R. Irvine [eds]. 2011. State of physical, biological, and selected fishery resources of Pacific Canadian marine ecosystems in 2010. Fisheries Oceanography Working Group (FOWG). Fisheries and Oceans Canada, Sidney, BC.
- Cummins, P. and R. Haigh. 2010. Ecosystem Status and Trends Report for North Coast and Hecate Strait ecozone. Fisheries and Oceans Canada Research Document 2010/045, Sidney, BC.
- Hall, A. 2008. State of the Ocean in the Pacific North Coast Integrated Management Area (PNCIMA). David Suzuki Foundation. 159pp.
- Jamieson, G. 2009. British Columbian corals. Fisheries and Oceans Canada, Nanimo, British Columbia. Accessed at: <http://www.geog.ubc.ca/biodiversity/efauna/BritishColumbianCorals.html>
- Johannessen, D.I., Macdonald, J.S., Harris, K.A., and Ross, P.S. 2007. Marine environmental quality in the Pacific North Coast Integrated Management Area (PNCIMA), British Columbia, Canada: A summary of contaminant sources, types and risks. Can. Tech. Rep. Fish. Aquat. Sci. 2716: xi + 53p.
- Natural Resources Canada. 2008. Sponge Reefs.
- Okey, T.A., H.M. Alidina, A. Montenegro, V. Lo, S. Agbayani, S. Jessen. 2012. Climate change impacts and Vulnerabilities in Canada's Pacific marine ecosystems. CPAWS BC and WWF Canada, Vancouver, BC.
- Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org
- Prescott-Allen, R. 2005. Coast Information Team: Review Report. Accessed at <http://www.citbc.org/>

Herring

- Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.
- British Columbia Marine Conservation Analysis. 2011. Marine Atlas of Pacific Canada: a product of the British Columbia Marine Conservation Analysis (BCMCA). British Columbia Marine Conservation Analysis, 2011.
- Fisheries and Oceans Canada. 2005. Herring Important Areas.

Fisheries and Oceans Canada. 2008. Commercial herring landings in British Columbia. Available from: <http://www.pac.dfo-mpo.gc.ca/stats/comm/summ-somm/her-har/chart-tab/index-eng.htm> [accessed 29 May 2012].

Fisheries and Oceans Canada. 2011. Stock Assessment Report on Pacific Herring in British Columbia in 2011. Canadian Science Advisory Secretariat Science Advisory Report. 15 pp.

G3 Consulting Ltd. 2003. Guidebook: environmentally sustainable log handling facilities in British Columbia. Report prepared for Fisheries and Oceans Canada, Pacific and Yukon Region, Habitat and Enhancement Branch by G3 Consulting Ltd., Burnaby BC. Pp 96.

Hay, D.E., McCarter, P.B. and Daniel, K.S. 2001. Tagging of Pacific herring *Clupea pallasii* from 1936-1992: a review with comments on homing, geographic fidelity, and straying. Canadian Journal of Fisheries and Aquatic Sciences. 58 (7): 1356-1370.

Hourston, A.S. 1980. The decline and recovery of Canada's Pacific herring stocks. International Council for the Exploration of the Sea. 177: 143-153.

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Moody, R. 2007. Heiltsuk Tribal Council Herring Spawn of Kelp Coordinator. Personal Communication, June 9, 2007. In: Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.

Province of British Columbia. 2011. Capture Fisheries Statistics. Available from: <http://www.env.gov.bc.ca/omfd/fishstats/capture/index.html> [accessed 28 May 2012].

Schweigert, J., Funk, F., Oda, K. And Moore, T. 2002. Herring size-at-age variation in the North Pacific. PICES-Globec International Program on Climate Change and Carrying Capacity. 20: 47-57.

Ware, D.M. and Tovey, C. 2004. Pacific herring spawn disappearance and recolonization events. Canadian Science Advisory Secretariat. Research Document 2004/008. Pp. 52.

Wilson, B., Batty, R.S. and Dill, L.M. 2004. Pacific and Atlantic herring produce burst pulse sounds. Proceedings of the Royal Society B: Biological Sciences, Supplement. 271: S95-S97.

Eulachon

Beacham, T.D., Hay, D.E. and Le, K.D. 2005. Population Structure and Stock Identification of Eulachon (*Thaleichthys pacificus*), an Anadromous Smelt, in the Pacific Northwest. Marine Biotechnology. 7: 363-372.

Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.

- Clarke, A.D., Lewis, A., Telmer, K.H. and Shrimpton, J.M. 2007. Life history and age at maturity of an anadromous smelt, the eulachon *Thaleichthys pacificus* (Richardson). *Journal of Fish Biology*. 71: 1479-1493.
- Clarke, C.L. and Jamieson, G.S. 2006. Identification of Ecologically and Biologically Significant Areas in the Pacific North Coast Integrated Management Area: Phase I - Identification of IAs. Canadian Technical Report of Fisheries and Aquatic Sciences. 2678: 1-97.
- Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2011. Fishes (marine). Available from:
http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm?id=1163&StartRow=1&boxStatus=All&boxTaxonomic=All&location=All&change=All&board=All&commonName=eulachon&scienceName=&returnFlag=0&Page=1 [accessed 11 May 2012].
- Fisheries and Oceans Canada. 2012. Species at Risk Consultations – Pacific Region Calendar. Available from: <http://www.pac.dfo-mpo.gc.ca/consultation/sara-lep/cal-eng.htm> [accessed 26 May 2012].
- Government of Canada. 2011. Species at Risk Public Registry. Available from:
http://www.sararegistry.gc.ca/sar/index/default_e.cfm?type=species&lng=e&index=1&cosid=&common=eulachon&scientific=&population=&taxid=0&locid=0&desid=0&schid=0&desid2=0& [accessed 26 May 2012].
- Fisheries and Oceans Canada. 2005. Eulachon Important Areas.
- Hay, D. and McCarter, P.B. 2000. Status of the eulachon *Thaleichthys pacificus* in Canada. Canadian Stock Assessment Secretariat. 2000/145. 99 pp.
- Kuhnlein, H., Yeboah, F., Sedgemore, M., Sedgemore, S. and Chan, H. 1996. Nutritional qualities of ooligan grease: a traditional food fat of British Columbia First Nations. *Journal of Food Composition and Analysis*. 9(4): 18-31.
- Moody, M.F. and Pitcher, T.J. 2010. Eulachon (*Thaleichthys pacificus*): past and present. Fisheries Centre Research Reports. 18(2): 1-199.
- Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org
- NatureServe. 2012. Explorer. Available from:
http://www.natureserve.org/explorer/servlet/NatureServe?loadTemplate=tabular_report.wmt&aging=home&save=all&sourceTemplate=reviewMiddle.wmt [accessed 26 May 2012].
- Province of British Columbia. BC Species and Ecosystems Explorer. Available from:
<http://a100.gov.bc.ca/pub/eswp/search.do;jsessionid=0ddac6a86b3e25fa0e100b2fd6071ffa5bea582ca3b42fe20d5e68ebda50ca35.e3uMah8KbhmLe34Tb3aOa3uOaN90n6jAmljGr5XDqQLvpAe> [accessed 28 May 2012].

Federal Marine protected areas

Coastal Resources Mapping. 2012. Protected Area Summary Tables. Generated for the Marine Planning Process (MaPP) for the North Pacific Coast, 12 May 2012. Available from MaPP.

Fisheries and Oceans Canada. 2005. Canada's federal Marine Protected Areas Strategy. Government of Canada, Ottawa. 18 pp. Available from <http://www.dfo-mpo.gc.ca/oceans/publications/fedmpa-zpmfed/pdf/mpa-eng.pdf> [accessed 26 May 2012].

Fisheries and Oceans Canada. 2006. Canada's Submission on Efforts and Action taken pursuant to Paragraphs 66 to 69 of United Nations General Assembly Resolution 59/25 to Identify, Manage and Protect Sensitive Marine Ecosystems and Species. Available from: http://www.dfo-mpo.gc.ca/misc/un59_25-eng.htm [accessed 27 May 2012].

Fisheries and Oceans Canada. 2007. Sponge reef closures, rockfish conservation areas.

Fisheries and Oceans Canada. 2009. Rockfish Conservation Areas. Available from: <http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/restricted-restreint/rca-acsc-eng.htm> [accessed 27 May 2012].

Fisheries and Oceans Canada. 2011. Marine Protection. Available from: <http://www.dfo-mpo.gc.ca/oceans/management-gestion/marineprotection-protectionmarine/index-eng.htm> [accessed 26 May 2012].

Fisheries and Oceans Canada. 2011. National framework for Canada's Network of Marine Protected Areas. Government of Canada, Ottawa. 31 pp. Available from <http://www.dfo-mpo.gc.ca/oceans/publications/dmpaf-eczpm/docs/framework-cadre2011-eng.pdf> [accessed 26 May 2012].

Fisheries and Oceans Canada. 2011. Socio-economic implications of the proposed Glass Sponge Reef Marine Protected Area. Policy and Economic Analysis Branch. Pacific Region. Pp. 25.

Government of Canada. 1985. *Fisheries Act*.

Government of Canada and Province of British Columbia. 2011. DRAFT Canada – British Columbia Marine Protected Area Network Strategy. Available from: <http://www.mccpacific.org/MPA-DraftCanadaBCStrat-May27.pdf> [accessed 27 May 2012].

Lester, S.E. and Halpern, B.S. 2008. Biological responses in marine no-take reserves versus partially protected areas. *Marine Ecology-Progress Series*. 367: 49-56.

Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Aïramé, S. and Warner, R.R. 2009. Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*. 384: 33-46.

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Provincial Marine protected areas

Coastal Resources Mapping. 2012. Protected Area Summary Tables. Generated for the Marine Planning Process (MaPP) for the North Pacific Coast, 12 May 2012. Available from MaPP.

Fisheries and Oceans Canada. 2011. National framework for Canada's Network of Marine Protected Areas. Government of Canada, Ottawa. 31 pp. Available from <http://www.dfo-mpo.gc.ca/oceans/publications/dmpaf-eczpm/docs/framework-cadre2011-eng.pdf> [accessed 26 May 2012].

Government of BC. 1996. *Ecological Reserve Act*.

Government of BC. 1996. *Park Act*.

Government of Canada and Province of British Columbia. 1998. Marine Protected Areas Discussion Paper. Available from: <http://archive.ilmb.gov.bc.ca/cis/rpts/disppap.htm> [accessed 26 May 2012].

Government of Canada and Province of British Columbia. 2011. DRAFT Canada – British Columbia Marine Protected Area Network Strategy. Available from: <http://www.mccpacific.org/MPA-DraftCanadaBCStrat-May27.pdf> [accessed 27 May 2012].

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Province of British Columbia. 2009. Marine Protected Areas. Available from: <http://www.env.gov.bc.ca/omfd/ocean-resources/mpa.html> [accessed 26 May 2012].

Province of British Columbia. 2011. Provincial Parks, Ecological Reserves, Protected Areas, Conservancies.

Shellfish Aquaculture

Barton, A., Hales, B., Waldbusser, G.G., Langdon, C. and Freely, R. 2012. The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects. *Limnology and Oceanography* 57 (3): 698 DOI: [10.4319/lo.2012.57.3.0698](https://doi.org/10.4319/lo.2012.57.3.0698)

British Columbia. Ministry of Agriculture. 2011. British Columbia Seafood Industry 2010 year in Review. 216 pp. <http://www.bcseafood.ca/PDFs/YIR-2010.pdf>

Edwin Blewett & Associates Inc. and Nelson Brothers Fisheries Ltd. 2003. North Islands Straits Shellfish Aquaculture Barriers & Constraints Study. April 2003. 40 pp. http://www.agf.gov.bc.ca/fisheries/cabinet/NI_Shellfish_Aq_Barriers_Constraints.pdf

Fisheries and Oceans Canada. 2011. Pacific Region Integrated Management of Aquaculture Plan–Shellfish Interim. November 29, 2011. 36 pp.

Government of Canada. Fisheries and Oceans Canada. Pacific Region. 2011. Pacific Region Integrated Management of Aquaculture Plan Shellfish Interim Nov 29, 2011. 36 pp. http://www.agf.gov.bc.ca/fisheries/cabinet/NI_Shellfish_Aq_Barriers_Constraints.pdf

Kingzett Professional Services Ltd. and AXYS Environmental Consulting Ltd. 2002. First nations Shellfish Aquaculture Regional Business Strategy: B.C. Central and Northern Coast. Prepared for Aboriginal Relations and Economic measures, Land and Water British Columbia Inc. November 2002. 254 pp. http://www.agf.gov.bc.ca/fisheries/cabinet/FN_Shellfish_Aquaculture_North_Coast_Strategy.pdf

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Snuneymuxw First Nation. 2008. Traditional Sea Foods Balancing Health Benefits with Pollution Risks. Final Report. Proceedings of a Workshop at Malaspina Campus, Vancouver Island University. April 5 2008. <http://www.indigenousfoodsvi.ca/wp-content/uploads/2011/08/2008-final-report.pdf>

APPENDIX 1: SUB-REGIONAL BASE CASE TABLE OF CONTENTS (DRAFT - MAY 2012)

1 The Central Coast

Human Communities	X
-------------------------	---

2 Coastal and Benthic Habitats

Eelgrass beds	X
Kelp forest	X
Estuaries	X
Rocky reefs	X
Deep water corals and sponges.....	X
Deep water canyons	X
Continental shelf break & slope	X

2 Marine Species

Zooplankton.....	X
Herring.....	X
Eulachon	X
Abalone.....	X
Salmon.....	X
Groundfish	X
Seabirds	X
Cetaceans	X
Species Guilds	X

4 Ecosystem Structure and Function

Gyres, eddies, seasonal flows and general currents.....	X
Coastal upwelling.....	X
Areas of high tidal flow	X
Hypoxia	X
Degraded areas.....	X
Freshwater inputs	X

5 Existing Marine protected areas

Federal Marine protected areas	X
Provincial Marine protected areas	X
Provincially Designated First Nations sites	X

6 Marine Uses and Activities

Fisheries and Seafood Processing, Marketing and Branding.....	X
Marine Pollution	
Shellfish, Seaweed and Finfish Aquaculture and Processing.....	X
Marine Recreation and Tourism	X
Marine Transportation and Vessel Safety	X
Logging-related Marine Activities.....	X
Marine and Coastal Infrastructure	X
Shore-based Marine Tenures	

Renewable Ocean Energy X

7 Governance and Collaborative Management

Information Sharing X

Monitoring, Compliance and Enforcement X

Collaborative Management Agreements X

Integrated Economic Development.....

Marine Policies and Regulations..... X

Legislation..... X

APPENDIX 2: CENTRAL COAST BASE CASE OVERVIEW (DRAFT - MAY 2012)

Overview

This is a project description and Executive Summary for deliverables produced for the Marine Planning Partnership.

Executive Summary:

- A base case is a description, or profile, of current biophysical and socioeconomic conditions within a planning area relevant to a plan's goals. As it pertains to the Marine Planning Partnership for the North Pacific Coast (MaPP), four sub-regional base cases are under development to provide snapshots of the current conditions and trends within the MaPP planning area. The MaPP sub-regional base cases follow the Ecosystem-based Management (EBM) framework and address the three EBM Elements: Ecological Integrity, Human Wellbeing and Governance.
- The features selected for the base case provide essential information that support several MaPP outputs including characterise and analyse existing conditions and trends (regional and sub-regional MaPP outputs), analyses that support EBM outputs such as marine spatial plans (regional and sub-regional MaPP outputs) and cumulative effects (regional and sub-regional MaPP outputs).
- Draft 2 of the Sub-regional base case (May 2012) follows the Base Case Table of Contents and includes eight features that address EBM Ecological Integrity and EBM Human Wellbeing: human communities, eelgrass beds, deepwater corals and sponges, herring, eulachon, Federal Marine Protected Areas, Provincial Marine Protected Areas and shellfish aquaculture.

Date first draft of report was completed: 14 May 2012

Date report was finalised: TBD

Prepared for: Marine Coordination Team, Central Coast Marine Tech Team

Prepared by: MaPP Contracted Support

Roles of other Parties: Review by MCT, Sub-regional Marine Technical Teams, Science Coordinator and Science Advisory Committee.

Constraints, Risks and Dependencies: Completion of all the features in the base case Table of Contents is constrained by data availability and QA/QC for relevant data layers.

Resources (human, \$, other): Coast Resources Mapping Ltd for developing all maps and tabulated data outputs.

Timelines, Outputs and Responsibilities:

Timeline	Output	Who is responsible for this
----------	--------	-----------------------------

14 May 2012	Completion of the first drafts of the sub-regional base cases	Contracted Support
14-25 May 2012	Review of the first drafts of the sub-regional base cases	Marine Coordination Team, Sub-regional Technical Teams, Science Coordinator
26-29 May 2012	Incorporate edits and update sub-regional base cases	Contracted support
30 May 2012	Distribution of the draft sub-regional base cases to Sub-regional Marine Plan Advisory Committee (MPAC) members	Sub-Regional Technical Team Co-leads
June 2012	Next draft of the final draft of the sub-regional base cases	Contracted Support
June–July 2012	Review of the final drafts of the sub-regional base cases. Distribution to MPAC.	Marine Coordination Team, Sub-regional Technical Teams, Science Coordinator, Science Advisory Committee
Summer 2012	Completion of base case project (final edits)	Contracted Support



CENTRAL COAST BASE CASE

Susan Dean

MAPP CONTRACTED SUPPORT

- The objectives of this presentation are to:
 - Explain what a base case is and how it can be used within the Central Coast's marine planning process
 - Present the Draft Central Coast Base Case for your review
 - Obtain your feedback on the Draft Central Coast Base Case



OUTLINE



- What is a base case?
- Where does information come from?
- How is it used?
- Who is responsible for it?
- When will it be finished?
- Base case walk through



WHAT IS A BASE CASE: DEFINITION



A base case is a description, or snapshot, of current biophysical and socioeconomic conditions within a planning area relevant to its goals



WHERE DOES INFORMATION COME FROM



- Best available data
- Published sources
- Traditional and local knowledge



WHERE DOES INFORMATION COME FROM



- Information Quality

- Reliable and transparent sources

- Information Type

- Ecological value
 - Human value
 - MaPP value

- Information Scale

- Sub-regional, regional, provincial, federal



HOW IS A BASE CASE USED: APPLICATION IN MaPP



- **Provides an initial picture of the planning area**
 - What indicators do we want to track?
- **Supports analysis and plan development leading to completion of MaPP outputs**
 - How can we influence indicators in a positive way?
- **Supports assessment of plan success**
 - How have indicators changed? Did we accomplish what we set out to do?



WHO IS RESPONSIBLE: BASE CASE DEVELOPMENT



- Development: MaPP Contracted Support
- Information: You and your sub-region
 - Tell us what information is needed
 - Be an information source
 - Help direct us to other information sources



WHEN WILL THE BASE CASE BE FINISHED: TIMELINE




- The first draft of the Central Coast Base Case will be completed for Summer 2012
- Base cases are living documents and information will be added throughout the planning process



- Please consider the following questions during our review of the Draft Central Coast Base Case:
 - Are there any errors or obvious omissions in the information presented?
 - Is there information included that requires clarification?
 - Are there any additional sources or reports that should be included in the Central Coast Base Case?

MAPP BASE CASE: TABLE OF CONTENTS



- Central Coast
 - Human Communities
- Ecological Integrity
 - Marine Species
 - Coastal and Benthic Habitats
 - Ecosystem Structure and Function
 - Existing Marine Protected Areas
- Human Well-being
 - Marine Uses and Activities
- Governance and Collaborative Management

Central Coast Sub-Region Human Wellbeing



Human Communities


- Description
- Current Population and Trends
- Marine Uses and Activities
- Education
- Employment
- Health and Wellbeing
- Income



Eelgrass Beds

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints


Central Coast Sub-Region Sea Floor Habitats



Deep Water Corals and Sponges

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints


Central Coast Sub-Region Species (Herring)



Herring

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints

Central Coast Sub-Region Species (Eulachon)




Eulachon

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints

ECOLOGICAL INTEGRITY: EXISTING MARINE PROTECTED AREAS



Central Coast Sub-Region Protected Areas (Federal)




Federal Marine Protected Areas

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints

ECOLOGICAL INTEGRITY: EXISTING MARINE PROTECTED AREAS



Central Coast Sub-Region Protected Areas (Provincial)



Provincial Marine Protected Areas

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints

Central Coast Sub-Region Marine Aquaculture (Shellfish)



Shellfish Aquaculture

- Importance for MaPP
- Description
- Current Conditions and Trends
- Social, Cultural, Economic Value
- Current Management Actions
- Climate Change Impacts
- Opportunities and Constraints

- Are there any errors or obvious omissions in the information presented?
- Is there information included that requires clarification?
- Are there any additional sources or reports that should be included in the Central Coast Base Case?



THANK YOU! QUESTIONS?





Central Coast Base Case Resources

To: Central Coast Marine Technical Team

From: Marine Contracted Support

Date: 30 May 2012

CONTEXT

This document provides (1) the resources used to create the maps and text of the Draft Central Coast Base Case as well as (2) a compilation of the ecological integrity and human wellbeing resources collected to date by MaPP Contracted Support for use in ongoing base case development. Individual resources are included within all the topic areas to which they apply and, as a result, may be listed more than once.

The intent of the document is to:

- Facilitate the identification of text resources that MPAC members feel are currently missing and that they would like to see included in future iterations of the Central Coast Base Case.
- Clearly outline the dates and sources of spatial data so that MPAC members can indicate if they are aware of different or more up-to-date spatial data resources.

SPATIAL DATA RESOURCES

Human Communities

- PNCIMA Atlas (2011)
- Province of British Columbia (2011) – Census Subdivisions

Deep Water Corals and Sponges

- BCMCA Atlas (2011) – Bathymetry
- Fisheries and Oceans Canada (2007) – Sponge/Coral Important Areas
- Natural Resources Canada (2008) – Sponge Reefs
- PNCIMA Atlas (2011)

Eelgrass Beds

- BCMCA Atlas (2011)

- BCMCA Project Team (2010)
- Community Mapping Network – Eelgrass Surveys
- Fisheries and Oceans Canada – Eelgrass Surveys
- Living Oceans Society – Merged Eelgrass Datasets
- Parks Canada – Central Coast Marine Plants
- PNCIMA Atlas (2011)
- Province of British Columbia (2008) – Shorezone Mapping System, Bioband
- Province of British Columbia – Eelgrass Surveys

Herring

- BCMCA Atlas (2011) – Herring Spawn
- Fisheries and Oceans Canada (2005) – Herring Important Areas
- PNCIMA Atlas (2011)

Eulachon

- Fisheries and Oceans Canada (2005) – Eulachon Important Areas
- PNCIMA Atlas (2011) – Eulachon run

Existing Federal Protected Areas

- Fisheries and Oceans Canada (2007) – Sponge Reef Closures, Rockfish Conservation Areas
- Parks Canada (2010) – National Parks
- PNCIMA Atlas (2011)

Existing Provincial Protected Areas

- PNCIMA Atlas (2011)
- Province of British Columbia (2011) – Provincial Parks, Ecological Reserves and Protected Areas

First Nations Protected Areas

- PNCIMA Atlas (2011)
- Province of British Columbia (2011) – Conservancies

Shellfish Aquaculture

- PNCIMA Atlas (2011) - Aquaculture
- Province of British Columbia – Tantalus, Crown Tenures

TEXT RESOURCES

Human Communities

British Columbia. 2010. Overall Regional Socio-economic Index. BC Stats. Links to data and methodology reports available from BC Stats at:

<http://www.bcstats.gov.bc.ca/StatisticsBySubject/SocialStatistics/SocioEconomicProfilesIndices/SocioEconomicIndices/LHAREports.aspx>

British Columbia. 2011. Census Subdivisions.

British Columbia. 2012. 2011 Sub-Provincial Population Estimates. BC Stats.

British Columbia. 2012. Community Facts. BC Stats. Links to reports for Regional Districts and communities available from BC Stats at:
<http://www.bcstats.gov.bc.ca/StatisticsBySubject/SocialStatistics/CommunityFacts.aspx>

Ference Weicker & Company. 2009. Social and Economic Assessment and Analysis of First Nation Communities and Territorial Natural Resources for Integrated Marine Use Planning in the Pacific North Coast Integrated Management Area. Report can be accessed from:
<http://ccira.ca/media/documents/pdf/marine-sector-report-f-w.pdf>

Government of Canada. 2009. 2006 Aboriginal Community Data Initiative. Statistics Canada. Links to individual reports by aboriginal community are provided by BC Stats:
www.bcstats.gov.bc.ca/StatisticsBySubject/AboriginalPeoples/AboriginalReports.aspx

Government of Canada. 2010. First Nation and Inuit Community Well-Being : Describing Historical Trends (1981-2006). Department of Indian Affairs and Northern Development.

Government of Canada. 2012. Registered Indian Population by Sex and Residence 2010. Department of Indian Affairs and Northern Development.

Sheltair Group. 2009. Ecosystem Based Management Human Wellbeing Indicators 2006 Baseline Report – Final. Prepared for Ecosystem Based Management Working Group for the North and Central Coast. Report can be accessed from:
http://archive.ilmb.gov.bc.ca/slrp/lrmp/nanaimo/cen coast/ebmwg_docs/hw02_baseline_final_report.pdf

Eelgrass Beds

Booth, J., N. Dale, and D. Haggarty. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences.

British Columbia Marine Conservation Analysis. 2011. Marine Atlas of Pacific Canada: a product of the British Columbia Marine Conservation Analysis (BCMCA), British Columbia Marine Conservation Analysis, 2011.

Dunster, K. 2003. Eelgrass Mapping Review: Eelgrass Mapping Initiatives in Coastal British Columbia. Prepared for Canadian Wildlife Service. Delta, BC. Accessed May 2012.
http://www.shim.bc.ca/eelgrass/Eelgrass_Mapping_Inventory_Final_v1.pdf.

Hall, A. 2008. State of the Ocean in the Pacific North Coast Integrated Management Area (PNCIMA). David Suzuki Foundation. 159pp.

Hutchings, J.A., Cote, I.M., Dodson, J.J., Fleming, I.A., Jennings, S., Mantua, N.J., Peterman, R.M., Riddell, B.E., Weaver, A.J., and D.L. VanderZwaag. 2012. Sustaining Canadian marine biodiversity: responding to the challenges pose by climate change, fisheries, and aquaculture. Expert panel report prepared for the Royal Society of Canada, Ottawa.

Lucas, B.G., Johannessen, D. and Lindstrom, S. 2007. Appendix E: Marine plants. In Ecosystem overview: Pacific North Coast Integrated Management Area (PNCIMA). Edited by Lucas, B.G., Verrin, S., and Brown, R. Can. Tech. Rep. Fish. Aquat. Sci. 2667: iv + 23 p.

Okey, T.A., H.M. Alidina, A. Montenegro, V. Lo, S. Agbayani, S. Jessen. 2012. Climate change impacts and Vulnerabilities in Canada's Pacific marine ecosystems. CPAWS BC and WWF Canada, Vancouver, BC.

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Seagrass Conservation Working Group. 2004. The BC Coastal Eelgrass Stewardship Project. Accessed at: http://www.stewardshipcentre.bc.ca/static/eelgrass/eelgrassfinalreport12_04.pdf

Seagrass Conservation Working Group. 2005. Communities connecting to place: A strategy for eelgrass restoration in British Columbia. Accessed at: <http://www.stewardshipcentre.bc.ca>.

Sloan, N.A., and P.M. Bartier. 2000. Living Marine Legacy of Gwaii Haanas I: Marine Plant Baseline to 1999 and Plant-related Management Issues. Technical Reports in Ecosystem Science; no. 27.

Deepwater Corals and Sponges

Ardron, J.A. 2005. Protecting British Columbia's Corals and Sponges from Bottom Trawling. A report by Living Oceans Society (v. 1.0). Box 320, Sointula, BC, V0N 3E0.

British Columbia Marine Conservation Analysis. 2011. Marine Atlas of Pacific Canada: a product of the British Columbia Marine Conservation Analysis (BCMCA). British Columbia Marine Conservation Analysis, 2011.

Crawford, W.R. and J.R. Irvine [eds]. 2011. State of physical, biological, and selected fishery resources of Pacific Canadian marine ecosystems in 2010. Fisheries Oceanography Working Group (FOWG). Fisheries and Oceans Canada, Sidney, BC.

Cummins, P. and R. Haigh. 2010. Ecosystem Status and Trends Report for North Coast and Hecate Strait ecozone. Fisheries and Oceans Canada Research Document 2010/045, Sidney, BC.

Hall, A. 2008. State of the Ocean in the Pacific North Coast Integrated Management Area (PNCIMA). David Suzuki Foundation. 159pp.

Jamieson, G. 2009. British Columbian corals. Fisheries and Oceans Canada, Nanaimo, British Columbia. Accessed at: <http://www.geog.ubc.ca/biodiversity/efauna/BritishColumbianCorals.html>

Johannessen, D.I., Macdonald, J.S., Harris, K.A., and Ross, P.S. 2007. Marine environmental quality in the Pacific North Coast Integrated Management Area (PNCIMA), British Columbia, Canada: A summary of contaminant sources, types and risks. Can. Tech. Rep. Fish. Aquat. Sci. 2716: xi + 53p.

Okey, T.A., H.M. Alidina, A. Montenegro, V. Lo, S. Agbayani, S. Jessen. 2012. Climate change impacts and Vulnerabilities in Canada's Pacific marine ecosystems. CPAWS BC and WWF Canada, Vancouver, BC.

Pacific North Coast Integrated Management Area Initiative. 2011. Atlas of the Pacific North Coast Integrated Management Area. Available at www.pncima.org

Prescott-Allen, R. 2005. Coast Information Team: Review Report. Accessed at <http://www.citbc.org/>

Herring

Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.

- Fisheries and Oceans Canada. 2008. Commercial herring landings in British Columbia. Available from: <http://www.pac.dfo-mpo.gc.ca/stats/comm/summ-somm/her-har/chart-tab/index-eng.htm> [accessed 29 May 2012].
- Fisheries and Oceans Canada. 2011. Stock Assessment Report on Pacific Herring in British Columbia in 2011. Canadian Science Advisory Secretariat Science Advisory Report. 15 pp.
- G3 Consulting Ltd. 2003. Guidebook: environmentally sustainable log handling facilities in British Columbia. Report prepared for Fisheries and Oceans Canada, Pacific and Yukon Region, Habitat and Enhancement Branch by G3 Consulting Ltd., Burnaby BC. Pp 96.
- Hay, D.E., McCarter, P.B. and Daniel, K.S. 2001. Tagging of Pacific herring *Clupea pallasii* from 1936-1992: a review with comments on homing, geographic fidelity, and straying. Canadian Journal of Fisheries and Aquatic Sciences. 58 (7): 1356-1370.
- Hourston, A.S. 1980. The decline and recovery of Canada's Pacific herring stocks. International Council for the Exploration of the Sea. 177: 143-153.
- Moody, R. 2007. Heiltsuk Tribal Council Herring Spawn of Kelp Coordinator. Personal Communication, June 9, 2007. In: Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.
- Province of British Columbia. 2011. Capture Fisheries Statistics. Available from: <http://www.env.gov.bc.ca/omfd/fishstats/capture/index.html> [accessed 28 May 2012].
- Schweigert, J., Funk, F., Oda, K. And Moore, T. 2002. Herring size-at-age variation in the North Pacific. PICES-Globec International Program on Climate Change and Carrying Capacity. 20: 47-57.
- Ware, D.M. and Tovey, C. 2004. Pacific herring spawn disappearance and recolonization events. Canadian Science Advisory Secretariat. Research Document 2004/008. Pp. 52.
- Wilson, B., Batty, R.S. and Dill, L.M. 2004. Pacific and Atlantic herring produce burst pulse sounds. Proceedings of the Royal Society B: Biological Sciences, Supplement. 271: S95-S97.

Eulachon

- Beacham, T.D., Hay, D.E. and Le, K.D. 2005. Population Structure and Stock Identification of Eulachon (*Thaleichthys pacificus*), an Anadromous Smelt, in the Pacific Northwest. Marine Biotechnology. 7: 363-372.
- Booth, J., Dale, N. and Haggarty, D. 2007. Ecosystem Assessment Pacific North Coast Management Area (PNCIMA). Canadian Technical Report of Fisheries and Aquatic Sciences. 215 pp.
- Clarke, A.D., Lewis, A., Telmer, K.H. and Shrimpton, J.M. 2007. Life history and age at maturity of an anadromous smelt, the eulachon *Thaleichthys pacificus* (Richardson). Journal of Fish Biology. 71: 1479-1493.
- Clarke, C.L. and Jamieson, G.S. 2006. Identification of Ecologically and Biologically Significant Areas in the Pacific North Coast Integrated Management Area: Phase I - Identification of IAs. Canadian Technical Report of Fisheries and Aquatic Sciences. 2678: 1-97.
- Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2011. Fishes (marine). Available from: http://www.cosewic.gc.ca/eng/sct1/searchdetail_e.cfm?id=1163&StartRow=1&boxStatus=All&

[boxTaxonomic=All&location=All&change=All&board=All&commonName=eulachon&scienceName=&returnFlag=0&Page=1](#) [accessed 11 May 2012].

Fisheries and Oceans Canada. 2012. Species at Risk Consultations – Pacific Region Calendar. Available from: <http://www.pac.dfo-mpo.gc.ca/consultation/sara-lep/cal-eng.htm> [accessed 26 May 2012].

Government of Canada. 2011. Species at Risk Public Registry. Available from: http://www.sararegistry.gc.ca/sar/index/default_e.cfm?type=species&lng=e&index=1&cosid=&common=eulachon&scientific=&population=&taxid=0&locid=0&desid=0&schid=0&desid2=0& [accessed 26 May 2012].

Hay, D. and McCarter, P.B. 2000. Status of the eulachon *Thaleichthys pacificus* in Canada. Canadian Stock Assessment Secretariat. 2000/145. 99 pp.

Kuhnlein, H., Yeboah, F., Sedgemore, M., Sedgemore, S. and Chan, H. 1996. Nutritional qualities of ooligan grease: a traditional food fat of British Columbia First Nations. Journal of Food Composition and Analysis. 9(4): 18-31.

Moody, M.F. and Pitcher, T.J. 2010. Eulachon (*Thaleichthys pacificus*): past and present. Fisheries Centre Research Reports. 18(2): 1-199.

NatureServe. 2012. Explorer. Available from: http://www.natureserve.org/explorer/servlet/NatureServe?loadTemplate=tabular_report.wmt&paging=home&save=all&sourceTemplate=reviewMiddle.wmt [accessed 26 May 2012].

Province of British Columbia. BC Species and Ecosystems Explorer. Available from: <http://a100.gov.bc.ca/pub/eswp/search.do;jsessionid=0ddac6a86b3e25fa0e100b2fd6071ffa5bea582ca3b42fe20d5e68ebda50ca35.e3uMah8KbhmLe34Tb3aOa3uOaN90n6jAmljGr5XDqQLvpAe> [accessed 28 May 2012].

Federal Marine Protected Areas

Coastal Resources Mapping. 2012. Protected Area Summary Tables. Generated for the Marine Planning Process (MaPP) for the North Pacific Coast, 12 May 2012. Available from MaPP.

Fisheries and Oceans Canada. 2005. Canada's federal Marine Protected Areas Strategy. Government of Canada, Ottawa. 18 pp. Available from <http://www.dfo-mpo.gc.ca/oceans/publications/fedmpa-zpmfed/pdf/mpa-eng.pdf> [accessed 26 May 2012].

Fisheries and Oceans Canada. 2006. Canada's Submission on Efforts and Action taken pursuant to Paragraphs 66 to 69 of United Nations General Assembly Resolution 59/25 to Identify, Manage and Protect Sensitive Marine Ecosystems and Species. Available from: http://www.dfo-mpo.gc.ca/misc/un59_25-eng.htm [accessed 27 May 2012].

Fisheries and Oceans Canada. 2009. Rockfish Conservation Areas. Available from: <http://www.pac.dfo-mpo.gc.ca/fm-gp/rec/restricted-restraint/rca-acs-eng.htm> [accessed 27 May 2012].

Fisheries and Oceans Canada. 2011. Marine Protection. Available from: <http://www.dfo-mpo.gc.ca/oceans/management-gestion/marineprotection-protectionmarine/index-eng.htm> [accessed 26 May 2012].

Fisheries and Oceans Canada. 2011. National framework for Canada's Network of Marine Protected Areas. Government of Canada, Ottawa. 31 pp. Available from <http://www.dfo->

mpo.gc.ca/oceans/publications/dmpaf-eczpm/docs/framework-cadre2011-eng.pdf [accessed 26 May 2012].

Fisheries and Oceans Canada. 2011. Socio-economic implications of the proposed Glass Sponge Reef Marine Protected Area. Policy and Economic Analysis Branch. Pacific Region. Pp. 25.

Government of Canada. 1985. *Fisheries Act*.

Government of Canada and Province of British Columbia. 2011. DRAFT Canada – British Columbia Marine Protected Area Network Strategy. Available from: <http://www.mccpacific.org/MPA-DraftCanadaBCStrat-May27.pdf> [accessed 27 May 2012].

Lester, S.E. and Halpern, B.S. 2008. Biological responses in marine no-take reserves versus partially protected areas. *Marine Ecology-Progress Series*. 367: 49-56.

Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Aíramé, S. and Warner, R.R. 2009. Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*. 384: 33-46.

Provincial Marine Protected Areas

Coastal Resources Mapping. 2012. Protected Area Summary Tables. Generated for the Marine Planning Process (MaPP) for the North Pacific Coast, 12 May 2012. Available from MaPP.

Fisheries and Oceans Canada. 2011. National framework for Canada's Network of Marine Protected Areas. Government of Canada, Ottawa. 31 pp. Available from <http://www.dfo-mpo.gc.ca/oceans/publications/dmpaf-eczpm/docs/framework-cadre2011-eng.pdf> [accessed 26 May 2012].

Government of BC. 1996. *Ecological Reserve Act*.

Government of BC. 1996. *Park Act*.

Government of Canada and Province of British Columbia. 1998. Marine Protected Areas Discussion Paper. Available from: <http://archive.ilmb.gov.bc.ca/cis/rpts/disppap.htm> [accessed 26 May 2012].

Government of Canada and Province of British Columbia. 2011. DRAFT Canada – British Columbia Marine Protected Area Network Strategy. Available from: <http://www.mccpacific.org/MPA-DraftCanadaBCStrat-May27.pdf> [accessed 27 May 2012].

Province of British Columbia. 2009. Marine Protected Areas. Available from: <http://www.env.gov.bc.ca/omfd/ocean-resources/mpa.html> [accessed 26 May 2012].

Shellfish Aquaculture

Barton, A., Hales, B., Waldbusser, G.G., Langdon, C., and Freely, R.A. 2012. The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects. *Limnology and Oceanography*, 57 (3): 698 DOI: [10.4319/lo.2012.57.3.0698](https://doi.org/10.4319/lo.2012.57.3.0698)

British Columbia. Ministry of Agriculture. 2011. British Columbia Seafood Industry 2010 year in Review. 216 pp. <http://www.bcseafood.ca/PDFs/YIR-2010.pdf>

- Edwin Blewett & Associates Inc. and Nelson Brothers Fisheries Ltd. 2003. North Islands Straits Shellfish Aquaculture Barriers & Constraints Study. April 2003. 40 pp.
http://www.agf.gov.bc.ca/fisheries/cabinet/NI_Shellfish_Aq_Barriers_Constraints.pdf
- Fisheries and Oceans Canada. 2011. Pacific Region Integrated Management of Aquaculture Plan—Shellfish Interim. November 29, 2011. 36 pp.
- Government of Canada. Fisheries and Oceans Canada. Pacific Region. 2011. Pacific Region Integrated Management of Aquaculture Plan Shellfish Interim Nov 29, 2011. 36 pp.
http://www.agf.gov.bc.ca/fisheries/cabinet/NI_Shellfish_Aq_Barriers_Constraints.pdf
- Kingzett Professional Services Ltd. and AXYS Environmental Consulting Ltd. 2002. First nations Shellfish Aquaculture Regional Business Strategy: B.C. Central and Northern Coast. Prepared for Aboriginal Relations and Economic measures, Land and Water British Columbia Inc. November 2002. 254 pp.
http://www.agf.gov.bc.ca/fisheries/cabinet/FN_Shellfish_Aquaculture_North_Coast_Strategy.pdf
- Snuneymuxw First Nation. 2008. Traditional Sea Foods Balancing Health Benefits with Pollution Risks. Final Report. Proceedings of a Workshop at Malaspina Campus, Vancouver Island University. April 5 2008. <http://www.indigenousfoodsvi.ca/wp-content/uploads/2011/08/2008-final-report.pdf>

RESOURCES COMPILED FOR BASE CASE DEVELOPMENT

Ecological Integrity

- Central Coast First Nations. Draft Central Coast First Nations Marine Use Plan. Executive Summary. Pp 32.
- Christensen, V., Lai, S., Palomares, M.L.D., Zeller, D. and Pauly, D (eds). 2011. The state of biodiversity and fisheries in regional seas. Fisheries Centre Research Reports. 19(3): 1-85.
- Côté, I.M., Dodson, J.J., Fleming, I.A., Hutchings, J.A., Jennings, S., Mantua, N.J., Peterman, R.M., Riddell, B.E., Weaver, A.J. and VanderZwagg, D.L. 2012. Sustaining Canadian Marine Biodiversity. An expert panel report on sustaining Canada's marine biodiversity: responding to the challenges posed by climate change, fisheries, and aquaculture. The Royal Society of Canada: The Academies of Arts, Humanities and Sciences of Canada. Pp 316.
- Fisheries and Oceans Canada. 2005. Guidelines on evaluating ecosystem overviews and assessments: necessary documentation. Canadian Science Advisory Secretariat Science Advisory Report. 2005/026: 1-10.
- Fisheries and Oceans Canada. 2010. State of the Pacific Ocean 2009. Canadian Science Advisory Secretariat Science Advisory Report. 2010/034: 1-23.
- Fisheries and Oceans Canada. 2011. State of the Pacific Ocean 2010. Canadian Science Advisory Secretariat Science Advisory Report. 2011/032: 1-23.
- Gregg, E.J and Coatta, R. 2008. Environmental data for the eastern North Pacific and Bering Sea. Fisheries Centre Research Reports. 16(6): 1-79.

- Haggarty, D.R., McCorquodale, B., Johannessen, D.I., Levings, C.D and Ross, P.S. 2003. Marine environmental quality in the central coast of British Columbia, Canada: a review of contaminant sources, types and risks. Canadian Technical Report of Fisheries and Aquatic Sciences. 2507: 1-165.
- Irvine, J.R. and Riddell, B.E. 2007. Salmon as status indicators for North Pacific Ecosystems. North Pacific Anadromous Fish Commission Bulletin. 4: 285-287.
- Jamieson, G.S and McCorquodale, B. 2004. Proceedings of the Central Coast Marine Environmental Quality Indicators Workshop. Canadian Science Advisory Secretariat. 2004/025: 1-58.
- Johannessen, D. 2005. Oceanographic overview of the Queen Charlotte Strait Area. Prepared for Ecostrust Canada and Namgi's First Nations. Pp 60.
- Levi, T., Darimont, C.T., MacDuffee, M., Mangel, M., Paquet, P. and Wilmers, C.C. 2012. Using Grizzly Bears to assess harvest-ecosystem tradeoffs in salmon fisheries. Public Library of Science Biology.10(4): 1-10.
- Rubus EcoScience Alliance. 2007. Indicators of human well-being for the central and north coast: review and recommendations for Schedules C and G. A report commissioned by the Ecosystem based Management Working Group. Pp 129.
- Ruskule, A., Kuris, M., Leiputė, G., Vetemaa, M. and Zableckis, S. 2009. See the Baltic Sea: unique assets we share. Baltic Environmental Forum, Latvia. p 82.

Human Wellbeing

- Central Coast First Nations. Central Coast First Nations Marine Use Plan. Executive Summary. Pp 32.
- Haggarty, D.R., McCorquodale, B., Johannessen, D.I., Levings, C.D and Ross, P.S. 2003. Marine environmental quality in the central coast of British Columbia, Canada: a review of contaminant sources, types and risks. Canadian Technical Report of Fisheries and Aquatic Sciences. 2507: 1-165.
- Helliwell, J., Layard, R. and Sachs, J. (eds). 2012. World Happiness Report. The Earth Institute, Canadian Institute for Advanced Research and Centre for Economic Performance.. Pp 158.
- Impact Assessment, Inc. 2011. Social and economic assessment of major oil spill litigation and settlement. Final Technical Report. Prepared for the US Department of the Interior. Pp 178.
- Indian and Northern Affairs Canada. 2010. First Nation and Inuit community well-being: describing historical trends (1981-2006). Strategic Research and Analysis Directorate. Pp 27.
- MacConnachie, S., Hillier, J. and Butterfield, S. 2007. Marine use analysis of the Pacific North Coast Integrated Management Area. Canadian Technical Report of Fisheries and Aquatic Sciences. 2677: 1-198.
- Robinson Consulting and Associates Ltd. 2012. Socio-economic and cultural overview and assessment report for the Pacific North Coast Integrated Management Area. Prepared for Fisheries and Oceans Canada. Pp 230.
- Rubus EcoScience Alliance. 2007. Indicators of human well-being for the central and north coast: review and recommendations for Schedules C and G. A report commissioned by the Ecosystem based Management Working Group. Pp 129.

- Smith, R. 2004. Review of well-being assessment of North and Central Coastal British Columbia DRAFT Report dated 27 January 2004. Environment Canada. Pp 10.
- Terry, E., Boothemp, J. and Komori, V. 2000. Central Coast land and coastal resource management plan. Socio-economic and environmental/marine base case: final report. Province of British Columbia. Pp 300.
- The Aboriginal Aquaculture Association. Year Unknown. Aquaculture development: an economic opportunity for BC Coastal First Nations. Available from:
www.cohencommission.ca/DownloadExhibit.php?ExhibitID=1917 [accessed 29 May 2012].