


Western balsam bark beetle,
Dryocoetes confusus, and subalpine-fir
forests in southern B.C.

Lorraine Maclauchlan

Min. of Forests, Lands & Natural Resource
Operations
Thompson Okanagan Region
Kamloops, B.C.



Outline

- *Dryocoetes confusus* life history tidbits
- Ecosystems at risk - subalpine-fir succession and *D. confusus* attack patterns
- Impacts observed in past decade - baseline data???
- Climate change and possible effects on *D. confusus* and subalpine-fir

When does chronic become epidemic?

Is mortality from *D. confusus* attack increasing?

Are the periods of high in-stand mortality longer?





Subalpine fir, *Abies lasiocarpa*, is well adapted to cold, high-elevation ecosystems ($\pm 1,600$ m in B.C.)

Succession is a slow process in these subalpine-fir ecosystems, with single or small groups of tree dying and falling out of the stand

Spruce or subalpine-fir will regenerate in gaps

± 300 years, catastrophic fires may destroy large areas of subalpine-fir forests (*rare event*)

Females construct egg galleries in phloem

Brood 1



Late summer – some adults re-emerge (mostly ♀)



Larvae feed in phloem

Dryocoetes confusus life cycle

Overwinter as late instar larvae

Require cold snap

Males initiate attack in June – attracting and mating multiple females



1st year (June – September)



Brood 1

Overwinter as adults

Summer

Females within galleries continue extending galleries, expelling frass & sawdust out of nuptial chamber in **spring** & may lay second brood.



Larvae continue feeding; pupate; and, transform to adults
Spring-early summer

Brood 2

Overwinter as larvae



2nd year (May – September)



New adults emerge from tree,
seek new host and initiate attack



Brood 1



Brood 2
Overwinter as
adults



Larvae continue feeding;
pupate; transform to adults

3rd year (May – September)



Brood 2

Adults emerge from tree & seek new host and mates


Trees fade to grey and loose needles over time



4th year (June)

Stand Dynamics

- some subalpine-fir die
- some subalpine-fir blow down (live & dead trees) creating gaps in the forest canopy



What is the sequence and magnitude of subalpine-fir mortality and blow down as it relates to insect colonization and stand succession?

The annual spatial signature of *Dryocoetes confusus* is typically small and disperse. This coupled with cyclical, severe pockets of attack, slowly converts old subalpine-fir stands to younger stands.



The studies I describe build upon others work:


Art Stock

Leroy Harder

Kathy Bleiker



Ha affected by western balsam bark beetle in southern B.C. (1987-2011)




Percent attack by *Dryocoetes* (reds and greys) in all stands surveyed by air, grouped by ecosystem ($\pm$ SE)



65,767 ha surveyed (805 polygons)

- Highest mortality (56%) observed in an ESSFwc2 polygon.
- Mortality no different among age classes 7, 8, & 9 (ANOVA, $P < 0.05$),
- Age class 6 significantly lower mortality.


Highest and most consistent levels of mortality in ESSF (740 polygons surveyed)






Location, ecosystem and phase of ten 1 hectare plots - established 1998- 2002


Location	BEC	Phase	Year established
Martin Cr.	ESSFwc	early	2000
Torrent Cr.	ESSFwc	mid- to late-	1998
Cherry Ridge	ESSFwc	mid- to late-	1998
Sicamous Cr.	ESSFwc	mid- to late-	1998
Scotch Creek	ESSFwc	mid- to late-	2002
Home Lk-1	ESSFxc	early to mid-	1999
Home Lk-2	ESSFxc	early to mid-	1999
Buck Mtn.	ESSFxc	early to mid-	1999
Spius Cr-1	ESSFmw	early	2002
Spius Cr-2	ESSFmw	early	2002




Live subalpine fir per hectare at plot establishment & final assessment in three ESSF subzones




Dryocoetes-attacked subalpine-fir per hectare at plot establishment & final assessment




Annual blowdown rate of live and dead subalpine-fir (sph) in seven PSP's (plot establishment-2004)



Stems per hectare of down subalpine fir (since plot establishment) noting presence/absence of *Dryocoetes* attack





Change of sub-alpine fir status over 10 year interval (average no. subalpine-fir/ha $\pm$ SE) in 10 PSP's




Hectares affected by
Dryocoetes as mapped in aerial
overview survey: 1996-2011

Okanagan Shuswap




2 to >3% average annual
attack rate in both plots

Scotch Creek



Home Lake-2




Hectares affected by *Dryocoetes* as mapped in aerial overview survey: 1996-2011


Cascades





Spius Cr-1




Spius Cr-2




Mean diameters of
subalpine-fir (from PSP's):
live; attacked by *WBBB*:
and, dead from other
causes (2004 data).






Current attack v-plot



Reds v-plot





Greys v-plot





Martin Creek Plot

Snags v-plot









Pissodes galleries in subalpine-fir

A *Pissodes* found killing trees -
alone and in combination with
Dryocoetes

e.g. Buck Plot - 2009
assessment 32 of 298 trees
killed were *Pissodes* (10%)



Pissodes adult



Summary

- Subalpine-fir stands are rapidly losing volume and succession is being driven by *D. confusus* attack
- Stands >70 years are suitable and over a 10-20 years period undergo *select harvest* by WBBB
- Moderating climate could allow *D. confusus* and associated insects greater success as host trees become stressed
- impacts can be severe after only 10 years of *D. confusus* activity in a stand