


Uncoupling of seed availability and regeneration processes in endangered limber pine populations of Alberta

Vern Peters

The King's University College

Edmonton Alberta



Conservation Issue:

- Mortality exceeds regeneration
- “Endangered species” – 2009 in Alberta



Disease

Limber Pine Ecology

Mutualism

White Pine Blister Rust

Clark's Nutcracker

More Predation

Cone Predation

Grizzly Bears

Red squirrels



-

+

-

Objectives

Seed Predation



- To determine whether cone (seed) escape differs in light versus severely infested landscapes
 - Hyp: More seeds escape in light infections

Regeneration

- To determine whether recent regeneration differs in light versus severely infected landscapes
 - Hyp: More regeneration occurs in light infections



Existing Limber pine field sites with stand inventory, disease, and prior cone production data.



Alberta, Canada

SAMPLING DESIGN – LANDSCAPE LEVEL CONE REMOVAL

- 2 metapopulations separated by 400 km with 8-9 limber pine populations in each

High WPBR - Porcupine Hills




2 Stand Types
(Limber Pine and Douglas Fir Mix)

Low WPBR - Kootenay Plains



3 Stand Types - Limber Pine,
Spruce Mix, Lodgepole Pine Mix


WPBR increase in Northern Study Areas



Data from
Smith, C. Parks Canada
Langor, D., CFS
WPBR Survey


Figure 1: Change in WPBR levels in two metapopulations in Alberta over time ((n =3-4)

Basal Area of Dead and Live Limber Pine (n = 8–9 populations per study area)



SAMPLING DESIGN – POPULATION

- 17 Populations Sampled
- Cone Removal by squirrel
- Low (2008, 2009) and High (2010) cone years
- 4 plots, 10 trees/plot
- Squirrel Abundance
- 6 Midden transects – 3 km (60 ha)/population
- Regeneration - ■ 12 plots per stand



Methodology – Cone Production

Pre-dispersal: Easy



Post-dispersal: Tough




Risk – underestimating cone removal

Methodology – Midden Use

Active Middens vs. feeding stations




Interannual variation in cone production (n = 9 and 8 populations, respectively)



Squirrel Abundance in low versus high BR study areas


(+/- std. error, n = 9 and 8 populations, respectively)



More Squirrels occurred in the low WPBR region


Cone predation amongst years

(n = 9 and 8 populations, respectively)



Interannual Cone escape

(n = 9 and 8 populations, respectively)



Nuts, Nuts, Get 'em!

Squirrel predation and WPBR

Tree

- Squirrels reduce the amount of seed available for dispersers
- Interannual differences in cone availability are more important than WPBR levels between landscapes
- Mast years are likely necessary to attract nutcrackers (1000 cones/ha)



Likely reduces gravity-dispersed regeneration

Regeneration



Fundamental Question in Seed Ecology

■ Is regeneration limited by:

1) seed availability

or

2) substrate availability

or

3) disease

- Production

- Predation

- Dispersal


- Seedbed

- Disturbance

- Succession


Regeneration and WPBR

(n = 9 and 8 populations, respectively)



Why is regeneration so low where more seed is available (i.e. escaping squirrels?)


Regeneration and Substrate Availability



- Substrates may be limiting regeneration at high WPBR

Timing of Limber Pine Regeneration

(n = 9 and 8 populations, respectively)




-Substrates are still available

-Grazing effects?
-Recruitment rates are lower


Size Structure of Seedlings

(n = 9, and 8 populations, respectively)

a) Low WPBR



b) High WPBR



Proportionally fewer trees in larger size classes at Low WPBR sites

- Seedlings are “recruiting” into larger size classes in both study areas

Preliminary Conclusions

- Seed is primarily available in select years.
 - Interannual availability is more important than differences in availability relative to WPBR
- Seed available for nutcrackers is limited in high- and low-WPBR landscapes.
- Regeneration is inadequate in high WPBR landscapes
 - Regeneration is uncoupled from seed production processes (more seed but less suitable substrate)

Acknowledgements:

Funding

- NSERC-USRA
- AB Tourism, Parks, and Protected Areas
- AB Conservation Association

WPBR data

- Parks Canada
- Canadian Forest Service



Questions??

