

**PESTICIDE USE**  
**IN BRITISH COLUMBIA FORESTRY**  
**1990**

by  
P. Humphreys  
Silviculture Branch

British Columbia Ministry of Forests  
Victoria, British Columbia  
Forest Service Internal Report  
PM-PB-79

Not for Publication

July 1991

(The contents of this report may not be cited  
in whole or part without the written approval of the  
Director, Silviculture Branch, Ministry of Forests  
31 Bastion Square, Victoria, B. C. V8W 3E7



## TABLE OF CONTENTS

|                                                            | <u>PAGE</u> |
|------------------------------------------------------------|-------------|
| LIST OF FIGURES AND TABLES.....                            | ii          |
| FORWARD.....                                               | iv          |
| INTRODUCTION.....                                          | 1           |
| RESULTS OF THE 1990 B.C. FORESTRY PESTICIDE USE SURVEY.... | 2           |
| REGIONAL SUMMARIES                                         |             |
| VANCOUVER.....                                             | 12          |
| PRINCE RUPERT.....                                         | 23          |
| PRINCE GEORGE.....                                         | 30          |
| KAMLOOPS.....                                              | 40          |
| NELSON.....                                                | 49          |
| CARIBOO.....                                               | 55          |
| DISTRICT SUMMARIES                                         |             |
| VANCOUVER.....                                             | 14          |
| PRINCE RUPERT.....                                         | 25          |
| KAMLOOPS.....                                              | 32          |
| NELSON.....                                                | 42          |
| CARIBOO.....                                               | 57          |
| DISCUSSION AND CONCLUSIONS.....                            | 69          |
| ACKNOWLEDGEMENTS.....                                      | 70          |

LIST OF TABLES AND FIGURESTABLES

| <u>NUMBER</u> |                                                                                                             | <u>PAGE</u> |
|---------------|-------------------------------------------------------------------------------------------------------------|-------------|
| I             | The Status of 1990 Pesticide Use Applications and Permits issued by the Pesticide Control Branch.....       | 3           |
| II            | The Status of Appealed 1990 Pesticide Use Permits as Determined by the B.C. Environmental Appeal Board..... | 4           |
| III           | Provincial Summary of Area (ha) Treated and Active Ingredient (kg a.i.) Used by Pesticide, 1990.....        | 5           |
| IV            | Regional Summary of Area (ha) Treated by Pesticide, 1990.....                                               | 8           |
| V             | Regional Summary of Area (ha) Treated by Control Purpose, 1990.....                                         | 9           |
| VI            | Regional Summary of Area (ha) Treated by Method of Application, 1990.....                                   | 10          |
| VII           | Provincial Summary of Area (ha) Treated by Control Purpose, 1990.....                                       | 60          |
| VIII          | Provincial Summary of Area (ha) Treated by Method of Application, 1990.....                                 | 63          |
| IX            | Area (ha) Treated with Herbicides and Insecticides for Forestry Purposes, 1980-1990.....                    | 66          |

| <u>NUMBER</u> | <u>FIGURES</u>                                                                                 | <u>PAGE</u> |
|---------------|------------------------------------------------------------------------------------------------|-------------|
| 1.            | Pesticide Use in Percent by Area (ha) Treated, 1990...                                         | 6           |
| 2.            | Pesticide Use in Percent by Active Ingredient<br>(kg a.i.) .....                               | 7           |
| 3.            | Provincial Summary of Area (ha) Treated with<br>Pesticides by Forest Region, 1990.....         | 11          |
| 4.            | Provincial Summary of Area (ha) Treated with<br>Pesticides by Control Purpose, 1990.....       | 62          |
| 5.            | Provincial Summary of Area (ha) Treated with<br>Pesticides by Method of Application, 1990..... | 65          |
| 6.            | Area (ha) Treated with Herbicides for Forestry<br>Purposes, 1980-1990.....                     | 67          |
| 7.            | Area (ha) Treated with Insecticides for Forestry<br>Purposes, 1980-1990.....                   | 68          |

**FOREWORD**

This report is part of a continuing series initiated in 1981. At that time the B.C. Ministry of Forests (Protection Branch), the Council of Forest Industries of B.C., and the B.C. Ministry of Environment (Pesticide Control Branch) began a cooperative project to collect data on forestry pesticide use in British Columbia. Information contained in these reports is gathered from all forestry pesticide permit holders, on an annual basis, by the Pesticide Control Branch, and represents only those permits that were active in 1990. Data includes quantities (kg of active ingredient) of each pesticide applied and total area (ha) treated. Area treated is broken down by control purpose and method of application. Also included in this report are forest region and district summaries of pesticide use, as well as herbicide and insecticide use trends from 1980 - 1990.

## INTRODUCTION

This report is part of a continuing series that began in 1981. At that time a cooperative project was initiated by the B.C. Ministry of Forests (Protection Branch), the Council of Forest Industries of B.C., and the B.C. Ministry of Environment (Pesticide Control Branch) to establish annual records of forestry pesticide use on crown land in British Columbia.

In 1983, the Pesticide Control Branch initiated a requirement of all pesticide use permit holders to submit follow-up reports on a calendar year basis. This data is summarized in tables and graphs on the following pages and includes the following information:<sup>1</sup>

1. Numbers of 1990 forestry permit applications submitted; denied; cancelled and appealed.
2. Environmental Appeal Board decisions on all appealed permits.
3. Provincial summary of area (ha) treated and active ingredient (kg a.i.) used by individual pesticide.
4. Forest region and district summaries of area (ha) treated by pesticide, control purpose and method of application.
5. Provincial summaries of area (ha) treated by control purpose and method of application.
6. Area (ha) of forest land treated with herbicides and insecticides, 1980-1990.

The above information is also divided by the user - i.e., B.C. Forest Service or forest industry.

<sup>1</sup>This report was based on all follow-up reports received by April 1991.

Totals in hectares may differ slightly in each table and graph because of rounding. Totals in forest region summaries are slightly lower than provincial summaries because provincial summaries reflect permits that were used to apply pesticides in several regions.

### RESULTS OF THE 1990 SURVEY

Pesticide Control Branch reviewed 335 forestry pesticide use applications in 1990 and issued 321 permits (Table I).

Sixteen permits were appealed in 1990 and Environmental Appeal Board decisions were as follows: nine supported and, seven amended (Table II).

A total of 57 662 hectares (ha) of B.C. forest and range (Crown) land were treated with pesticides in 1990. The B.C. Forest Service applied pesticides to 21 058 ha (36%) and forest industry to 36 603 ha (64%).

The herbicides glyphosate and picloram were the most widely used products. Glyphosate was applied to 48 976 ha, 85% of the total area treated in the province in 1990, and picloram to 6 476 ha. These were followed by 2,4-D (1 052 ha), MSMA (431 ha), *Bacillus thuringiensis* var. *kurstaki* (B.t.k.) (360 ha) and hexazinone (308 ha) (Table III, Figure 1).

Quantities of pesticides applied in kilograms of active ingredient (kg a.i.) are shown in Table III and Figure 2.

Summaries of pesticide use by individual forest regions and districts are shown on pages 12 - 59.

The majority of pesticide applications occurred in the Prince George Forest Region (27 143 ha), followed by Vancouver (9 782 ha), Cariboo (5 092 ha) and Prince Rupert (4 778 ha) (Table IV, Figure 3).

Herbicide applications for the purpose of controlling vegetation (i.e. crop establishment (brushing), crop growth enhancement (conifer release)) remain the most common type of forestry operation involving pesticides.

Herbicides were applied to 43 569 ha for brushing and conifer release operations and to 4 389 ha to prepare harvested sites for planting. Pre-harvest and site rehabilitation treatments were conducted on 1 210 ha, and 425 ha, respectively. Herbicides were also applied to 6 479 ha for range land management (noxious weed control). Insecticides were applied to 845 ha (Table VII, Figure 4).

Pesticides were aerially applied to 39 010 ha of forest land. This area was divided almost equally between helicopter operations and fixed-wing aircraft operations. Tree injection application methods were used over 7 623 ha, while backpack sprayers and vehicle mounted sprayers were used to apply pesticides to 11 029 ha (Tables VIII, Figure 5).

Herbicides were applied to a total of 56 810 ha, of forest and rangeland in 1990. Insecticides were applied to 845 ha (Table IX, Figures 6 and 7).

TABLE I  
THE STATUS OF 1990 PESTICIDE USE APPLICATIONS AND PERMITS  
ISSUED BY THE PESTICIDE CONTROL BRANCH

|                                                                              | B.C. FOREST<br>SERVICE | FOREST<br>INDUSTRY | TOTAL |
|------------------------------------------------------------------------------|------------------------|--------------------|-------|
| APPLICATIONS<br>SUBMITTED                                                    | 157                    | 178                | 335   |
| APPLICATIONS WITHDRAWN<br>OR DENIED                                          | 4                      | 10                 | 14    |
| PERMITS ISSUED<br>(TOTAL)                                                    | 153                    | 168                | 321   |
| PERMITS CANCELLED REVOKED<br>OR SUSPENDED BY THE<br>PESTICIDE CONTROL BRANCH | 0                      | 0                  | 0     |

TABLE II  
THE STATUS OF APPEALED 1990 PESTICIDE USE PERMITS  
AS DETERMINED BY THE B.C. ENVIRONMENTAL APPEAL BOARD

|                                          | B.C. FOREST<br>SERVICE | FOREST<br>INDUSTRY | TOTAL |
|------------------------------------------|------------------------|--------------------|-------|
| PERMITS APPEALED                         | 5                      | 11                 | 16    |
| APPEALED PERMITS<br>CANCELLED/REVOKED    | 0                      | 0                  | 0     |
| APPEALED PERMITS<br>AMENDED <sup>1</sup> | 5                      | 2                  | 7     |
| APPEALED PERMITS<br>SUPPORTED            | 0                      | 9                  | 9     |
| HEARINGS OR DECISIONS<br>PENDING         | 0                      | 0                  | 0     |

<sup>1</sup>Permits extended one year by the Appeal Board are not considered to be amended.


TABLE III  
PROVINCIAL SUMMARY OF AREA (ha)  
TREATED AND ACTIVE INGREDIENT (kg a.i.) USED  
BY PESTICIDE, 1990

| PESTICIDE                  | BC<br>FOREST SERVICE |                     | FOREST<br>INDUSTRY |                     |
|----------------------------|----------------------|---------------------|--------------------|---------------------|
|                            | ha<br>treated        | kg. a.i.<br>applied | ha<br>treated      | kg. a.i.<br>applied |
| 2, 4-D amine               | 90.800               | 436.175             | 937.000            | 551.950             |
| 2, 4-D ester               | 0.000                | 0.000               | 24.000             | 88.400              |
| acephate                   | 0.012                | 0.077               | 0.050              | 0.030               |
| <u>B.t.k.</u> <sup>1</sup> | 0.000                | 0.000               | 360.000            | 900.000             |
| carbaryl                   | 0.840                | 13.440              | 0.000              | 0.000               |
| chlorpyrifos               | 0.760                | 49.440              | 0.000              | 0.000               |
| cyclohexnon                | 16.000               | 70.000              | 0.000              | 0.000               |
| diphacinone                | 0.000                | 0.000               | 0.044              | 0.005               |
| glyphosate                 | 13 983.642           | 22 356.802          | 34 992.979         | 57 069.120          |
| hexazinone                 | 27.300               | 83.686              | 280.800            | 293.600             |
| metalazyl                  | 2.370                | 2.023               | 0.000              | 0.000               |
| mineral oil                | 0.160                | 80.000              | 0.000              | 0.000               |
| MSMA                       | 423.913              | 810.200             | 7.020              | 54.980              |
| oxydemeton                 | 0.012                | 0.073               | 0.000              | 0.000               |
| picloram                   | 6 476.172            | 3 529.725           | 0.393              | 0.220               |
| pirmicarb                  | 0.005                | 0.093               | 0.000              | 0.000               |
| putrid eggs                | 0.000                | 0.000               | 0.800              | 0.130               |
| treademefon                | 0.350                | 0.080               | 0.000              | 0.000               |
| triclopyr                  | 0.100                | 0.100               | 0.000              | 0.000               |
| verbenone                  | 36.000               | 29.440              | 0.000              | 0.000               |

---

<sup>1</sup>*Bacillus Thuringiensis* var. *kurstaki*


# PESTICIDE USE<sup>1</sup> IN PERCENT (ha) TREATED, 1990



<sup>1</sup> B.C. Forest Service & Industry Combined

<sup>2</sup> *Bacillus thuringiensis* var. *kurstaki*

# PESTICIDE USE IN PERCENT ACTIVE INGREDIENT (kg a.i.) APPLIED, 1990<sup>1</sup>



<sup>1</sup> B.C. Forest Service & Industry Combined

<sup>2</sup> *Bacillus thuringiensis* var. *kurstaki*

**TABLE IV**  
**REGIONAL SUMMARY OF AREA (ha) TREATED**  
**BY PESTICIDE, 1990**  
**BCFS AND FOREST INDUSTRY**

| <b>FOREST REGION</b> | <b>glyphosate</b> | <b>2, 4-D</b>  | <b>hexazinone</b> | <b>picloram</b>  | <b><u>B.t.k.</u></b> | <b>Other</b>   | <b>Totals</b>     |
|----------------------|-------------------|----------------|-------------------|------------------|----------------------|----------------|-------------------|
| VANCOUVER            | 8 458.128         | 937.000        | 0.000             | 0.010            | 360.000              | 26.644         | 9 781.782         |
| PRINCE RUPERT        | 4 750.860         | 0.000          | 0.000             | 13.380           | 0.000                | 13.860         | 4 778.100         |
| PRINCE GEORGE        | 26 727.412        | 0.000          | 51.600            | 37.221           | 0.000                | 326.973        | 27 143.206        |
| KAMLOOPS             | 1 551.116         | 0.000          | 0.000             | 6 334.063        | 0.000                | 94.360         | 7 979.539         |
| NELSON               | 593.930           | 0.000          | 0.000             | 91.891           | 0.000                | 34.070         | 719.891           |
| CARIBOO              | 4 834.765         | 0.000          | 256.500           | 0.000            | 0.000                | 1.010          | 5 092.275         |
| <b>TOTALS:</b>       | <b>46 916.211</b> | <b>937.000</b> | <b>308.100</b>    | <b>6 476.565</b> | <b>360.000</b>       | <b>496.917</b> | <b>55 494.793</b> |
| <b>(%)</b>           | <b>84.54</b>      | <b>1.69</b>    | <b>0.56</b>       | <b>11.67</b>     | <b>0.65</b>          | <b>0.90</b>    | <b>100.00</b>     |


**TABLE V**  
**REGIONAL SUMMARY OF AREA (ha) TREATED**  
**BY CONTROL PURPOSE, 1990**  
**BCFS AND FOREST INDUSTRY**

| <b>FOREST REGION</b> | <b>VEGETATION<br/>MGMNT</b> | <b>INSECT<br/>MGMNT</b> | <b>RANGE<br/>MGMNT</b> | <b>OTHER</b> | <b>TOTAL</b>      |
|----------------------|-----------------------------|-------------------------|------------------------|--------------|-------------------|
| VANCOUVER            | 9 419.238                   | 361.350                 | 0.000                  | 1.194        | 9 781.782         |
| PRINCE RUPERT        | 4 747.740                   | 13.360                  | 13.500                 | 3.500        | 4 778.100         |
| PRINCE GEORGE        | 26 778.590                  | 326.523                 | 38.093                 | 0.000        | 27 143.206        |
| KAMLOOPS             | 1 551.499                   | 93.590                  | 6 333.680              | 0.770        | 7 979.539         |
| NELSON               | 591.930                     | 33.470                  | 93.891                 | 0.600        | 719.891           |
| CARIBOO              | 5 091.265                   | 0.510                   | 0.000                  | 0.500        | 5 092.275         |
| <b>TOTALS:</b>       | <b>48 180.262</b>           | <b>828.803</b>          | <b>6 479.164</b>       | <b>6.564</b> | <b>55 494.793</b> |
| <b>(%)</b>           | <b>86.82</b>                | <b>1.49</b>             | <b>11.68</b>           | <b>0.01</b>  | <b>100.00</b>     |

**TABLE VI**  
**REGIONAL SUMMARY OF AREA (ha)**  
**TREATED BY METHOD OF APPLICATION, 1990**  
**BCFS AND FOREST INDUSTRY**

| <b>FOREST REGION</b> | <b>AERIAL<br/>SPRAY</b> | <b>GROUND<br/>APPLICATION</b> | <b>STEM<br/>INJECTION</b> | <b>TOTALS</b>     |
|----------------------|-------------------------|-------------------------------|---------------------------|-------------------|
| VANCOUVER            | 3 694.220               | 1 492.564                     | 4 594.998                 | 9 781.782         |
| PRINCE RUPERT        | 3 699.540               | 476.700                       | 601.860                   | 4 778.100         |
| PRINCE GEORGE        | 24 513.900              | 990.973                       | 1,638.333                 | 27 43.206         |
| KAMLOOPS             | 460.500                 | 7 086.230                     | 432.809                   | 7 979.539         |
| NELSON               | 415.630                 | 103.741                       | 200.520                   | 719.891           |
| CARIBOO              | 4 356.065               | 735.700                       | 0.510                     | 5 092.275         |
| <b>TOTALS:</b>       | <b>37 139.855</b>       | <b>10 885.908</b>             | <b>7 469.030</b>          | <b>55 494.793</b> |
| (%)                  | 66.92                   | 19.62                         | 13.46                     | 100.00            |

# AREA TREATED WITH PESTICIDES<sup>1</sup> PROVINCIAL SUMMARY BY FOREST REGION 1990



<sup>1</sup> B.C. Forest Service & Industry Combined

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                                          | TOTAL (ha)<br>TREATED | PESTICIDES                              | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                          |                           |
|--------------------------------------------------|-----------------------|-----------------------------------------|---------------------------------------------|--------------------------|---------------------------|
|                                                  |                       |                                         | AERIAL<br>SPRAY                             | GROUND<br>APPLIC.        | STEM<br>INJECTION         |
| <b>REGION: VANCOUVER<br/>B.C. FOREST SERVICE</b> |                       |                                         |                                             |                          |                           |
| FOREST BRUSHING <sup>1</sup>                     | 1 310.570             | glyphosate                              | 1 004.830                                   | 305.740                  | 0.000                     |
| FOREST CONIFER<br>RELEASE <sup>2</sup>           | 895.600               | 2, 4-D amine<br>glyphosate<br>triclopyr | 0.000<br>137.500<br>0.000                   | 0.000<br>35.400<br>0.100 | 0.000<br>722.600<br>0.000 |
| FOREST NURSERY<br>FUNGUS CONTROL                 | 0.350                 | triademefon                             | 0.000                                       | 0.350                    | 0.000                     |
| FOREST SITE<br>PREPARATION                       | 36.000                | glyphosate                              | 14.000                                      | 22.000                   | 0.000                     |
| FOREST SITE<br>REHABILITATION                    | 66.800                | glyphosate                              | 66.800                                      | 0.000                    | 0.000                     |
| <b>SUB TOTALS:</b>                               | <b>2 309.320</b>      |                                         | <b>1 223.130</b>                            | <b>363.590</b>           | <b>722.600</b>            |

REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY

| PURPOSE                                | TOTAL ha<br>TREATED | PESTICIDES    | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------------|---------------------|---------------|---------------------------------------------|-------------------|-------------------|
|                                        |                     |               | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| REGION: VANCOUVER<br>FOREST INDUSTRY   |                     |               |                                             |                   |                   |
| FOREST BRUSHING <sup>1</sup>           |                     | 2, 4-D amine  | 0.000                                       | 0.000             | 772.000           |
|                                        |                     | glyphosate    | 1 388.140                                   | 163.200           | 2 353.000         |
|                                        |                     | picloram      | 0.000                                       | 0.000             | 0.010             |
|                                        | 4 676.350           |               |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE <sup>2</sup> |                     | 2, 4-D amine  | 0.000                                       | 0.000             | 136.000           |
|                                        |                     | glyphosate    | 340.350                                     | 496.750           | 611.338           |
|                                        | 1 584.438           |               |                                             |                   |                   |
| FOREST INSECT<br>CONTROL               |                     | acephate      | 0.000                                       | 0.000             | 0.050             |
|                                        |                     | <u>B.t.k.</u> | 360.000                                     | 0.000             | 0.000             |
|                                        |                     | MSMA          | 0.000                                       | 1.300             | 0.000             |
|                                        | 361.350             |               |                                             |                   |                   |
| FOREST RIGHT-OF WAY<br>MAINIENANCE     |                     | 2, 4-D amine  | 0.000                                       | 29.000            | 0.000             |
|                                        |                     | 2, 4-D ester  | 0.000                                       | 24.000            | 0.000             |
|                                        |                     | glyphosate    | 0.000                                       | 335.350           | 0.000             |
|                                        | 388.350             |               |                                             |                   |                   |
| FOREST SITE<br>PREPARATION             |                     | glyphosate    | 382.600                                     | 78.530            | 0.000             |
|                                        | 461.130             |               |                                             |                   |                   |
| FOREST VERTEBRATE<br>CONTROL           |                     | diphacinone   | 0.000                                       | 0.044             | 0.000             |
|                                        |                     | putrid eggs   | 0.000                                       | 0.800             | 0.000             |
|                                        | 0.844               |               |                                             |                   |                   |
| <hr/>                                  |                     |               |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>              | <b>7 472.462</b>    |               | <b>2 471.090</b>                            | <b>1 128.974</b>  | <b>3 872.398</b>  |
| <hr/>                                  |                     |               |                                             |                   |                   |
| <b><u>TOTALS:</u></b>                  | <b>9 781.782</b>    |               | <b>3 694.220</b>                            | <b>1 492.564</b>  | <b>4 594.998</b>  |

<sup>1</sup>Throughout this report Forest Brushing refers to treatments that ensure tree crop establishment and survival.

<sup>2</sup>Throughout this report Forest Conifer Release refers to treatments that enhance the growth of the established tree crop.

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: CHILLIWACK**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES             | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------|---------------------|------------------------|---------------------------------------------|-------------------|-------------------|
|                                    |                     |                        | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |                        |                                             |                   |                   |
| FOREST BRUSHING                    | 582.580             | glyphosate             | 580.380                                     | 2.200             | 0.000             |
| FOREST CONIFER<br>RELEASE          | 30.600              | glyphosate             | 0.000                                       | 0.000             | 30.600            |
| FOREST SITE<br>PREPARATION         | 14.000              | glyphosate             | 14.000                                      | 0.000             | 0.000             |
| FOREST SITE<br>REHABILITATION      | 66.800              | glyphosate             | 66.800                                      | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>                 | 693.980             |                        | 661.180                                     | 2.200             | 30.600            |
| FOREST INDUSTRY                    |                     |                        |                                             |                   |                   |
| FOREST BRUSHING                    | 536.850             | glyphosate<br>picloram | 536.840<br>0.000                            | 0.000<br>0.000    | 0.000<br>0.010    |
| FOREST CONIFER<br>RELEASE          | 158.488             | glyphosate             | 156.350                                     | 0.500             | 1.638             |
| FOREST INSECT<br>CONTROL           | 1.300               | MSMA                   | 0.000                                       | 1.300             | 0.000             |
| FOREST RIGHT-OF-WAY<br>MAINTENENCE | 24.000              | 2, 4-D ester           | 0.000                                       | 24.000            | 0.000             |
| FOREST SITE<br>PREPARATION         | 60.100              | glyphosate             | 60.100                                      | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>                 | 780.738             |                        | 753.290                                     | 25.800            | 1.648             |
| <u>TOTALS:</u>                     | 1 474.718           |                        | 1 414.47                                    | 28.000            | 32.248            |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: SQUAMISH**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                  |                     |            |                                             |                   |                   |
| FOREST BRUSHING           | 416.690             | glyphosate | 411.150                                     | 5.540             | 0.000             |
| FOREST CONIFER<br>RELEASE | 13.500              | glyphosate | 13.500                                      | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>        | 430.190             |            | 424.650                                     | 5.540             | 0.000             |
| FOREST INDUSTRY           |                     |            |                                             |                   |                   |
| FOREST BRUSHING           | 134.600             | glyphosate | 131.000                                     | 3.600             | 0.000             |
| FOREST CONIFER<br>RELEASE | 35.000              | glyphosate | 35.000                                      | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>        | 169.600             |            | 166.000                                     | 3.600             | 0.000             |
| <u>TOTALS:</u>            | 599.790             |            | 590.65                                      | 9.14              | 0.00              |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: SUNSHINE COAST**

| PURPOSE                        | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                |                   |
|--------------------------------|---------------------|------------|---------------------------------------------|----------------|-------------------|
|                                |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPL | STEM<br>INJECTION |
| B.C.F.S.                       |                     |            |                                             |                |                   |
| FOREST BRUSHING<br>AND WEEDING | 26.300              | glyphosate | 0.000                                       | 26.300         | 0.000             |
| FOREST CONIFER<br>RELEASE      | 169.000             | glyphosate | 124.000                                     | 0.000          | 45.000            |
| FOREST SITE<br>PREPARATION     | 22.000              | glyphosate | 0.000                                       | 22.000         | 0.000             |
| <b><u>SUB TOTALS:</u></b>      | <b>217.300</b>      |            | <b>124.000</b>                              | <b>48.300</b>  | <b>45.00</b>      |
| FOREST INDUSTRY                |                     |            |                                             |                |                   |
| FOREST BRUSHING<br>AND WEEDING | 112.100             | glyphosate | 112.100                                     | 0.000          | 0.000             |
| FOREST CONIFER<br>RELEASE      | 48.000              | glyphosate | 0.000                                       | 48.000         | 0.000             |
| FOREST SITE<br>PREPARATION     | 72.600              | glyphosate | 0.000                                       | 72.600         | 0.000             |
| <b><u>SUB TOTALS:</u></b>      | <b>232.700</b>      |            | <b>112.100</b>                              | <b>120.600</b> | <b>0.000</b>      |
| <b><u>TOTALS:</u></b>          | <b>450.000</b>      |            | <b>236.100</b>                              | <b>168.900</b> | <b>45.00</b>      |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: DUNCAN**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES                 | AREA (ha) BREAKDOWN BY<br>TREATMENT METHOD |                   |                   |
|------------------------------------|---------------------|----------------------------|--------------------------------------------|-------------------|-------------------|
|                                    |                     |                            | AERIAL<br>SPRAY                            | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |                            |                                            |                   |                   |
| FOREST NURSERY<br>FUNGUS CONTROL   | 0.050               | triademefon                | 0.000                                      | 0.050             | 0.000             |
| <u>SUB TOTALS:</u>                 | 0.050               |                            | 0.000                                      | 0.050             | 0.000             |
| FOREST INDUSTRY                    |                     |                            |                                            |                   |                   |
| FOREST BRUSHING                    | 349.400             | glyphosate                 | 150.000                                    | 106.100           | 93.300            |
| FOREST CONIFER<br>RELEASE          | 105.700             | glyphosate                 | 0.000                                      | 75.700            | 30.000            |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 2.700               | glyphosate                 | 0.000                                      | 2.700             | 0.000             |
| FOREST VERTEBRATE<br>CONTROL       | 0.844               | diphacinone<br>putrid eggs | 0.000<br>0.000                             | 0.044<br>0.800    | 0.000<br>0.000    |
| <u>SUB TOTALS:</u>                 | 458.644             |                            | 150.000                                    | 185.344           | 123.300           |
| <u>TOTALS:</u>                     | 458.694             |                            | 150.000                                    | 185.394           | 123.300           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: PORT ALBERNI**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES  | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------|---------------------|-------------|---------------------------------------------|-------------------|-------------------|
|                                    |                     |             | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |             |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE          | 0.100               | triclopyr   | 0.000                                       | 0.100             | 0.000             |
| <hr/>                              |                     |             |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>          | <b>0.100</b>        |             | <b>0.000</b>                                | <b>0.100</b>      | <b>0.000</b>      |
| <hr/>                              |                     |             |                                             |                   |                   |
| FOREST INDUSTRY                    |                     |             |                                             |                   |                   |
| FOREST BRUSHING                    | 1 554.700           | 2,4-D amine | 0.000                                       | 0.000             | 772.000           |
|                                    |                     | glyphosate  | 28.000                                      | 4.500             | 750.200           |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 19.000              | glyphosate  | 0.000                                       | 19.000            | 0.000             |
| <hr/>                              |                     |             |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>          | <b>1 573.700</b>    |             | <b>28.000</b>                               | <b>23.500</b>     | <b>1 522.200</b>  |
| <hr/>                              |                     |             |                                             |                   |                   |
| <b><u>TOTALS:</u></b>              | <b>1 573.800</b>    |             | <b>28.000</b>                               | <b>23.600</b>     | <b>1 522.200</b>  |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: CAMPBELL RIVER**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES                 | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                    |
|------------------------------------|---------------------|----------------------------|---------------------------------------------|-------------------|--------------------|
|                                    |                     |                            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION  |
| B.C.F.S.                           |                     |                            |                                             |                   |                    |
| FOREST BRUSHING                    | 87.800              | glyphosate                 | 11.500                                      | 76.300            | 0.000              |
| FOREST CONIFER<br>RELEASE          | 35.200              | 2, 4-D amine<br>glyphosate | 0.000<br>0.000                              | 0.000<br>35.200   | 0.000<br>0.000     |
| FOREST NURSERY<br>FUNGUS CONTROL   | 0.300               | treademefon                | 0.000                                       | 0.300             | 0.000              |
| <b><u>SUB TOTALS:</u></b>          | <b>123.300</b>      |                            | <b>11.500</b>                               | <b>111.800</b>    | <b>0.000</b>       |
| FOREST INDUSTRY                    |                     |                            |                                             |                   |                    |
| FOREST BRUSHING                    | 102.900             | glyphosate                 | 100.900                                     | 2.000             | 0.000              |
| FOREST CONIFER<br>RELEASE          | 888.150             | 2, 4-D amine<br>glyphosate | 0.000<br>0.000                              | 0.000<br>255.850  | 136.000<br>496.300 |
| FOREST INSECT<br>CONTROL           | 0.050               | acephate                   | 0.000                                       | 0.000             | 0.050              |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 237.500             | 2, 4-D amine<br>glyphosate | 0.000<br>0.000                              | 29.000<br>208.500 | 0.000<br>0.000     |
| FOREST SITE<br>PREPARATION         | 202.530             | glyphosate                 | 196.600                                     | 5.930             | 0.000              |
| <b><u>SUB TOTALS:</u></b>          | <b>1 431.130</b>    |                            | <b>297.500</b>                              | <b>501.280</b>    | <b>632.350</b>     |
| <b><u>TOTALS:</u></b>              | <b>1 554.43</b>     |                            | <b>309.000</b>                              | <b>613.080</b>    | <b>632.350</b>     |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: PORT MCNEILL**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHOS |                   |                   |
|------------------------------------|---------------------|------------|--------------------------------------------|-------------------|-------------------|
|                                    |                     |            | AERIAL<br>SPRAY                            | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |            |                                            |                   |                   |
| FOREST BRUSHING                    | 1.800               | glyphosate | 1.800                                      | 0.000             | 0.000             |
| FOREST CONIFER<br>RELEASE          | 475.000             | glyphosate | 0.000                                      | 0.000             | 475.000           |
| <u>SUB TOTALS:</u>                 | 476.800             |            | 1.800                                      | 0.000             | 475.000           |
| FOREST INDUSTRY                    |                     |            |                                            |                   |                   |
| FOREST BRUSHING                    | 1 477.600           | glyphosate | 282.600                                    | 47.000            | 1,148.000         |
| FOREST CONIFER<br>RELEASE          | 349.100             | glyphosate | 149.000                                    | 116.700           | 83.400            |
| FOREST INSECT<br>CONTROL           | 360.000             | B.t.k.     | 360.000                                    | 0.000             | 0.000             |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 105.150             | glyphosate | 0.000                                      | 105.150           | 0.000             |
| <u>SUB TOTALS:</u>                 | 2 291.850           |            | 791.600                                    | 268.850           | 1 231.400         |
| <u>TOTALS:</u>                     | 2 768.650           |            | 793.400                                    | 268.850           | 1 706.400         |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: MID-COAST**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |            |                                             |                   |                   |
| FOREST BRUSHING            | 175.800             | glyphosate | 0.000                                       | 175.800           | 0.000             |
| <u>SUB TOTALS:</u>         |                     |            | 0.000                                       | 175.800           | 0.000             |
| FOREST INDUSTRY            |                     |            |                                             |                   |                   |
| FOREST BRUSHING            | 46.700              | glyphosate | 46.700                                      | 0.000             | 0.000             |
| FOREST SITE<br>PREPARATION | 125.900             | glyphosate | 125.900                                     | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>         |                     |            | 172.600                                     | 0.000             | 0.000             |
| <u>TOTALS:</u>             |                     |            | 172.600                                     | 175.800           | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: VANCOUVER**

**DISTRICT: QUEEN CHARLOTTE ISLANDS**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                  |                     |            |                                             |                   |                   |
| FOREST BRUSHING           | 19.600              | glyphosate | 0.000                                       | 19.600            | 0.000             |
| FOREST CONIFER<br>RELEASE | 172.000             | glyphosate | 0.000                                       | 0.000             | 172.000           |
| <u>SUB TOTALS:</u>        | 191.600             |            | 0.000                                       | 19.600            | 172.000           |
| FOREST INDUSTRY           |                     |            |                                             |                   |                   |
| FOREST BRUSHING           | 361.500             | glyphosate | 0.000                                       | 0.000             | 361.500           |
| <u>SUB TOTALS:</u>        | 361.500             |            | 0.000                                       | 0.000             | 361.500           |
| <u>TOTALS:</u>            | 553.100             |            | 0.000                                       | 19.600            | 533.500           |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                                              | TOTAL ha<br>TREATED | PESTICIDES             | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------------------------|---------------------|------------------------|---------------------------------------------|-------------------|-------------------|
|                                                      |                     |                        | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <b>REGION: PRINCE RUPERT<br/>B.C. FOREST SERVICE</b> |                     |                        |                                             |                   |                   |
| FOREST BRUSHING                                      | 617.240             | glyphosate             | 547.740                                     | 69.500            | 0.000             |
| FOREST CONIFER<br>RELEASE                            | 0.200               | glyphosate             | 0.000                                       | 0.200             | 0.000             |
| FOREST FUNGUS<br>CONTROL                             | 3.000               | glyphosate             | 0.000                                       | 0.000             | 3.000             |
| FOREST INSECT<br>CONTROL                             | 13.360              | MSMA                   | 0.000                                       | 0.000             | 13.360            |
| FOREST NURSERY<br>FUNGUS CONTROL                     | 0.500               | metalaxyl              | 0.000                                       | 0.500             | 0.000             |
| FOREST RANGE<br>MANAGEMENT                           | 13.500              | glyphosate<br>picloram | 0.000<br>0.000                              | 0.120<br>13.350   | 0.000<br>0.000    |
| FOREST SITE<br>REHABILITATION                        | 250.700             | glyphosate             | 174.700                                     | 76.000            | 0.000             |
| <b><u>SUB TOTALS:</u></b>                            | <b>898.500</b>      |                        | <b>722.440</b>                              | <b>159.700</b>    | <b>16.360</b>     |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|--------------------------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <b>REGION: PRINCE RUPERT<br/>FOREST INDUSTRY</b> |                     |            |                                             |                   |                   |
| FOREST BRUSHING                                  | 2 382.800           | glyphosate | 1 782.100                                   | 15.200            | 585.500           |
| FOREST CONIFER<br>RELEASE                        | 1 310.000           | glyphosate | 1 023.200                                   | 286.800           | 0.000             |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE               | 15.000              | glyphosate | 0.000                                       | 15.000            | 0.000             |
| FOREST SITE<br>PREPARATION                       | 171.800             | glyphosate | 171.800                                     | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>                        | <b>3 879.600</b>    |            | <b>2 977.100</b>                            | <b>317.000</b>    | <b>585.500</b>    |
| <b><u>TOTALS:</u></b>                            | <b>4 778.100</b>    |            | <b>3 699.540</b>                            | <b>476.700</b>    | <b>601.860</b>    |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE RUPERT**

**DISTRICT: LAKES**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| FOREST INDUSTRY           |                     |            |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE | 90.000              | glyphosate | 90.000                                      | 0.000             | 0.000             |
| <u>TOTALS:</u>            | 90.000              |            | 90.000                                      | 0.000             | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE RUPERT**

**DISTRICT: MORICE**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                    |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |            |                                             |                   |                   |
| FOREST BRUSHING                    | 155.000             | glyphosate | 85.500                                      | 69.500            | 0.000             |
| FOREST INSECT<br>CONTROL           | 11.360              | MSMA       | 0.000                                       | 0.000             | 11.360            |
| FOREST RANGE<br>MANAGEMENT         | 13.500              | glyphosate | 0.000                                       | 0.120             | 0.000             |
|                                    |                     | picloram   | 0.000                                       | 13.380            | 0.000             |
| <b><u>SUB TOTALS:</u></b>          | <b>179.860</b>      |            | <b>85.500</b>                               | <b>83.000</b>     | <b>11.360</b>     |
| FOREST INDUSTRY                    |                     |            |                                             |                   |                   |
| FOREST BRUSHING                    | 750.400             | glyphosate | 735.200                                     | 15.200            | 0.000             |
| FOREST CONIFER<br>RELEASE          | 1 220.000           | glyphosate | 933.200                                     | 286.800           | 0.000             |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 15.000              | glyphosate | 0.000                                       | 15.000            | 0.000             |
| FOREST SITE<br>PREPARATION         | 160.000             | glyphosate | 160.000                                     | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>          | <b>2 145.400</b>    |            | <b>1 828.400</b>                            | <b>317.000</b>    | <b>0.000</b>      |
| <b><u>TOTALS:</u></b>              | <b>2 325.260</b>    |            | <b>1 913.900</b>                            | <b>400.00</b>     | <b>11.360</b>     |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE RUPERT**

**DISTRICT: BULKLEY**

| PURPOSE                       | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|-------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                               |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                      |                     |            |                                             |                   |                   |
| FOREST BRUSHING               | 121.140             | glyphosate | 121.140                                     | 0.000             | 0.000             |
| FOREST FUNGUS<br>CONTROL      | 3.000               | glyphosate | 0.000                                       | 0.000             | 3.000             |
| FOREST INSECT<br>CONTROL      | 2.000               | MSMA       | 0.000                                       | 0.000             | 2.000             |
| FOREST SITE<br>REHABILITATION | 174.700             | glyphosate | 174.700                                     | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>     | <b>300.840</b>      |            | <b>295.840</b>                              | <b>0.000</b>      | <b>5.000</b>      |
| FOREST INDUSTRY               |                     |            |                                             |                   |                   |
| FOREST BRUSHING               | 1 009.200           | glyphosate | 1 009.200                                   | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>     | <b>1 009.200</b>    |            | <b>1 009.200</b>                            | <b>0.000</b>      | <b>0.000</b>      |
| <b><u>TOTALS:</u></b>         | <b>1 310.040</b>    |            | <b>1 305.040</b>                            | <b>0.000</b>      | <b>5.000</b>      |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE RUPERT**

**DISTRICT: KALUM**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                         |                     |            |                                             |                   |                   |
| FOREST BRUSHING                  | 341.100             | glyphosate | 341.100                                     | 0.000             | 0.000             |
| FOREST CONIFER<br>RELEASE        | 0.200               | glyphosate | 0.000                                       | 0.200             | 0.000             |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.500               | metalaxyl  | 0.000                                       | 0.500             | 0.000             |
| <u>TOTALS:</u>                   | 341.800             |            | 341.100                                     | 0.700             | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE RUPERT**

**DISTRICT: NORTH COAST**

| PURPOSE                       | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|-------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                               |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                      |                     |            |                                             |                   |                   |
| FOREST SITE<br>REHABILITATION | 76.000              | glyphosate | 0.000                                       | 76.000            | 0.000             |
| <u>SUB TOTALS:</u>            | 76.000              |            | 0.000                                       | 76.000            | 0.000             |
| FOREST INDUSTRY               |                     |            |                                             |                   |                   |
| FOREST BRUSHING               | 623.200             | glyphosate | 37.700                                      | 0.000             | 585.500           |
| FOREST SITE<br>PREPARATION    | 11.800              | glyphosate | 11.800                                      | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>            | 635.000             |            | 49.500                                      | 0.000             | 585.500           |
| <u>TOTALS:</u>                | 711.000             |            | 49.500                                      | 76.000            | 585.000           |

REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY

| PURPOSE                                              | TOTAL ha<br>TREATED | PESTICIDES  | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------------------------|---------------------|-------------|---------------------------------------------|-------------------|-------------------|
|                                                      |                     |             | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <b>REGION: PRINCE GEORGE<br/>B.C. FOREST SERVICE</b> |                     |             |                                             |                   |                   |
| FOREST BRUSHING                                      |                     | glyphosate  | 7 149.600                                   | 320.220           | 0.000             |
|                                                      |                     | hexazinone  | 27.300                                      | 0.000             | 0.000             |
|                                                      | 7 497.120           |             |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE                            | 1 022.500           | glyphosate  | 686.400                                     | 108.400           | 227.700           |
| FOREST INSECT<br>CONTROL                             |                     | mineral oil | 0.000                                       | 0.160             | 0.000             |
|                                                      |                     | MSMA        | 0.000                                       | 0.000             | 324.593           |
|                                                      | 324.753             |             |                                             |                   |                   |
| FOREST RANGE<br>MANAGEMENT                           |                     | glyphosate  | 0.000                                       | 0.872             | 0.000             |
|                                                      |                     | picloram    | 0.000                                       | 37.051            | 0.170             |
|                                                      | 38.093              |             |                                             |                   |                   |
| FOREST SITE<br>PREPARATION                           | 27.000              | glyphosate  | 27.000                                      | 0.000             | 0.000             |
| FOREST SITE<br>REHABILITATION                        | 108.000             | glyphosate  | 108.000                                     | 0.000             | 0.000             |
| <hr/>                                                |                     |             |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>                            | <b>9 017.466</b>    |             | <b>7 998.300</b>                            | <b>466.703</b>    | <b>552.463</b>    |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                    |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <u>REGION: PRINCE GEORGE</u>       |                     |            |                                             |                   |                   |
| <u>FOREST INDUSTRY</u>             |                     |            |                                             |                   |                   |
| FOREST BRUSHING                    |                     | glyphosate | 9 474.190                                   | 365.400           | 41.500            |
|                                    |                     | hexazinone | 0.000                                       | 24.300            | 0.000             |
|                                    | 9 905.390           |            |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE          | 4 754.710           | glyphosate | 4 751.110                                   | 3.600             | 0.000             |
| FOREST INSECT<br>CONTROL           | 1.770               | MSMA       | 0.000                                       | 0.000             | 1.770             |
| FOREST PRE-HARVEST<br>TREATMENT    | 1 042.600           | glyphosate | 0.000                                       | 0.000             | 1 042.600         |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 130.970             | glyphosate | 0.000                                       | 130.970           | 0.000             |
| FOREST SITE<br>PREPARATION         | 2 290.300           | glyphosate | 2 290.300                                   | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>                 | 18 125.740          |            | 16 515.600                                  | 524.270           | 1 085.870         |
| <u>TOTALS:</u>                     | 27 143.206          |            | 24 513.900                                  | 990.973           | 1 638.333         |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: PRINCE GEORGE**

| PURPOSE                            | TOTAL ha<br>TREATED | PESTICIDES               | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|------------------------------------|---------------------|--------------------------|---------------------------------------------|-------------------|-------------------|
|                                    |                     |                          | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                           |                     |                          |                                             |                   |                   |
| FOREST BRUSHING                    | 1 253.700           | glyphosate               | 1 192.000                                   | 61.700            | 0.000             |
| FOREST CONIFER<br>RELEASE          | 108.400             | glyphosate               | 0.000                                       | 108.400           | 0.000             |
| FOREST INSECT<br>CONTROL           | 3.773               | MSMA                     | 0.000                                       | 0.000             | 3.773             |
| FOREST RANGE<br>MANAGEMENT         | 2.000               | picloram                 | 0.000                                       | 2.000             | 0.000             |
| FOREST SITE<br>REHABILITATION      | 108.000             | glyphosate               | 108.000                                     | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>          | <b>1 475.843</b>    |                          | <b>1 300.000</b>                            | <b>172.100</b>    | <b>3.773</b>      |
| FOREST INDUSTRY                    |                     |                          |                                             |                   |                   |
| FOREST BRUSHING                    | 6 499.890           | glyphosate<br>hexazinone | 6 419.090<br>0.000                          | 15.000<br>24.300  | 41.500<br>0.000   |
| FOREST CONIFER<br>RELEASE          | 4 268.510           | glyphosate               | 4 264.910                                   | 3.600             | 0.000             |
| FOREST INSECT<br>CONTROL           | 0.270               | MSMA                     | 0.000                                       | 0.000             | 0.270             |
| FOREST RIGHT-OF-WAY<br>MAINTENANCE | 130.970             | glyphosate               | 0.000                                       | 130.970           | 0.000             |
| FOREST SITE<br>PREPARATION         | 2 014.900           | glyphosate               | 2 014.900                                   | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>          | <b>12 914.540</b>   |                          | <b>12 698.900</b>                           | <b>173.870</b>    | <b>41.770</b>     |
| <b><u>TOTALS:</u></b>              | <b>14 390.413</b>   |                          | <b>13 998.900</b>                           | <b>345.970</b>    | <b>45.543</b>     |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: ROBSON VALLEY**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL   | 118.000             | MSMA       | 0.000                                       | 0.000             | 118.000           |
| FOREST RANGE<br>MANAGEMENT | 17.660              | glyphosate | 0.000                                       | 0.870             | 0.000             |
|                            |                     | picloram   | 0.000                                       | 16.790            | 0.000             |
| <hr/>                      |                     |            |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>  | <b>135.660</b>      |            | <b>0.000</b>                                | <b>17.660</b>     | <b>118.000</b>    |
| <hr/>                      |                     |            |                                             |                   |                   |
| FOREST INDUSTRY            |                     |            |                                             |                   |                   |
| FOREST BRUSHING            | 1 199.700           | glyphosate | 1 199.700                                   | 0.000             | 0.000             |
| <hr/>                      |                     |            |                                             |                   |                   |
| <b><u>SUB TOTALS:</u></b>  | <b>1 199.700</b>    |            | <b>1 199.700</b>                            | <b>0.000</b>      | <b>0.000</b>      |
| <hr/>                      |                     |            |                                             |                   |                   |
| <b><u>TOTALS:</u></b>      | <b>1 335.360</b>    |            | <b>1 199.700</b>                            | <b>17.660</b>     | <b>118.000</b>    |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: VANDERHOOF**

| PURPOSE                         | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                 |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                        |                     |            |                                             |                   |                   |
| FOREST BRUSHING                 | 465.200             | glyphosate | 465.200                                     | 0.000             | 0.000             |
| FOREST INSECT<br>CONTROL        | 0.200               | MSMA       | 0.000                                       | 0.000             | 0.200             |
| <u>SUB TOTALS:</u>              | 465.400             |            | 465.200                                     | 0.000             | 0.200             |
| FOREST INDUSTRY                 |                     |            |                                             |                   |                   |
| FOREST BRUSHING                 | 263.900             | glyphosate | 79.500                                      | 184.400           | 0.000             |
| FOREST PRE-HARVEST<br>TREATMENT | 472.600             | glyphosate | 0.000                                       | 0.000             | 472.600           |
| <u>SUB TOTALS:</u>              | 736.500             |            | 79.500                                      | 184.400           | 472.600           |
| <u>TOTALS:</u>                  | 1 201.900           |            | 544.700                                     | 184.400           | 472.800           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: FORT ST. JAMES**

| PURPOSE                         | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                 |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                        |                     |            |                                             |                   |                   |
| FOREST BRUSHING                 | 928.970             | glyphosate | 846.400                                     | 82.570            | 0.000             |
| FOREST INSECT<br>CONTROL        | 201.800             | MSMA       | 0.000                                       | 0.000             | 201.800           |
| <u>SUB TOTALS:</u>              | 1 130.770           |            | 846.400                                     | 82.570            | 201.800           |
| FOREST INDUSTRY                 |                     |            |                                             |                   |                   |
| FOREST BRUSHING                 | 199.600             | glyphosate | 124.600                                     | 75.000            | 0.000             |
| FOREST INSECT<br>CONTROL        | 1.500               | MSMA       | 0.000                                       | 0.000             | 1.500             |
| FOREST PRE-HARVEST<br>TREATMENT | 570.000             | glyphosate | 0.000                                       | 0.000             | 570.000           |
| FOREST SITE<br>PREPARATION      | 275.400             | glyphosate | 275.400                                     | 0.000             | 0.000             |
| <u>SUB TOTALS:</u>              | 1 046.500           |            | 400.000                                     | 75.000            | 571.500           |
| <u>TOTALS:</u>                  | 2 177.270           |            | 1 246.400                                   | 157.570           | 773.300           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: MACKENZIE**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES  | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|-------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |             | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                  |                     |             |                                             |                   |                   |
| FOREST INSECT<br>CONTROL  | 0.980               | mineral oil | 0.000                                       | 0.160             | 0.000             |
|                           |                     | MSMA        | 0.000                                       | 0.000             | 0.820             |
| <hr/>                     |                     |             |                                             |                   |                   |
| <u>SUB TOTALS:</u>        | 0.980               |             | 0.000                                       | 0.160             | 0.820             |
| <hr/>                     |                     |             |                                             |                   |                   |
| FOREST INDUSTRY           |                     |             |                                             |                   |                   |
| FOREST BRUSHING           | 592.200             | glyphosate  | 501.200                                     | 91.000            | 0.000             |
| FOREST CONIFER<br>RELEASE | 486.200             | glyphosate  | 486.200                                     | 0.000             | 0.000             |
| <hr/>                     |                     |             |                                             |                   |                   |
| <u>SUB TOTALS:</u>        | 1 078.400           |             | 987.400                                     | 91.000            | 0.000             |
| <hr/>                     |                     |             |                                             |                   |                   |
| <u>TOTALS:</u>            | 1 079.380           |             | 987.400                                     | 91.160            | 0.820             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: DAWSON CREEK**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES             | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |                        | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |                        |                                             |                   |                   |
| FOREST BRUSHING            | 2 697.400           | glyphosate             | 2 521.900                                   | 175.500           | 0.000             |
| FOREST RANGE<br>MANAGEMENT | 14.933              | glyphosate<br>picloram | 0.000<br>0.000                              | 0.002<br>14.761   | 0.000<br>0.170    |
| <u>TOTALS:</u>             | 2 712.333           |                        | 2 521.900                                   | 190.263           | 0.170             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: FORT ST. JOHN**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |            |                                             |                   |                   |
| FOREST BRUSHING            | 1 129.650           | glyphosate | 1 129.200                                   | 0.450             | 0.000             |
| FOREST CONIFER<br>RELEASE  | 862.100             | glyphosate | 634.400                                     | 0.000             | 227.700           |
| FOREST RANGE<br>MANAGEMENT | 3.500               | picloram   | 0.000                                       | 3.500             | 0.000             |
| FOREST SITE<br>PREPARATION | 27.000              | glyphosate | 27.000                                      | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>  | <b>2 022.250</b>    |            | <b>1 790.600</b>                            | <b>3.950</b>      | <b>227.700</b>    |
| FOREST INDUSTRY            |                     |            |                                             |                   |                   |
| FOREST BRUSHING            | 1 150.100           | glyphosate | 1 150.100                                   | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b>  | <b>1 150.100</b>    |            | <b>1 150.100</b>                            | <b>0.000</b>      | <b>0.000</b>      |
| <b><u>TOTALS:</u></b>      | <b>3 172.350</b>    |            | <b>2 940.700</b>                            | <b>3.950</b>      | <b>227.700</b>    |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: PRINCE GEORGE**

**DISTRICT: FORT NELSON**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                  |                     |            |                                             |                   |                   |
| FOREST BRUSHING           | 1 022.200           | glyphosate | 994.900                                     | 0.000             | 0.000             |
|                           |                     | hexazinone | 27.300                                      | 0.000             | 0.000             |
| FOREST CONIFER<br>RELEASE | 52.000              | glyphosate | 52.000                                      | 0.000             | 0.000             |
| <u>TOTALS:</u>            | 1 074.200           |            | 1 074.200                                   | 0.000             | 0.000             |

REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY

| PURPOSE                          | TOTAL (ha)<br>TREATED | PESTICIDES   | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|-----------------------|--------------|---------------------------------------------|-------------------|-------------------|
|                                  |                       |              | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| REGION: KAMLOOPS                 |                       |              |                                             |                   |                   |
| B.C. FOREST SERVICE              |                       |              |                                             |                   |                   |
| FOREST BRUSHING                  | 83.820                | glyphosate   | 11.400                                      | 72.420            | 0.000             |
| FOREST CONIFER<br>RELEASE        | 170.650               | glyphosate   | 63.900                                      | 0.000             | 106.750           |
| FOREST INSECT<br>CONTROL         | 89.640                | chlorpyrifos | 0.000                                       | 0.760             | 0.000             |
|                                  |                       | cyclohexnon  | 16.000                                      | 0.000             | 0.000             |
|                                  |                       | MSMA         | 0.000                                       | 36.200            | 0.680             |
|                                  |                       | verbenone    | 36.000                                      | 0.000             | 0.000             |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.770                 | metalaxyl    | 0.000                                       | 0.770             | 0.000             |
| FOREST RANGE<br>MANAGEMENT       | 6 333.680             | picloram     | 0.000                                       | 6 333.680         | 0.000             |
| FOREST WEED<br>SEED TREE CONTROL | 199.850               | glyphosate   | 0.000                                       | 0.000             | 199.850           |
| <u>SUB TOTALS:</u>               | 6 878.410             |              | 127.300                                     | 6 443.830         | 307.280           |

REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY

| PURPOSE                                     | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                             |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <b>REGION: KAMLOOPS<br/>FOREST INDUSTRY</b> |                     |            |                                             |                   |                   |
| FOREST BRUSHING                             | 362.180             | glyphosate | 30.000                                      | 211.600           | 120.530           |
|                                             |                     | picloram   | 0.000                                       | 0.000             | 0.50              |
| FOREST CONIFER<br>RELEASE                   | 357.300             | glyphosate | 0.000                                       | 357.300           | 0.000             |
| FOREST CONIFER<br>THINNING                  | 0.999               | glyphosate | 0.000                                       | 0.000             | 0.666             |
|                                             |                     | picloram   | 0.000                                       | 0.000             | 0.333             |
| FOREST INSECT<br>CONTROL                    | 3.950               | MSMA       | 0.000                                       | 0.000             | 3.950             |
| FOREST SITE<br>PREPARATION                  | 376.700             | glyphosate | 303.200                                     | 73.500            | 0.000             |
| <b><u>SUB TOTALS:</u></b>                   | <b>1 101.129</b>    |            | <b>333.200</b>                              | <b>642.400</b>    | <b>125.529</b>    |
| <b><u>TOTALS:</u></b>                       | <b>7 979.539</b>    |            | <b>460.500</b>                              | <b>7 086.230</b>  | <b>432.809</b>    |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: CLEARWATER**

| <b>PURPOSE</b>             | <b>TOTAL ha<br/>TREATED</b> | <b>PESTICIDES</b> | <b>AREA (ha) BREAKDOWN BY<br/>TREATMENT METHODS</b> |                                  |                                  |
|----------------------------|-----------------------------|-------------------|-----------------------------------------------------|----------------------------------|----------------------------------|
|                            |                             |                   | <b><u>AERIAL<br/>SPRAY</u></b>                      | <b><u>GROUND<br/>APPLIC.</u></b> | <b><u>STEM<br/>INJECTION</u></b> |
| <b>B.C.F.S.</b>            |                             |                   |                                                     |                                  |                                  |
| FOREST BRUSHING            | 0.720                       | glyphosate        | 0.000                                               | 0.720                            | 0.000                            |
| FOREST INSECT<br>CONTROL   | 0.500                       | MSMA              | 0.000                                               | 0.000                            | 0.500                            |
| FOREST RANGE<br>MANAGEMENT | 18.600                      | picloram          | 0.000                                               | 18.600                           | 0.000                            |
| <b><u>TOTALS:</u></b>      | <b>19.820</b>               |                   | <b>0.000</b>                                        | <b>19.320</b>                    | <b>0.500</b>                     |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: KAMLOOPS**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES   | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|--------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |              | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                         |                     |              |                                             |                   |                   |
| FOREST INSECT<br>CONTROL         | 36.940              | chlorpyrifos | 0.000                                       | 0.760             | 0.000             |
|                                  |                     | MSMA         | 0.000                                       | 0.000             | 0.180             |
|                                  |                     | verbenone    | 36.000                                      | 0.000             | 0.000             |
| FOREST RANGE<br>MANAGEMENT       | 456.500             | picloram     | 0.000                                       | 456.500           | 0.000             |
| FOREST WEED<br>SEED TREE CONTROL | 168.750             | glyphosate   | 0.000                                       | 0.000             | 168.750           |
| <u>TOTALS:</u>                   | 662.190             |              | 36.000                                      | 457.260           | 168.930           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: SALMON ARM**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                         |                     |            |                                             |                   |                   |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.660               | metalaxyl  | 0.000                                       | 0.660             | 0.000             |
| FOREST RANGE<br>MANAGEMENT       | 6.900               | picloram   | 0.000                                       | 6.900             | 0.000             |
| <b><u>SUB TOTALS:</u></b>        |                     |            | 0.000                                       | 7.560             | 0.000             |
| FOREST INDUSTRY                  |                     |            |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE        | 73.200              | glyphosate | 0.000                                       | 73.200            | 0.000             |
| FOREST SITE<br>PREPARATION       | 334.200             | glyphosate | 303.200                                     | 31.000            | 0.000             |
| <b><u>SUB TOTALS:</u></b>        |                     |            | 303.200                                     | 104.200           | 0.000             |
| <b><u>TOTALS:</u></b>            |                     |            | 303.200                                     | 111.760           | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

| <u>REGION: KAMLOOPS</u>    |                             | <u>DISTRICT: VERNON</u> |                                                     |                           |                           |
|----------------------------|-----------------------------|-------------------------|-----------------------------------------------------|---------------------------|---------------------------|
| <u>PURPOSE</u>             | <u>TOTAL ha<br/>TREATED</u> | <u>PESTICIDES</u>       | <u>AREA (ha) BREAKDOWN BY<br/>TREATMENT METHODS</u> |                           |                           |
|                            |                             |                         | <u>AERIAL<br/>SPRAY</u>                             | <u>GROUND<br/>APPLIC.</u> | <u>STEM<br/>INJECTION</u> |
| <b>B.C.F.S.</b>            |                             |                         |                                                     |                           |                           |
| FOREST BRUSHING            | 71.700                      | glyphosate              | 0.000                                               | 71.700                    | 0.000                     |
| FOREST CONIFER<br>RELEASE  | 0.500                       | glyphosate              | 0.000                                               | 0.000                     | 0.500                     |
| FOREST NURSERY<br>FUNGUS   | 0.110                       | metalaxyl               | 0.000                                               | 0.110                     | 0.000                     |
| FOREST RANGE<br>MANAGEMENT | 5 531.680                   | picloram                | 0.000                                               | 5 531.680                 | 0.000                     |
| <b><u>SUB TOTALS:</u></b>  | <b>5 603.990</b>            |                         | <b>0.000</b>                                        | <b>5 603.490</b>          | <b>0.500</b>              |
| <b>FOREST INDUSTRY</b>     |                             |                         |                                                     |                           |                           |
| FOREST BRUSHING            | 95.400                      | glyphosate              | 0.000                                               | 95.400                    | 0.000                     |
| FOREST CONIFER<br>RELEASE  | 284.100                     | glyphosate              | 0.000                                               | 284.100                   | 0.000                     |
| FOREST SITE<br>PREPARATION | 42.500                      | glyphosate              | 0.000                                               | 42.500                    | 0.000                     |
| <b><u>SUB TOTALS:</u></b>  | <b>422.000</b>              |                         | <b>0.000</b>                                        | <b>422.000</b>            | <b>0.000</b>              |
| <b><u>TOTALS:</u></b>      | <b>6 025.990</b>            |                         | <b>0.000</b>                                        | <b>6 025.490</b>          | <b>0.500</b>              |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: PENTICTON**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES             | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |                        | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |                        |                                             |                   |                   |
| FOREST CONIFER<br>RELEASE  | 106.250             | glyphosate             | 0.000                                       | 0.000             | 106.250           |
| <hr/>                      |                     |                        |                                             |                   |                   |
| <u>SUB TOTALS:</u>         | 106.250             |                        | 0.000                                       | 0.000             | 106.250           |
| <hr/>                      |                     |                        |                                             |                   |                   |
| FOREST INDUSTRY            |                     |                        |                                             |                   |                   |
| FOREST BRUSHING            | 0.080               | glyphosate             | 0.000                                       | 0.000             | 0.030             |
|                            |                     | picloram               | 0.000                                       | 0.000             | 0.050             |
| FOREST CONIFER<br>THINNING | 0.999               | glyphosate<br>picloram | 0.000<br>0.000                              | 0.000<br>0.000    | 0.666<br>0.333    |
| FOREST INSECT<br>CONTROL   | 3.950               | MSMA                   | 0.000                                       | 0.000             | 3.950             |
| <hr/>                      |                     |                        |                                             |                   |                   |
| <u>SUB TOTALS:</u>         | 5.029               |                        | 0.000                                       | 0.000             | 5.029             |
| <hr/>                      |                     |                        |                                             |                   |                   |
| <u>TOTALS:</u>             | 111.279             |                        | 0.000                                       | 0.000             | 111.279           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: MERRITT**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                         |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL         | 36.200              | MSMA       | 0.000                                       | 36.200            | 0.000             |
| FOREST RANGE<br>MANAGEMENT       | 320.000             | picloram   | 0.000                                       | 320.000           | 0.000             |
| FOREST WEED<br>SEED TREE CONTROL | 31.100              | glyphosate | 0.000                                       | 0.000             | 31.100            |
| <b><u>SUB TOTALS:</u></b>        | <b>387.300</b>      |            | <b>0.000</b>                                | <b>356.200</b>    | <b>31.100</b>     |
| FOREST INDUSTRY                  |                     |            |                                             |                   |                   |
| FOREST BRUSHING                  | 236.700             | glyphosate | 0.000                                       | 116.200           | 120.500           |
| <b><u>SUB TOTALS:</u></b>        | <b>236.700</b>      |            | <b>0.000</b>                                | <b>116.200</b>    | <b>120.500</b>    |
| <b><u>TOTALS:</u></b>            | <b>624.000</b>      |            | <b>0.000</b>                                | <b>472.400</b>    | <b>151.600</b>    |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: KAMLOOPS**

**DISTRICT: LILLOOET**

| PURPOSE                   | TOTAL ha<br>TREATED | PESTICIDES  | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------|---------------------|-------------|---------------------------------------------|-------------------|-------------------|
|                           |                     |             | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                  |                     |             |                                             |                   |                   |
| FOREST BRUSHING           | 11.400              | glyphosate  | 11.400                                      | 0.000             | 0.000             |
| FOREST CONIFER<br>RELEASE | 63.900              | glyphosate  | 63.900                                      | 0.000             | 0.000             |
| FOREST INSECT<br>CONTROL  | 16.000              | cyclohexnon | 16.000                                      | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b> | <b>91.300</b>       |             | <b>91.300</b>                               | <b>0.000</b>      | <b>0.000</b>      |
| FOREST INDUSTRY           |                     |             |                                             |                   |                   |
| FOREST BRUSHING           | 30.000              | glyphosate  | 30.000                                      | 0.000             | 0.000             |
| <b><u>SUB TOTALS:</u></b> | <b>30.000</b>       |             | <b>30.000</b>                               | <b>0.000</b>      | <b>0.000</b>      |
| <b><u>TOTALS:</u></b>     | <b>121.300</b>      |             | <b>121.300</b>                              | <b>0.000</b>      | <b>0.000</b>      |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| REGION: NELSON                   |                     |            |                                             |                   |                   |
| B.C. FOREST SERVICE              |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL         | 33.470              | MSMA       | 0.000                                       | 0.000             | 33.470            |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.600               | metalaxyl  | 0.000                                       | 0.600             | 0.000             |
| FOREST PRE-HARVEST<br>TREATMENT  | 67.300              | glyphosate | 0.000                                       | 0.000             | 67.300            |
| FOREST RANGE<br>MANAGEMENT       | 93.891              | glyphosate | 0.000                                       | 2.000             | 0.000             |
|                                  |                     | picloram   | 0.000                                       | 91.891            | 0.000             |
| <hr/>                            |                     |            |                                             |                   |                   |
| SUB TOTALS:                      | 195.261             |            | 0.000                                       | 94.491            | 100.770           |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                         | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                 |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <u>REGION: NELSON</u>           |                     |            |                                             |                   |                   |
| FOREST INDUSTRY                 |                     |            |                                             |                   |                   |
| FOREST PRE-HARVEST<br>TREATMENT | 99.750              | glyphosate | 0.000                                       | 0.000             | 99.750            |
| FOREST SITE<br>PREPARATION      | 424.880             | glyphosate | 415.630                                     | 9.250             | 0.000             |
| <u>SUB TOTALS:</u>              | 524.630             |            | 415.630                                     | 9.250             | 99.750            |
| <u>TOTALS:</u>                  | 719.891             |            | 415.630                                     | 103.741           | 200.520           |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: NELSON**

**DISTRICT: CRANBROOK**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |            |                                             |                   |                   |
| FOREST RANGE<br>MANAGEMENT | 21.400              | picloram   | 0.000                                       | 21.400            | 0.000             |
|                            |                     |            |                                             |                   |                   |
| <u>TOTALS:</u>             | 21.400              |            | 0.000                                       | 21.400            | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: NELSON**

**DISTRICT: INVERMERE**

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                   |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL   | 33.150              | MSMA       | 0.000                                       | 0.000             | 33.150            |
| FOREST RANGE<br>MANAGEMENT | 10.441              | picloram   | 0.000                                       | 10.441            | 0.000             |
|                            | ..                  |            |                                             |                   |                   |
| <u>TOTALS:</u>             | 43.591              |            | 0.000                                       | 10.441            | 33.150            |

SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY

REGION: NELSON

DISTRICT: GOLDEN

| PURPOSE                    | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                            |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| FOREST INDUSTRY            |                     |            |                                             |                   |                   |
| FOREST SITE<br>PREPARATION | 424.880             | glyphosate | 415.630                                     | 9.250             | 0.000             |
|                            |                     |            |                                             |                   |                   |
| <b>TOTALS:</b>             | <b>424.880</b>      |            | <b>415.630</b>                              | <b>9.250</b>      | <b>0.000</b>      |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: NELSON**

**DISTRICT: BOUNDARY**

| PURPOSE                         | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|---------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                 |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                        |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL        | 0.320               | MSMA       | 0.000                                       | 0.000             | 0.320             |
| FOREST PRE-HARVEST<br>TREATMENT | 67.300              | glyphosate | 0.000                                       | 0.000             | 67.300            |
| FOREST RANGE<br>MANAGEMENT      | 62.050              | glyphosate | 0.000                                       | 2.000             | 0.000             |
|                                 |                     | picloram   | 0.000                                       | 60.050            | 0.000             |
| <b><u>SUB TOTALS:</u></b>       | <b>129.670</b>      |            | <b>0.000</b>                                | <b>62.050</b>     | <b>67.620</b>     |
| FOREST INDUSTRY                 |                     |            |                                             |                   |                   |
| FOREST PRE-HARVEST<br>TREATMENT | 99.750              | glyphosate | 0.000                                       | 0.000             | 99.750            |
| <b><u>SUB TOTALS:</u></b>       | <b>99.750</b>       |            | <b>0.000</b>                                | <b>0.000</b>      | <b>99.750</b>     |
| <b><u>TOTALS:</u></b>           | <b>229.420</b>      |            | <b>0.000</b>                                | <b>62.050</b>     | <b>167.370</b>    |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <b>REGION: CARIBOO</b>           |                     |            |                                             |                   |                   |
| <b>B.C. FOREST SERVICE</b>       |                     |            |                                             |                   |                   |
| FOREST BRUSHING                  | 614.000             | glyphosate | 372.000                                     | 242.000           | 0.000             |
| FOREST CONIFER<br>RELEASE        | 652.500             | glyphosate | 524.300                                     | 128.200           | 0.000             |
| FOREST INSECT<br>CONTROL         | 0.510               | MSMA       | 0.000                                       | 0.000             | 0.510             |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.500               | metalaxyl  | 0.000                                       | 0.500             | 0.000             |
| <b><u>SUB TOTALS:</u></b>        | <b>1 267.510</b>    |            | <b>896.300</b>                              | <b>370.700</b>    | <b>0.510</b>      |

**REGIONAL SUMMARY OF PESTICIDE USE, 1990  
BCFS AND FOREST INDUSTRY**

| PURPOSE                                   | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|-------------------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                           |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| <u>REGION: CARIBOO</u><br>FOREST INDUSTRY |                     |            |                                             |                   |                   |
| FOREST BRUSHING                           | 1 723.265           | glyphosate | 1 643.465                                   | 64.000            | 0.000             |
|                                           |                     | hexazinone | 0.000                                       | 15.800            | 0.000             |
| FOREST CONIFER<br>RELEASE                 | 1 963.200           | glyphosate | 1 816.300                                   | 0.000             | 0.000             |
|                                           |                     | hexazinone | 0.000                                       | 146.900           | 0.000             |
| FOREST SITE<br>PREPARATION                | 138.300             | glyphosate | 0.000                                       | 44.500            | 0.000             |
|                                           |                     | hexazinone | 0.000                                       | 93.800            | 0.000             |
| <u>SUB TOTALS:</u>                        |                     |            | 3 459.765                                   | 365.000           | 0.000             |
| <u>TOTALS:</u>                            |                     |            | 4 356.065                                   | 735.700           | 0.510             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: CARIBOO**

**DISTRICT: QUESNEL**

| PURPOSE                          | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|----------------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                                  |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                         |                     |            |                                             |                   |                   |
| FOREST BRUSHING                  | 614.000             | glyphosate | 372.000                                     | 242.000           | 0.000             |
| FOREST CONIFER<br>RELEASE        | 652.500             | glyphosate | 524.300                                     | 128.200           | 0.000             |
| FOREST NURSERY<br>FUNGUS CONTROL | 0.500               | metalaxyl  | 0.000                                       | 0.500             | 0.000             |
| <u>SUB TOTALS:</u>               | 1 267.000           |            | 896.300                                     | 370.700           | 0.000             |
| FOREST INDUSTRY                  |                     |            |                                             |                   |                   |
| FOREST BRUSHING                  | 1 659.265           | glyphosate | 1 643.465                                   | 0.000             | 0.000             |
|                                  |                     | hexazinone | 0.000                                       | 15.800            | 0.000             |
| FOREST CONIFER<br>RELEASE        | 1 963.200           | glyphosate | 1 816.300                                   | 0.000             | 0.000             |
|                                  |                     | hexazinone | 0.000                                       | 146.900           | 0.000             |
| FOREST SITE<br>PREPARATION       | 138.300             | glyphosate | 0.000                                       | 44.500            | 0.000             |
|                                  |                     | hexazinone | 0.000                                       | 93.800            | 0.000             |
| <u>SUB TOTALS:</u>               | 3 760.765           |            | 3 459.765                                   | 301.000           | 0.000             |
| <u>TOTALS:</u>                   | 5 027.765           |            | 4 356.065                                   | 671.700           | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: CARIBOO**

**DISTRICT: 100 MILE HOUSE**

| PURPOSE         | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|-----------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                 |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| FOREST INDUSTRY |                     |            |                                             |                   |                   |
| FOREST BRUSHING | 64.000              | glyphosate | 0.000                                       | 64.000            | 0.000             |
|                 |                     |            |                                             |                   |                   |
| <u>TOTALS:</u>  | 64.000              |            | 0.000                                       | 64.000            | 0.000             |

**SUMMARY OF PESTICIDE USE BY FOREST REGION AND DISTRICT, 1990  
BCFS AND FOREST INDUSTRY**

**REGION: CARIBOO**

**DISTRICT: CHILCOTIN**

| PURPOSE                  | TOTAL ha<br>TREATED | PESTICIDES | AREA (ha) BREAKDOWN BY<br>TREATMENT METHODS |                   |                   |
|--------------------------|---------------------|------------|---------------------------------------------|-------------------|-------------------|
|                          |                     |            | AERIAL<br>SPRAY                             | GROUND<br>APPLIC. | STEM<br>INJECTION |
| B.C.F.S.                 |                     |            |                                             |                   |                   |
| FOREST INSECT<br>CONTROL | 0.510               | MSMA       | 0.000                                       | 0.000             | 0.510             |
| TOTALS:                  | 0.510               |            | 0.000                                       | 0.000             | 0.510             |


TABLE VII Cont.  
PROVINCIAL SUMMARY OF AREA (ha)  
TREATED BY CONTROL PURPOSE, 1990

| PURPOSE                                   | PESTICIDE                                  | B.C. FOREST<br>SERVICE<br>ha Treated | FOREST<br>INDUSTRY<br>ha Treated |
|-------------------------------------------|--------------------------------------------|--------------------------------------|----------------------------------|
| FOREST<br>PRE-HARVEST<br>TREATMENT        | glyphosate                                 | 67.300                               | 1 142.350                        |
| TOTAL ha.:                                | 1 209.650                                  |                                      |                                  |
| FOREST RANGE<br>MANAGEMENT                | glyphosate<br>picloram                     | 2.992<br>6 476.172                   | 0.000<br>0.000                   |
| TOTAL ha.:                                | 6 479.164                                  |                                      |                                  |
| FOREST<br>RIGHT-OF-WAY<br>MAINTENANCE     | 2, 4-D amine<br>2, 4-D ester<br>glyphosate | 0.000<br>0.000<br>0.000              | 29.000<br>24.000<br>483.320      |
| TOTAL ha.:                                | 536.320                                    |                                      |                                  |
| FOREST SEED<br>AND CONE<br>INSECT CONTROL | acephate<br>oxydemeton                     | 0.012<br>0.012                       | 0.000<br>0.000                   |
| TOTAL ha.:                                | 0.024                                      |                                      |                                  |
| FOREST SITE<br>PREPARATION                | glyphosate<br>hexazinone                   | 166.600<br>0.000                     | 4 128.910<br>93.800              |
| TOTAL ha.:                                | 4 389.310                                  |                                      |                                  |
| FOREST SITE<br>REHABILITATION             | glyphosate                                 | 425.500                              | 0.000                            |
| TOTAL ha.:                                | 425.500                                    |                                      |                                  |
| FOREST<br>VERTEBRATE<br>CONTROL           | diphacinone<br>putrid eggs                 | 0.000<br>0.000                       | 0.044<br>0.800                   |
| TOTAL ha.:                                | 0.844                                      |                                      |                                  |
| FOREST WEED<br>SEED TREE CONTROL          | glyphosate                                 | 199.850                              | 0.000                            |
| TOTAL ha.:                                | 199.850                                    |                                      |                                  |
| <u>GRAND TOTALS:</u>                      |                                            | 21 058.436                           | 36 603.086                       |

TABLE VII Cont.  
PROVINCIAL SUMMARY OF AREA (ha)  
TREATED BY CONTROL PURPOSE, 1990

| PURPOSE                                   | PESTICIDE                                  | B.C. FOREST<br>SERVICE<br>ha Treated | FOREST<br>INDUSTRY<br>ha Treated |
|-------------------------------------------|--------------------------------------------|--------------------------------------|----------------------------------|
| FOREST<br>PRE-HARVEST<br>TREATMENT        | glyphosate                                 | 67.300                               | 1 142.350                        |
| TOTAL ha.:                                | 1 209.650                                  |                                      |                                  |
| FOREST RANGE<br>MANAGEMENT                | glyphosate<br>picloram                     | 2.992<br>6 476.172                   | 0.000<br>0.000                   |
| TOTAL ha.:                                | 6 479.164                                  |                                      |                                  |
| FOREST<br>RIGHT-OF-WAY<br>MAINTENANCE     | 2, 4-D amine<br>2, 4-D ester<br>glyphosate | 0.000<br>0.000<br>0.000              | 29.000<br>24.000<br>483.320      |
| TOTAL ha.:                                | 536.320                                    |                                      |                                  |
| FOREST SEED<br>AND CONE<br>INSECT CONTROL | acephate<br>oxydemeton                     | 0.012<br>0.012                       | 0.000<br>0.000                   |
| TOTAL ha.:                                | 0.024                                      |                                      |                                  |
| FOREST SITE<br>PREPARATION                | glyphosate<br>hexazinone                   | 166.600<br>0.000                     | 4 128.910<br>93.800              |
| TOTAL ha.:                                | 4 389.310                                  |                                      |                                  |
| FOREST SITE<br>REHABILITATION             | glyphosate                                 | 425.500                              | 0.000                            |
| TOTAL ha.:                                | 425.500                                    |                                      |                                  |
| FOREST<br>VERTEBRATE<br>CONTROL           | diphacinone<br>putrid eggs                 | 0.000<br>0.000                       | 0.044<br>0.800                   |
| TOTAL ha.:                                | 0.844                                      |                                      |                                  |
| FOREST WEED<br>SEED TREE CONTROL          | glyphosate                                 | 199.850                              | 0.000                            |
| TOTAL ha.:                                | 199.850                                    |                                      |                                  |
| <u>GRAND TOTALS:</u>                      |                                            | 21 058.436                           | 36 603.086                       |

# PROVINCIAL SUMMARY OF AREA (ha) TREATED WITH PESTICIDES BY CONTROL PURPOSE, 1990<sup>1</sup>



## CONTROL PURPOSE

<sup>1</sup> B.C. Forest Service & Industry Combined

**TABLE VIII**  
**PROVINCIAL SUMMARY OF AREA (ha)**  
**TREATED BY CONTROL METHOD, 1990**

| METHOD                                                                   | PESTICIDE        | B.C. FOREST<br>SERVICE | FOREST<br>INDUSTRY |
|--------------------------------------------------------------------------|------------------|------------------------|--------------------|
|                                                                          |                  | ha TREATED             | ha TREATED         |
| VEHICLE MOUNTED<br>SPRAY BOOM                                            | glyphosate       | 98.070                 | 379.270            |
|                                                                          | hexazinone       | 0.000                  | 93.800             |
|                                                                          | metalaxyl        | 2.260                  | 0.000              |
|                                                                          | picloram         | 6 094.020              | 0.000              |
|                                                                          | treademefon      | 0.350                  | 0.000              |
| <b>TOTAL ha.:</b>                                                        | <b>6 667.770</b> |                        |                    |
| BACK PACK<br>SPRAYER AND<br>HAND HELD<br>SPRAY<br>EQUIPMENT <sup>1</sup> | 2, 4-D amine     |                        | 29.000             |
|                                                                          | carbaryl         | 0.840                  | 0.000              |
|                                                                          | chlorpyrifos     | 0.760                  | 0.000              |
|                                                                          | glyphosate       | 1 181.8801             | 1 688.800          |
|                                                                          | hexazinone       | 0.000                  | 154.400            |
|                                                                          | metalaxyl        | 0.110                  | 0.000              |
|                                                                          | mineral oil      | 0.160                  | 0.000              |
|                                                                          | MSMA             | 36.200                 | 1.300              |
|                                                                          | picloram         | 2.000                  | 0.000              |
|                                                                          | triclopyr        | 0.100                  | 0.000              |
| <b>TOTAL ha.:</b>                                                        | <b>3 095.550</b> |                        |                    |
| VEHICLE MOUNTED<br>SPRAY NOZZLES                                         | 2, 4-D ester     | 0.000                  | 24.000             |
|                                                                          | acephate         | 0.012                  | 0.000              |
|                                                                          | glyphosate       | 221.322                | 606.880            |
|                                                                          | hexazinone       | 0.000                  | 32.600             |
|                                                                          | oxydemeton       | 0.012                  | 0.000              |
|                                                                          | picloram         | 379.982                | 0.000              |
|                                                                          | pirmicarb        | 0.005                  | 0.000              |
| <b>TOTAL ha.:</b>                                                        | <b>1 264.813</b> |                        |                    |
| HAND BROADCAST                                                           | diphacinone      | 0.000                  | 0.044              |
| <b>TOTAL ha.:</b>                                                        | <b>0.044</b>     |                        |                    |


TABLE VIII Cont.  
PROVINCIAL SUMMARY OF AREA (ha)  
TREATED BY CONTROL METHOD, 1990

| METHOD                                                   | PESTICIDE     | B.C. FOREST<br>SERVICE<br>ha TREATED | FOREST<br>INDUSTRY<br>ha TREATED |
|----------------------------------------------------------|---------------|--------------------------------------|----------------------------------|
| TREE INJECTION/<br>CUT SURFACE<br>TREATMENT <sup>2</sup> | 2, 4-D amine  | 0.000                                | 60.000                           |
|                                                          | acephate      | 0.000                                | 0.050                            |
|                                                          | glyphosate    | 993.250                              | 3 384.184                        |
|                                                          | MSMA          | 387.713                              | 5.720                            |
|                                                          | picloram      | 0.000                                | 0.393                            |
| TOTAL ha.:                                               | 4 831.310     |                                      |                                  |
| STUMP<br>APPLICATION                                     | 2, 4-D amine  | 90.800                               | 848.000                          |
|                                                          | glyphosate    | 367.550                              | 1 484.900                        |
|                                                          | picloram      | 0.170                                | 0.000                            |
| TOTAL ha.:                                               | 2 791.420     |                                      |                                  |
| AERIAL<br>FIXED WING                                     | <u>B.t.k.</u> | 0.000                                | 360.000                          |
|                                                          | cyclohexnon   | 16.000                               | 0.000                            |
|                                                          | glyphosate    | 4 312.700                            | 17 033.930                       |
|                                                          | hexazinone    | 27.300                               | 0.000                            |
| TOTAL ha.:                                               | 21 749.930    |                                      |                                  |
| AERIAL<br>HELICOPTER                                     | glyphosate    | 6 808.870                            | 10 415.015                       |
|                                                          | verbenone     | 36.000                               | 0.000                            |
| TOTAL ha.:                                               | 17 259.885    |                                      |                                  |
| OTHER                                                    | putrid eggs   | 0.000                                | 0.800                            |
| TOTAL ha.:                                               | 0.800         |                                      |                                  |
| <u>GRAND TOTALS:</u>                                     |               | 21 058.436                           | 36 603.086                       |

<sup>1</sup>Includes mistblowers, ultra low volume sprayers, spot guns and granule spreaders.

<sup>2</sup>Involves hack and squirt, frill and squirt, and cut stump treatments. Includes brush saws, lance injectors and hypo hatchets.

# PROVINCIAL SUMMARY OF AREA (ha) TREATED WITH PESTICIDES BY METHOD OF APPLICATION 1990<sup>1</sup>



## METHOD OF APPLICATION

<sup>1</sup> B.C. Forest Service & Industry Combined


**TABLE IX**  
**TOTAL AREA (ha) TREATED WITH HERBICIDES AND INSECTICIDES**  
**FOR FORESTRY PURPOSES, 1980 - 1990**

| YEAR | B.C. FOREST SERVICE<br>HERB. | INSECT.              | FOREST INDUSTRY<br>HERB. | INSECT.          | TOTALS<br>HERB. | INSECT.  |
|------|------------------------------|----------------------|--------------------------|------------------|-----------------|----------|
| 1980 | 1 614.0                      | 1.0                  | 3 412.0                  | 70.0             | 5 026.0         | 71.0     |
| 1981 | 2 361.0                      | 134.0                | 4 232.0                  | 0.4              | 6 593.0         | 134.0    |
| 1982 | 2 630.0                      | 786.0                | 1 636.0                  | 0.1 <sup>1</sup> | 4 266.0         | 786.0    |
| 1983 | 2 586.0                      | 807.0                | 3 595.0                  | 5.0              | 6 181.0         | 812.0    |
| 1984 | 6 184.0                      | 199.0                | 7 709.0                  | 33.0             | 13 893.0        | 232.0    |
| 1985 | 8 018.0                      | 612.0                | 19 133.0                 | 26.0             | 27 151.0        | 638.0    |
| 1986 | 6 434.0                      | 162.0                | 18 487.0                 | 200.0            | 24 921.0        | 362.0    |
| 1987 | 15 705.0                     | 1 035.0 <sup>2</sup> | 26 791.0                 | 282.0            | 42 496.0        | 1 317.0  |
| 1988 | 17 378.0                     | 1 876.0              | 24 224.0                 | 270.0            | 41 602.0        | 2 146.0  |
| 1989 | 17 201.9                     | 566.03               | 40 901.13                | 734.4            | 58 103.03       | 1 300.43 |
| 1990 | 20 575.01                    | 478.0                | 36 235.2                 | 367.1            | 56 810.2        | 845.1    |


<sup>1</sup>Area treated was less than 0.1 ha.

<sup>2</sup>Insecticide figures for 1987 have been revised.

# AREA TREATED WITH HERBICIDES FOR FORESTRY PURPOSES 1980-1990



# AREA TREATED WITH INSECTICIDES FOR FORESTRY PURPOSES 1980-1990<sup>1</sup>



<sup>1</sup> B.C. Forest Service & Industry Combined

### DISCUSSIONS AND CONCLUSIONS

Pesticides were applied to 57 662 ha of B.C. forest and rangeland in 1990, approximately the same level of use as 1989 (59 425 ha). Forest companies were responsible for 64% of the total area treated and the Forest Service for 36%, as compared to 70% and 30%, respectively, in 1989.

Glyphosate was applied to over 80% of the provincial area treated with pesticides in 1990 and 1989. The use of 2,4-D has declined from 3 346 ha treated in 1989 to 1 052 ha in 1990.

Herbicide applications for the purpose of controlling unwanted vegetation remained at the same levels as 1989.

Helicopters and fixed-wing aircraft were used to apply pesticides to approximately 68% of the total provincial area treated in both 1989 and 1990.

**ACKNOWLEDGEMENTS**

The author would like to thank the Ministry of Environment, Pesticide Control Branch for its ongoing cooperation in providing this data.