

PROBE: PROTOCOL FOR OPERATIONAL BRUSHING EVALUATIONS

(FIRST APPROXIMATION)

Land Management
Report NUMBER **86**

ISSN 0702-9861

1993



Province of British Columbia
Ministry of Forests

PROBE

PROTOCOL FOR OPERATIONAL BRUSHING EEVALUATIONS

(FIRST APPROXIMATIONS)

S. W. Simard



Ministry of Forests
Research Program

1 9 9 3

Canadian Cataloguing in Publication Data

Simard, S. W.

PROBE : protocol for operational brushing evaluations
(first approximation)

(Land management report, ISSN 0702-9861 , no. 86)

Includes bibliographical references: p.
ISBN 0-7726-1891-7

1. Brush - Control - British Columbia. 2. Trees -
Weed control - British Columbia. I. British Columbia.
Ministry of Forests. II. Title. III. Series.

SD391.S55 1993 634.9'55'09711 C93-092323-5

Prepared by S.W. Simard
Forest Sciences Section
Kamloops Forest Region
515 Columbia Street
Kamloops, B.C. V2C 2T7
For the Research Program,
Ministry of Forests

© 1993 Province of British Columbia
Published by the
Research Branch
Ministry of Forests
31 Bastion Square
Victoria, B.C. V8W 3E7

Copies of this and other Ministry of Forests titles are
available from Crown Publications Inc., 546 Yates
Street, Victoria, B.C. V8W 1K8.

ACKNOWLEDGEMENTS

This protocol was developed for monitoring operational brushing treatments in the Kamloops and Nelson Forest Regions. The need for the protocol was endorsed by the Forest Districts in the Kamloops Forest Region at a meeting held on February 27, 1991, attended by R. Gray, R. Enfield, B. Nash, D. Horne, D. Purdy, J. Wright and S. Simard.

PROBE builds on the concepts presented in other vegetation management research design protocols, particularly those by Herring and Pollack (1985), Hays (1990, unpubl.) and Northern Interior Vegetation Management Association (1990, unpubl.). The author wishes to make special acknowledgement of their contribution. PROBE includes options for monitoring habitat conditions using the procedures of the Habitat Monitoring Committee (1990), as well as options for assessing impacts of broadcast applications of prescribed fire and mechanical site preparation, using modifications of procedures described by Trowbridge *et al.* (1986) and Curran and Thompson (1991, unpubl.), respectively.

The PROBE Data Entry System and SAS programs were developed by D. Sachs.

Thanks go to P. Comeau, W. Bergerud, W. Hayes, C. Thompson, A. Vyse, J. Pollack, G. Hope, D. Coates, T. Newsome, B. D'Anjou, D. Clark and G. Reichenbach for valuable suggestions and advice in manuscript revision.

The field procedures were tested and improved by J. Mather and F. Newhouse. The author wishes to make special acknowledgement of their work. The protocol will be subject to further revisions and improvements as more experience is gained and new situations are encountered.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
1 INTRODUCTION	1
2 ESTABLISHMENT PROCEDURE	2
2.1 Introduction	2
2.2 Site Selection	2
2.3 Treatment and Control Plot Location	3
2.4 Subplot Location	3
3 ASSESSMENTS	6
3.1 Introduction	6
3.2 Target Crop Trees	6
3.2.1 Growth	6
3.2.2 Density and stocking	7
3.3 Target Non-crop Vegetation	7
3.3.1 Competitive effect	7
3.3.2 Efficacy of treatment	7
3.4 Wildlife Habitat (Optional)	8
3.5 Prescribed Fire (Optional)	8
3.5.1 Sample design	9
3.5.2 Slash load	9
3.5.3 Forest floor and mineral soil exposure	9
3.5.4 Planting	9
3.6 Mechanical Site Preparation (Optional)	9
3.6.1 Planting	10
3.6.2 Sampling design and disturbance survey	10
4 RECORDS	10
5 PHOTOGRAPHS	10
6 MAPPING	10
7 TIMING	11
8 MAINTENANCE	11
9 PERSON-DAY REQUIREMENTS	12
10 RESPONSIBILITIES	12
11 ANALYSES	13
11.1 Introduction	13
11.2 T-tests	13
11.3 Analysis of Variance	13
11.4 Regression Analysis	14
11.5 Contingency Tables	14

12	DATA MANAGEMENT	15
13	PERSONNEL QUALIFICATIONS	15
14	LIST OF MATERIALS	15

APPENDICES

1	Species composition of major vegetation complexes in the southern Interior of British Columbia (from Kimmins and Comeau 1990)	17
2	PROBE forms and coding instructions	19
3	Codes for competitive status, vigour, condition, damage, phenology, and tree symbols	29
4	Silviculture Survey forms and coding instructions (from B.C. Ministry of Forests 1989)	34
5	Habitat Monitoring forms (from Habitat Monitoring Committee 1990)	37
6	Fuel Assessment form and tally rules (from Trowbridge <i>et al.</i> 1986)	45
7	MSP Soil Disturbance Survey forms and Soil Disturbance Categories (from Curran and Thompson 1991, unpubl.)	47
8	Analysis of Variance table	52
	REFERENCES	53

TABLES

1	Sample matrix for PROBE installations in the Kamloops and Nelson Forest Regions	4
2	Criteria for selecting comparable Treatment (T) and Control (C) plots	5
3	List of forms for operational brushing evaluations	11
4	Estimate of crew-day requirements (2 persons) per installation	12
5	Competition indices	14

FIGURES

1	Treatment or Control plot design	5
---	--	---

1 INTRODUCTION

An expanding brushing program in the southern Interior of British Columbia has created a need for better information on the short- and long-term effectiveness of vegetation management treatments that have been applied on an operational basis. Qualitative evaluations of treatments were useful to operational foresters in the formative stages of the program, but today have little value in defending prescriptions before the public. A quantitative, technically valid, and yet simple monitoring system is now necessary to meet the needs of operational staff.

In the past, several *ad hoc*, quantitative methods for evaluating operational vegetation management treatments were developed and implemented in the southern Interior. Each method, however, had specific objectives for evaluating specific vegetation management treatments. The narrow scope and lack of consistency among evaluation methods gave operational foresters limited opportunities for comparing results among sites and treatments. Development of a standardized monitoring system was subsequently requested.

This Land Management Handbook presents a design protocol that defines minimum standards for an objective, low intensity monitoring system. It is called Protocol for Operational Bushing Evaluations, or PROBE. The purpose of PROBE is to provide a monitoring framework for use in a wide array of vegetation complexes, ecosystems (site series), and operational vegetation management treatments. The framework standardizes installation location, response measurements and statistical analyses.

PROBE is intended solely for voluntary operational monitoring. It provides for extensive, quantitative evaluations of common treatments in the most frequently treated vegetation complexes; and is detailed enough that an individual who is familiar with research plot layout and monitoring can easily complete a high quality installation. A large number of PROBE installations can be readily established by a variety of individuals. PROBE is *not* designed for qualitative evaluation of the extent and success of all vegetation management activities in operational programs, nor is it designed to replace replicated research trials which may have different or more detailed objectives. PROBE should be considered an adjunct to both program-level evaluations *and* research trials, and used to fill rapidly the large information gap on the effectiveness of operational treatments.

The PROBE method satisfies the basic objective of most operational brushing evaluations, which is to determine the effects of vegetation management treatments on:

- the survival and growth of crop trees
- the abundance of target non-crop species

This it does by comparing crop tree and non-crop vegetation response in a designated treated area (called after this "Treatment" and abbreviated "T"), with response in a designated control area ("Control" and abbreviated "C"). When the user selects Treatment and Control plots, criteria are provided to ensure that the plots are located in areas that are similar in ecosystem characteristics and site history. Thirty-six tree-centred subplots are located in each of the Treatment and Control plots in which assessments are made. The subplots are assessed immediately before treatment and at pre-determined intervals up to 10 years after treatment.

The condition of wildlife habitat can be monitored within the PROBE framework, in addition to the basic assessments of target crop tree growth and target non-crop vegetation abundance. To that end, PROBE includes an option for assessing the abundance and distribution of all vascular plant species, using procedures developed by the Habitat Monitoring Committee (1990).

In addition to brushing treatments, site preparation treatments can also be monitored using PROBE. PROBE is best suited to broadcast treatments, such as broadcast burning or mechanical site preparation (MSP) using a brush blade. PROBE is less well suited to selective site preparation treatments, such as disc trenching or excavating, where treatments are applied in rows or to small, discrete microsites. Where PROBE is run to evaluate the effects of prescribed fire and MSP on the vegetation community, pre- and post-treatment conditions are evaluated using modifications of procedures described by Trowbridge *et al.* (1986) and Curran and Thompson,¹ respectively. The results are used to quantify treatment intensity, in much the same way as application rate quantifies the intensity of a herbicide treatment.

¹ Curran, M. and S. Thompson. 1991. Measuring soil disturbance following mechanical site preparation. B.C. Min. For., Nelson, B.C. Unpubl. report.

The Treatment and Control for a particular complex/ecosystem/treatment combination are replicated at least three times, once on each of three sites (i.e., three installations). Each installation serves as a block in a randomized block design (RBD). Comparisons of crop tree performance and non-crop vegetation abundance are then made between the Treatment and Control using Analysis of Variance (ANOVA). This approach requires that all blocks within the RBD be identical in treatment, plot layout and measurements. It is a powerful approach, in that conclusions may be drawn about treatments across a broad range of sites. Anyone establishing PROBE installations must be aware, however, that even a small, apparently slight deviation from the protocol may make a particular installation unusable to the whole (W. Bergerud, pers. comm., June 1991, B.C. Min. For., Victoria, B.C.). The procedures outlined in this protocol help to ensure that methods and measurements are standard and comparable. Complementary annual training sessions are also necessary to ensure that consistency is maintained among "probers."

In addition to comparing treatment means using ANOVA, the neighbourhood approach is used to build predictive models, which relate the performance of crop trees to the degree of crowding by neighbouring non-crop vegetation (Goldberg and Werner 1983; Weiner 1984; Wagner and Radosevich 1991). More specifically, simple regression models are constructed to relate the size, and possibly survival, of target crop trees to indices of non-crop vegetation competition. The competition indices are calculated from easily measured non-crop vegetation parameters, including height, cover and, for hardwoods only, density.

Two other monitoring systems are currently in use in the northern Interior of British Columbia: (1) Operational Herbicide Monitoring (OHM);² and (2) Unified System of Vegetation Monitoring (Northern Interior Vegetation Management Association (NIVMA)).³ OHM is designed for the establishment of benchmark installations, and includes detailed measurements of crop trees, competing vegetation, wildlife forage and environmental conditions. NIVMA is less comprehensive and objective. Its aim is to provide operational foresters with a simple, subjective monitoring system and a broad, shared database (NIVMAN). NIVMA is intended to complement OHM, so that forest managers are provided with both an extensive database (i.e., NIVMA) and objective treatment comparisons (i.e., OHM) (L. Herring, pers. comm., March 1991, B.C. Min. For., Prince George, B.C.).

The monitoring system presented in this report fulfils similar objectives to those of NIVMA and OHM. However, PROBE is more objective than NIVMA and less comprehensive than OHM, and hence represents a compromise between the two. NIVMA's system is unsuitable for the southern Interior for several reasons: (1) it does not permit objective evaluations of treatment effectiveness because a control is not required; (2) it does not meet the objectives of the Habitat Monitoring Committee (1990) for monitoring wildlife habitat; and (3) methods of data analysis are unclear. OHM also is considered unsuitable because of the large sample size and large number of variables it investigates. The monitoring system for the southern Interior, as with NIVMA's system in the north, however, would benefit from additional information gathered in OHM benchmark installations.

2 ESTABLISHMENT PROCEDURE

2.1 Introduction

This section describes the installation establishment procedure, which consists of three steps:

1. site selection
2. location of Treatment and Control plots based on similarity criteria
3. location of subplots within Treatment and Control plots

2.2 Site Selection

Site selection for operational brushing evaluations should be guided by the vegetation complexes, ecosystems and treatment regimes that are a high priority in the Kamloops and Nelson Forest Regions. The high priority vegetation complexes represent the early seral vegetation communities which are considered to compete with

² Hayes, W.J. 1990. Operational monitoring in the Prince George Forest Region. B.C. Min For., Prince George, B.C. Unpubl. report.

³ Northern Interior Vegetation Management Association (NIVMA). 1990. Unified system of silviculture monitoring. Prince George, B.C. Unpubl. report.

conifers in the southern Interior, as identified by Kimmins and Comeau (1990). The species characterizing those vegetation complexes are listed in Appendix 1. The high priority ecosystems are the site series of medium soil moisture regime (i.e., submesic, mesic, subhygric) and medium soil nutrient regime within the Biogeoclimatic Ecosystem Classification zones (Lloyd *et al.* 1990; Braumandl and Curran 1992) in which the vegetation complexes occur (Kimmins and Comeau 1990). The high priority treatments to be targeted by PROBE include all brushing and most site preparation treatments operational in the southern Interior. Sites selected should have been disturbed (i.e., harvested or site prepared) within the past 15 years.

Common complexes, ecosystems and treatment regimes in the Kamloops and Nelson Forest Regions are summarized in the sampling matrix shown in Table 1. Each cell within the sampling matrix represents a complex/ecosystem/treatment combination. Planted tree species are not included for the sake of simplicity. An example of one such combination is Dry Alder Complex/mesic site in the Montane Spruce zone, to which ground foliar glyphosate is applied at a rate of 2.14 kg a.i./ha to release interior spruce. The program objective is to replicate each combination (cell) three times.

Some combinations shown in Table 1 are rarely used, but are shown for completeness. These combinations will be given low priority through inter-region co-ordination. The regions will direct site selection to ensure that high priority combinations are sampled and that there is no unnecessary replication.

2.3 Treatment and Control Plot Location

Each installation should consist of a maximum of one Treatment and one Control plot. These are located in areas representative of the average treatment, and are homogeneous in ecosystem and site history. The Treatment and Control plots are located on the same opening where possible, to minimize between-plot variation. They must be similar to one another in the criteria described in Table 2. To ensure Treatment and Control plots are comparable, all ecosystem characteristics must be evaluated in concert (W. Hayes, pers. comm., June 1991, Prince George, B.C.). Buffer zones or areas that differ from the average conditions of the opening (e.g., gullies, creeks, rock outcrops, highly disturbed) must be avoided. The Treatment and Control plots must be located at least 30 m from treatment edges (e.g., timber edge, landings, roads). Where the treatment is an aerial herbicide application, however, (at least a minimum of) 50 m is required.

The Treatment and Control plots can be laid out in either a square or rectangular configuration. The dimensions of a square plot are 90 × 90 m (0.81 ha). Where a rectangular configuration is more suitable, any other dimensions may be used, provided the area is at least 0.80 ha (e.g., 120 × 70 m = 0.84 ha). The same plot configuration should be used for both the Treatment and Control within an installation, where possible. These plot dimensions include a 20-m surrounding buffer (Figure 1). The area in the plot which includes the buffer is referred to as the "treatment plot," and that which does not include the buffer is referred to as the "measurement plot."

The four corners of the Treatment and Control plots should be permanently marked with pressure-treated fence posts. The top of the posts are painted red in the Control and blue in the Treatment. Each post is marked with an aluminum tag detailing the installation number, treatment, corner, and distance and bearing to the two adjacent corners. Before treatment, the Control perimeter is clearly marked at 3-m intervals with winter-weight pink and blue flagging, and the Treatment perimeter with yellow or lime-green flagging. In the case of aerial herbicide, the corners are marked with large, orange, plastic garbage bags, or according to instructions provided by the pilot.

A point of commencement (POC) is located at an easily recognizable point and marked with an aluminum tag on a pressure-treated fence post. The tag details the installation number, date of establishment, Ministry contact, and distance and bearing to the first subplot in each of the Treatment and Control.

2.4 Subplot Location

Within each Treatment and Control plot, 36 tree-centred, circular subplots are located in a square 6 × 6 grid (or a rectangular grid such as 9 × 4, depending on the plot configuration). This is done by systematically locating six gridlines, each comprised of six gridpoints, inside the measurement plot. The first gridpoint, which marks the beginning of the first gridline, is randomly located inside one of the measurement plot corners.

TABLE 1. Sample matrix for Probe installations in the Kamloops and Nelson Forest Regions

Complex	Zone	TREATMENT ^a												
		M	GGA	CSG	CSS	HS	BBT	G	SG	CG	BB	BM	CS	BrB
Pinegrass	IDF	— ^b	X ^c	—	—	—	—	—	X	X	X	X	X	X
	MS	—	X	—	—	—	—	—	X	X	X	X	X	X
	ESSF	—	X	—	—	—	—	—	X	X	X	X	X	X
Fireweed	ICH	X	X	—	—	—	—	—	X	X	X	X	—	—
	MS	X	X	—	—	—	—	—	X	X	X	X	—	—
	ESSF	X	X	—	—	—	—	—	X	X	X	X	—	—
Fern	ICH	X	X	—	—	—	—	—	X	X	X	X	—	—
Dry Alder	MS	X	X	X	X	X	X	—	X	X	X	X	—	X
	ESSF	X	X	X	X	X	X	—	X	X	X	X	—	X
Wet Alder	ICH	X	X	X	X	X	X	—	X	X	X	X	—	X
	ESSF	X	X	X	X	X	X	—	X	X	X	X	—	X
Mixed Shrub	ICH	X	X	—	—	—	—	—	X	X	X	X	—	X
	ESSF	X	X	—	—	—	—	—	X	X	X	X	—	X
Dry Shrub	IDF	X	X	—	—	—	—	—	X	X	X	X	—	X
	ICH	X	X	—	—	—	—	—	X	X	X	X	—	X
Ericaceous Shrub	ESSF	X	X	—	—	—	—	—	—	—	X	X	—	X
Subalpine Herb	ESSF	X	X	—	—	—	—	—	X	X	X	X	—	—
Aspen	IDF	X	X	X	X	X	X	X	X	X	X	X	—	X
	ICH	X	X	X	X	X	X	X	X	X	X	X	—	X
	MS	X	X	X	X	X	X	X	X	X	X	X	—	X
Cottonwood	IDF	X	X	X	X	X	X	X	X	X	X	X	—	X
	ICH	X	X	X	X	X	X	X	X	X	X	X	—	X
Mixed Hardwood	IDF	X	X	X	X	X	X	X	X	X	X	X	—	X
	ICH	X	X	X	X	X	X	X	X	X	X	X	—	X
	MS	X	X	X	X	X	X	X	X	X	X	X	—	X
Willow	IDF	X	X	X	X	X	—	—	X	X	X	X	—	X
	ICH	X	X	X	X	X	—	—	X	X	X	X	—	X
	MS	X	X	X	X	X	—	—	X	X	X	X	—	X
	ESSF	X	X	X	X	X	—	—	X	X	X	X	—	X
Domestic Grasses	IDF	X	X	—	—	—	—	—	X	X	X	X	X	—
	MS	X	X	—	—	—	—	—	X	X	X	X	X	—

^a M = manual cut (or bend fireweed)

GGA = spray glyphosate, ground or aerial application

CSG = manual cut and swab stump with glyphosate

CSS = manual cut and later spray sprouts with glyphosate

HS = hack-and-squirt or e-z-ject with glyphosate

BBT = basal bark treatment using triclopyr ester

G = girdle

SG = sheep grazing

CG = cattle grazing

BB = broadcast burn

BM = mechanical bunch-and-burn

CS = chain scarifier

BrB = brush blade

^b — = not applicable^c X = sample of 3 required




FIGURE 1. Treatment or Control plot design.

TABLE 2. Criteria for selecting comparable Treatment (T) and Control (C) plots

Characteristic	Similarity criteria
<i>Ecosystem</i>	
Mesoslope position	same class (crest, upper slope, middle slope, lower slope, toe, depression, level)
Slope	$\pm 10\%$ (e.g., if slope in T is 50%, then slope in C can range from 40 to 60%)
Aspect	$\pm 45^\circ$ (e.g., if aspect in T is 180° , then aspect in C can range from 135° to 245°)
Elevation	± 50 m
Soil texture	same broad class: sand, silt, clay, loam (e.g., loamy sand and silty sand are the same broad class)
Coarse fragment content	$\pm 25\%$
Soil moisture regime	same class (submesic [SM], mesic [M] or subhygric [SHG])
Site series	same
Vegetation complex	same dominant species
<i>Site history</i>	
Timber type	same dominant tree species
History	same (year and type of site prep. and planting treatment)

The remaining gridpoints are systematically located at 10-m intervals. Gridpoints are marked with either pin-flags or wooden pickets, which are red in the Control and blue in the Treatment plot.

The closest, healthy tree of the target crop species to each gridpoint serves as the subplot centre. The target crop species must be well represented in both the Treatment and Control to ensure that 36 subplots can be located in each. Overlap of subplots is acceptable because competitive effects are evaluated on individual tree basis. When selecting target crop trees, keep in mind the objectives of the treatment. A conifer release prescription may specify, for example, the minimum size of crop tree that is treated. The distance and azimuth from the gridpoint to the target crop tree are recorded at the time of crop tree selection, to facilitate both relocation and development of the installation map (see Section 6).

The presence of two target crop species planted in discrete areas of an opening may necessitate the establishment of two separate installations. Where the two target crop species are intimately mixed, however, one installation may be used to evaluate both. In this situation, a subplot is centred on each target crop tree (i.e., two subplots) so that each Treatment and Control plot has 72 rather than 36 subplots.

The tree-centred subplots are permanently marked with rebar stakes located 50 cm due north of the target seedling. Where two target species are intimately mixed, a rebar stake is used to mark each subplot. The stakes are 1 m tall and hooked at the top end to reduce the risk of damage to wildlife and cattle. The hooked end of each stake is painted (red in Control, blue in Treatment), flagged with winter-weight pink, and marked with an aluminum tag. The aluminum tags identify the installation number, treatment, and subplot number (consecutive 1–36, or where two species are intimately mixed, 1–72). Each target crop tree and target non-crop specimen (see Section 3.3.2) are also marked with aluminum tags, which are loosely wired onto a lateral branch. The aluminum tags are numbered consecutively throughout the installation. The first numbers are assigned to the target crop tree and target non-crop specimens in subplot 1 in the Control, and the last numbers to those in subplot 36 in the Treatment. To avoid injury to the treatment crew, rebar stakes and aluminum tags should not be installed in the Treatment plot until after the treatment has been completed.

3 ASSESSMENTS

3.1 Introduction

In all installations, assessments should be made of two community components: (1) crop trees; and (2) target non-crop vegetation. Habitat for wildlife or cattle may also be assessed using the procedures described by the Habitat Monitoring Committee (1990). Where the treatment is prescribed fire or mechanical site preparation, pre- and post-treatment forest floor and fuel conditions should be assessed using the procedures of Trowbridge *et al.* (1986) or Curran and Thompson,⁴ respectively. The results of those assessments are used to quantify the intensity of site preparation treatments. This section describes the sampling strategy for each component.

Assessments are made within two different sized circular crop tree centred subplots. Competitive effect of target non-crop vegetation is evaluated in either 10 m² ($r=1.78$ m) subplots when mean crop tree height is ≤ 2 m, or 20 m² ($r=2.52$ m) subplots when mean crop tree height is >2 m. Mean height is determined when crop trees are selected (see Section 2.4). The same subplot size is used for all subplots within an assessment period. Crop tree density and stocking, non-crop vegetation treatment efficacy, and wildlife habitat are evaluated in 50 m² ($r=3.99$ m) subplots.

3.2 Target Crop Trees

One target crop species is usually selected for each installation. Two may be selected, however, where the species are intimately mixed (see Section 2.4). The target species selected is defined by the program objectives, and may be either a conifer or a hardwood. Planted crop trees are preferred over naturals. Where naturally regenerated trees are selected, however, all target crop trees must be the same age.

3.2.1 Growth

Crop tree growth is assessed in all subplots in each of the Treatment and Control. Measurements are seedling height (cm), height increment (cm), root collar diameter (mm), crown diameter (cm), crown length (cm), vigour, condition, damage, and height (%) and competitive status relative to surrounding non-crop vegetation. If shoot elongation is still occurring at the time of the baseline assessment, height to the top whorl and previous year's height increment are measured. Vigour, competitive status, and condition of foliage, leader and stem are assessed with procedures described by Herring and Pollack (1985). Damage cause is described according to Silviculture Survey damage codes (B.C. Ministry of Forests 1989). It is

⁴ Curran and Thompson, 1991.

described only where it potentially can affect crop tree performance. The field data form (PROBE form 2) and details on recording conventions are located in Appendix 2. Vigour, condition, damage and competitive status codes are located in Appendix 3.

3.2.2 Density and stocking

Density of all conifer species and stocking of well-spaced and free-growing trees are determined in all subplots in each of the Treatment and Control. The subplots are 50 m² ($r=3.99$ m) and centred on the target crop tree. A modification of the standard Silviculture Surveys procedure is used (B.C. Ministry of Forests 1989). This involves completion of the FS 657 (Silviculture Survey) and FS 658 (Silviculture Survey Plot) forms (see Appendix 4 for forms and recording conventions). The FS 659 form (Silviculture Survey Summary) is not used because the data are summarized and analyzed using SAS programs (see Section 11).

3.3 Target Non-crop Vegetation

Species may be selected as targets for measurements if they (1) represent a competitive threat to crop trees, and (2) occur in at least 80% (i.e., 29 out of 36) of the subplots in each of the Treatment and Control. Target species are selected following an assessment of species composition in the Treatment and Control plots. To do this, all potential non-crop species present in each subplot are tallied at the time of crop tree selection. A minimum of one and maximum of five target non-crop species are selected.

In some cases, the abundance of a non-crop species that was not selected as a target in the baseline assessment can increase dramatically after treatment, posing a competitive threat to crop trees. Clearly, pre- and post-treatment measurements of such species are beneficial, although it is impractical to measure all species based on the possibility of their increased abundance. Rather, Research Silviculturists can identify non-crop species that are *likely* to increase in abundance following treatment, before the baseline assessment, using their experience and results of previous trials. With this approach, some important species shifts may be missed. However, the abundance, vigour and distribution of all vascular plant species will be monitored in the eight subplots serving as Habitat Monitoring plots (see Section 3.4), which can provide sufficient information for the qualitative assessment of species shifts.

3.3.1 Competitive effect

The competitive effect of target non-crop species on target crop tree performance is assessed in a 10 or 20 m² circular subplot (10 m² if crop tree ≤ 2 m tall; 20 m² if crop tree > 2 m tall), which is centred on the target crop tree. The modal height, cover and density (hardwoods only) of target non-crop vegetation are assessed in all tree centered subplots in each of the Treatment and Control. More specifically, the following are measured: subplot total cover (%) and subplot modal height (cm); for each target species, cover (%) and modal height (cm); and for target hardwood species (i.e., paper birch, trembling aspen and black cottonwood), density (stems per plot). The field data form (PROBE form 3) and details on recording conventions are in Appendix 2.

3.3.2 Efficacy of treatment

Efficacy of treatment is evaluated on one tagged specimen of each woody target non-crop species located within each 50 m² crop tree-centred subplot. Usually, the closest, healthy specimen to the crop tree is chosen. However, care must be taken to ensure that the specimen meets the target criteria (e.g., minimum size) specified in the treatment prescription. For example, where the prescription calls for girdling all hardwood stems > 5 cm dbh, select only target specimens with at least 5 cm dbh. Non-woody species are not tagged for this evaluation because of the difficulty in relocating specimens from year to year. Rather, treatment efficacy at controlling non-woody species is evaluated simply as change in percent cover and modal height (see Section 3.3.1).

In baseline assessments only, the location (distance [dm] and azimuth) of each tagged woody specimen is described. In baseline and subsequent re-assessments, the total height (cm), leader height (cm),

vigour, condition and phenology (see Appendix 3) are recorded. Leader height may be difficult to determine, particularly if elongation is not complete at the time of the assessment. Where elongation appears to be 80% complete, record the current year's leader height. Otherwise, record the previous year's leader height. Leader condition is assessed for the leader whose height is measured. Phenology recorded before treatment assists in the interpretation of efficacy results, and that recorded after treatment provides a qualitative assessment of treatment-imposed stress. In re-assessments only, the number and length (cm) of sprouts (all woody shrubs and hardwoods) or suckers (aspen only), damage cause, and ECW rating are also recorded. The field data form (PROBE form 4) and details on recording conventions are in Appendix 2.

3.4 Wildlife Habitat (Optional)

All vascular plant species can be monitored throughout the 10-year re-measurement period according to procedures recommended by the Habitat Monitoring Committee (1990). They are monitored in four 50 m² crop tree centred subplots ($r=3.99$ m), which are randomly selected from the 36 subplots in each of the Treatment and Control (total of eight plots per installation). This entails completion of the Project Description Form (EM-1), Site Description Form (EM-2), Vegetation Description Form (EM-3) and Herbicide/Fertilizer Form (EM-5). The field forms are given in Appendix 5. For details on completing the field forms, refer to the Habitat Monitoring Committee (1990).

Form EM-1 is used to describe the location, goals and objectives of the project area, and Form EM-2 to describe the ecosystem and habitat monitoring sample plot layout.

Form EM-3 is used to record the vegetation characteristics of the habitat monitoring sample plots. For each species, the stratum, percent cover, distribution and vigour are recorded. Optional attributes that may also be recorded include phenology, impact assessment, stem count, total percent cover over all strata, strata height, and the number of dead trees.

On form EM-5 are described treatment application details (e.g., method of treatment, environmental conditions at time of treatment), as well as the impact and recovery of a maximum of six dominant shrub species. The form was designed to record herbicide and fertilizer treatment assessments, but is used for all treatments assessed in PROBE installations. Impact and recovery of herbaceous species is assessed as in percent cover, and recorded on the Vegetation Description Form (EM-3).

Pre-treatment assessment involves completion of forms EM-1, EM-2, EM-3, and EM-5A. Post-treatment assessment involves completion of forms EM-3 and EM-5B (Impact Assessment) at the end of the first full growing season following treatment (i.e., first re-assessment), and EM-5C (Recovery Assessment) at the end of the third (i.e., second re-assessment) and fifth (i.e., third re-assessment) growing seasons. Assessments in the selected habitat plots are made of the same target non-crop species evaluated using PROBE form 4. The Impact Assessment consists simply of plot tallies of the number of stems of each target shrub and hardwood species. The Recovery Assessment consists of plot tallies of the number of stems and basal sprouts in each pre-defined Assessment Category.

3.5 Prescribed Fire (Optional)

Pre- and post-burn conditions should be assessed using procedures modified from Trowbridge *et al.* (1986). Modification to those procedures is in the sample design only, not in the data collection. This section provides a brief summary of the procedures; for more detail, refer to the original handbook by Trowbridge *et al.* (1986). A copy of the fuel assessment form is in Appendix 6.

Before the burn, environmental conditions and vegetation are assessed in both the Treatment and Control according to procedures described in Sections 3.1, 3.3 and, optionally, 3.4. Gridpoints serve as subplot centres. In the Treatment plot only, pre- and post-burn estimates of slash loading, forest floor depth and mineral soil exposure also are made. Re-measurements are done as soon after the burn as possible because rain or snow can disturb the forest floor.

3.5.1 Sample design

Slash load and mineral soil exposure are estimated by the line intersect method, where each of the six gridlines serves as a 50-m line transect (i.e., $n=6$). In addition, forest floor depth is estimated and burn-pins inserted at each of the 36 systematically located gridpoints.

3.5.2 Slash load

Pre- and post-burn slash load is estimated along each of the six 50-m gridlines (transects). The slope is determined at the beginning of each gridline. The number of slash pieces (twigs, branches, small stems) ≤ 7 cm in diameter are then tallied according to the following size classes and sample line lengths:

Diameter size class (cm)	Portion of sample line tallied (m)
0–0.5	0–5
0.6–1.0	0–10
1.1–3.0	0–15
3.1–5.0	0–20
5.1–7.0	0–25

The size class is the actual diameter of each piece at the point of intersection with the gridline. The percent species composition of all pieces ≤ 7 cm in diameter is estimated after the count for a particular line is completed. The diameter and species of all intersecting pieces > 7 cm in diameter are determined along the entire 50-m gridline. To relocate the same measurement point on a piece after the burn, a nail is driven into that point, leaving approximately 1 cm of the nail exposed. Special tally rules for pieces of all diameters are summarized in Appendix 6.

After the burn, sampling of the ≤ 7 cm size classes is repeated according to the same procedures as in the pre-burn. All > 7 cm nailed pieces are tallied with the pre-burn procedures, even if the pieces are smaller than 7 cm or have moved off the gridline. Exposed roots, or pieces that are not nailed but have moved onto the gridline, are not measured.

3.5.3 Forest floor and mineral soil exposure

Before the burn, four depth-of-burn pins are inserted at each of the 36 gridpoints (total of 144 pins). The four pins are located 1 m from the gridpoint, one in each cardinal direction. The pins are placed in undisturbed forest floor so that the crossbar is level with the surface of the litter, moss layer or decayed log. The depth of litter or moss layer is measured where the depth-of-burn pins are inserted.

After the burn, the depth-of-burn pins are re-located and the ash is scraped away. The bottom of the crossbar to the top of the unburned forest floor, and the top of the unburned forest floor to the mineral soil surface, are measured.

To estimate percent mineral soil exposure along the full length of each gridline, the length of exposed mineral soil is measured, divided by 50 and then multiplied by 100.

3.5.4 Planting

After post-burn fuel sampling, the opening is operationally planted. In the Treatment plot, gridpoints are relocated by a compass and chain and marked with pin-flags or wooden stakes. In the Treatment and Control plots, the closest seedling to each gridpoint is selected as the target crop tree. The tree-centred subplots are marked with rebar stakes, located approximately 50 cm due north of the target trees.

3.6 Mechanical Site Preparation (Optional)

Soil disturbance after mechanical site preparation should be measured according to a modification of the Gridpoint-Intercept method described by Curran and Thompson.⁵ This section briefly summarizes the procedures; for more detail, refer to the original handbook.⁶ The soil disturbance survey forms and Soil Disturbance Cause codes are in Appendix 7.

⁵ Curran and Thompson, 1991.

⁶ *Ibid.*

Before mechanical site preparation, environmental conditions and vegetation are assessed in both the Treatment and Control according to procedures described in Sections 3.1, 3.3 and, optionally, 3.4. Gridpoints serve as subplot centres. After site preparation and planting, soil disturbance is assessed in both the Treatment and Control plots.

3.6.1 Planting

After mechanical site preparation is completed, the opening is operationally planted. In the Treatment plot, gridpoints are relocated by a compass and chain and marked with pin-flags or wooden stakes. In the Treatment and Control plots, the closest seedling to each gridpoint is selected as the target crop tree. The tree-centred subplots are marked with rebar stakes, located approximately 50 cm due north of the target trees.

3.6.2 Sampling design and disturbance survey

After planting is completed, soil disturbance is estimated along the six gridlines (i.e., $n=6$). It is evaluated at four observation points in each tree-centred subplot. The four points are located 1 m from the target tree, one in each cardinal direction. The Soil Disturbance Cause code is recorded for each point on Form 1-MSP (grid intercept survey). In addition, slope gradient is measured at the beginning of each gridline. The survey results are summarized separately for the Treatment and Control on Form 3A-MSP (Calculation Card Gridpoint-Intercept). The MSP forms are in Appendix 7.

4 RECORDS

PROBE and Silviculture Survey forms must be completed for each basic installation. As options, habitat monitoring, prescribed fire assessment, and mechanical site preparation soil disturbance survey forms may also be completed. The list of forms and the appendices in which they and their coding instructions are located are given in Table 3.

5 PHOTOGRAPHS

Photographs are taken during the baseline assessment, immediately after treatment, and at each post-treatment assessment. Coloured, 35mm slide film and a wide-angle lens (28–35mm) are used. The date the photograph was taken, installation number, installation location, treatment (T or C), subplot number and subject must be recorded on each slide. The information can be written either directly on the slide after it has been developed, or on a whiteboard which is included in the picture.

A minimum of four photographs are taken in each of the eight habitat monitoring sample plots. During each assessment, photographs are taken from the same points, which are recorded during the baseline assessment (see photographic record in Appendix 2). The points should be carefully chosen to afford a good view of the subject. The four photographs show: (1) target crop tree, (2) general habitat condition, (3) target non-crop specimens (more than one photograph may be necessary, depending on the proximity of the specimens to one another), and (4) overview of ground cover (optional). In addition, one photograph is taken toward the centre of the plot from each of the four corner posts in the Treatment and Control.

6 MAPPING

A location (1:20 000) and installation (1:1000) map are drafted for each installation. The location map shows the location of the installation on the appropriate Forest Cover map. The map notation must be permanently included on the Forest Cover mylar in the appropriate Forest District. The installation map shows the POC, Treatment plot, Control plot, gridpoints, subplots and important topographical and man-made features.

TABLE 3. List of forms for operational brushing evaluations

Form number	Description	Appendix
<i>Mandatory</i>		
PROBE-1	Project description	2
PROBE-2	Crop tree growth	2
PROBE-3	Target non-crop vegetation	2
PROBE-4	Target specimens	2
PROBE-5	Photographic record	2
FS 657, 658	Silviculture survey	4
<i>Optional</i>		
<i>Habitat Monitoring:</i>		
EM-1	Project description	5
EM-2	Site description	5
EM-3	Vegetation description	5
EM-5A	Herbicide/fertilizer	5
EM-5B	Herbicide impact assessment	5
EM-5C	Herbicide recovery assessment	5
<i>Prescribed Fire Assessment:</i>		
PF-4	Fuel assessment	6
<i>Mechanical Site Preparation Soil Disturbance Survey:</i>		
1-MSP	Grid intercept survey	7
3A-MSP	Calculation Card Gridpoint-Intercept	7

7 TIMING

Ideally, sites should be selected during the fall before installation establishment. Establishment and baseline assessments are best done just before treatment, and hence will depend on the type of treatment being imposed. For example, assessments may best be done in early summer (e.g., mid-late July) for brushing treatments or late summer (e.g., mid-late August) for site preparation treatments. Subsequent assessments are conducted during the period of full vegetative development and before frost incidence (i.e., between July 15 and August 31). They are conducted five times over a 10-year period:

1. Baseline — immediately before treatment.
2. First re-assessment — one growing season following treatment.
3. Second re-assessment — three growing seasons following treatment.
4. Third re-assessment — five growing seasons following treatment.
5. Fourth re-assessment — 10 growing seasons following treatment.

All measurements are completed during each re-assessment, except crop tree stocking and density, which are assessed only during the baseline, third and fourth re-assessments.

8 MAINTENANCE

Signs are erected on or near the POC at the time of plot installation. Maintenance of the installations is done during each re-assessment. At that time, missing corner posts, subplot centres and tags are replaced. Corner posts and subplot stakes are re-painted and re-flagged where necessary.

9 PERSON-DAY REQUIREMENTS

The time estimate for establishment of a new PROBE installation is given in Table 4. An extra day or two will be required if a prescribed burn assessment or mechanical site preparation soil disturbance survey is required. The time estimate for a re-assessment is 2–3 days, depending on whether or not a silviculture survey is required. The estimates are based on a two-person crew familiar with the sampling procedures.

TABLE 4. Estimate of crew-day requirements (2 persons) per installation

Activity	Crew-days (10-hour day)
Site location, Treatment and Control boundary layout	1
Subplot layout, crop and target specimen tagging	1
Crop tree and target non-crop specimen assessments (PROBE-2, -4)	0.5–1
Percent cover, modal heights (PROBE-3), posts, POC	0.5–1
Habitat assessments (all EM forms), photos	1
Silviculture survey (FS 567 and 658)	1
Rebar installation, site description, access notes (PROBE-1)	0.5
Approximate number of crew-days	5–7

10 RESPONSIBILITIES

The establishment, monitoring and data entry of PROBE installations are contracted out. The Stand Tending Co-ordinators are responsible for the administration of the contracts and co-ordination of the program. The specific responsibilities of the co-ordinators are:

- identification of program objectives, through the construction of the sampling matrix;
- approval of candidate sites, on the basis of program objectives;
- selection of qualified contractors;
- administration of contracts;
- maintenance of records library, including PROBE and habitat monitoring databases, annual reports, and slides (the habitat monitoring information is also included in a provincial database, which is maintained by the Habitat Monitoring Committee [1990];
- dissemination of annual reports to the Forest Districts; and
- co-ordination of field trips with District Silviculturists to discuss monitoring results.

The Research Silviculturists are responsible for annual training, consulting, analysis and reporting; and the District Silviculturists, or appropriate designates, are responsible for selection of candidate sites for PROBE installations.

The PROBE Contractors are responsible for site selection (with assistance from District Silviculturists), layout, assessment and data entry. In addition, the PROBE Contractors must maintain contact with the District

Silviculturists and the brushing or site preparation contractors throughout the establishment and treatment phases, to ensure that Control plots are not mistakenly treated. As soon as a PROBE installation has been established, the PROBE Contractors must send the District Silviculturists the following: (1) a description of general location and opening number; (2) a Forest Cover map showing location of POC and Treatment and Control plots; and (3) a description of how the installation is marked on the ground (e.g., colour of corner posts, rebar and flagging).

11 ANALYSES

11.1 Introduction

The Research Silviculturists are responsible for data analysis and reporting within their region. Crop tree and non-crop vegetation data are analyzed with the use of *t*-tests, ANOVAs, regression techniques and contingency tables. The habitat data are summarized according to procedures described by the Habitat Monitoring Committee (1990). Prescribed fire and soil disturbance data are summarized according to procedures described by Trowbridge *et al.* (1986) and Curran and Thompson.⁷

11.2 T-tests

Within an installation, *t*-tests are used to compare crop tree performance and target non-crop vegetation abundance between the Treatment and Control. Subplots located within the Treatment and Control plots are used as treatment units (i.e., $n=36$). Because the treatments were not randomly assigned to randomly selected subplots, however, this represents “pseudo-replication” (Hurlbert 1984). The true treatment units are not the 36 subplots, but rather the Treatment and Control plots in which they are located, which are not replicated on-site. Consequently, the validity of the *t*-test depends on the assumption of no between-plot variation. The criteria for selection of comparable Treatment and Control plots help to ensure that between-plot variation is indeed minimal.

The variables that will be tested using *t*-tests are:

- Crop tree
 - total height, height increment, diameter, diameter increment, crown volume, relative height, stocking, density
- Non-crop vegetation
 - All vegetation
 - cover, modal height
 - Target species
 - cover, modal height, density (hardwoods only)
 - Tagged target specimen
 - total height, height increment, sprout number, sprout length, ECW control rating

where crown volume = crown length $\times$ (crown diameter/2)². Means and standard errors are summarized in tables and boxplots.

11.3 Analysis of Variance

Each complex/ecosystem/treatment combination (cell) will be replicated three times (i.e., three installations). Once this has been accomplished, the combination is analyzed as a RDB using two-way ANOVA (see Appendix 8 for ANOVA table). With data pooled from three or more installations, broad inferences may be made about a particular treatment across a wide range of sites.

⁷ Curran and Thompson, 1991.

The variables which will be tested using ANOVA are:

- Crop tree - relative height growth rate, relative diameter growth rate, stocking, density
- Non-crop vegetation
 - All vegetation - cover, modal height
 - Target species - cover, modal height, density (hardwoods only)
 - Tagged target specimen - total height, height increment, sprout number, sprout length, ECW control rating

where relative height (or diameter) growth rate = $(\text{height}_{(x)} - \text{height}_{(x-1)})/\text{height}_{(x-1)}$. Means and standard errors are summarized in tables and boxplots.

11.4 Regression Analysis

For each installation, simple regression models are constructed which relate the performance of crop trees (dependent variable) to a competition index (independent variable). The crop tree performance variables tested include height, height increment, diameter, crown volume, and vigour. The competition indices are calculated according to the non-crop vegetation abundance measures (see Section 3.3.1). The indices are expected to reflect the amount of competition a target crop tree experiences from neighbouring vegetation. The competition indices which will be tested are given in Table 5.

The best models are selected based on the distribution of residuals and the coefficient of determination (r^2).

TABLE 5. Competition indices

Competition index	Description
C_T	Subplot total cover
H_T	Subplot modal height
CH_T	$C_T \times H_T$
CRH_T	$CH_T/\text{crop tree height}$
C_{Sx}	Species specific cover
H_{Sx}	Species specific modal height
D_{Sx}	Species specific density (hardwoods)
CH_{Sx}	$C_{Sx} \times H_{Sx}$
DH_{Sx}	$D_{Sx} \times H_{Sx}$
CRH_{Sx}	$CH_{Sx}/\text{crop tree height}$
DRH_{Sx}	$DH_{Sx}/\text{crop tree height}$

11.5 Contingency Tables

Contingency tables are constructed to investigate relationships among the following categorical variables:

- Crop tree - vigour, condition, damage cause, and competitive status
- Non-crop specimen - condition and damage cause

12 DATA MANAGEMENT

The Research Silviculturists are responsible for data management within their region. The PROBE Contractors must send the Research Silviculturists floppy discs and either originals or clear photocopies of the field forms. The Contractors should retain one copy of all data on file. Results from all installations will be summarized annually, and reports made available through the Research Silviculturists and Stand Tending Coordinators in the Kamloops and Nelson Forest Regions.

Data collected on PROBE forms (PROBE-1, PROBE-2, PROBE-3, and PROBE-4) and Silviculture Survey cards (FS 657 and 658) are entered into a database using PROBE Data Entry System Software, which is an application of Alpha Four (Alpha Software Corporation 1991). The data are analyzed by the Research Silviculturists using SAS programs (SAS Inc. 1985) specifically developed for PROBE. Data collected on Habitat Monitoring forms (EM-1, EM-2, EM-3, EM-5A, EM-5B, EM-5C) are entered and summarized using the Habitat Monitoring Data Base System (Habitat Monitoring Committee 1990).

13 PERSONNEL QUALIFICATIONS

Typically, work can be completed by a two-person crew consisting of one professional and one technician. The professional must have experience in layout of research trials, field forestry sampling, ecosystem classification, and plant identification. Experience in evaluating brushing operations and prescriptions is an asset, and an education background in forestry (e.g., B.S.F. or R.P.F.) or botany (e.g., B.Sc.) is preferred. The technician must be familiar with field forestry sampling.

14 LIST OF MATERIALS

Supplies:

- pressure-treated fence posts (POC and corner posts, 9 per installation)
- 1m tall rebar (sample plot centres, 72 per installation)
- wooden stakes or pin-flags (gridpoints, 72 per installation)
- blue and red tree-marking paint
- metal tags (~200 per installation)
- wire
- nails or staples
- blue, pink, and yellow or lime-green flagging
- numbered aluminum tags (no. target species $\times$ 72 subplots per installation)
- film (~60 exposures per installation)
- data forms

Equipment:

- compass
- clinometer
- 50m nylon chain or tape
- hip chain + thread
- calipers
- height pole
- metric scale rule
- 1.78, 2.52 and 3.99m plot cords
- shovel
- axe
- plant press
- 35mm camera + wide-angle lens
- whiteboard + felt pen
- clipboard

Maps and references:

- topographic map
- subzone map
- forest cover map
- planting unit map
- site history records
- site identification guide (Lloyd *et al.* 1990; Braumandl and Curran 1992)
- plant guides (e.g., Angove and Bancroft 1983)

APPENDIX 1. Species composition of major vegetation complexes in the southern Interior of British Columbia (from Kimmins and Comeau 1990)

Species composition of major complexes of competing vegetation in the southern Interior (from Kimmins and Comeau 1990). (Zones indicated in parentheses are those in which the complex has limited occurrence.)

Complex	Major species	Biogeoclimatic zone	Sites
Pinegrass	<i>Calamagrostis rubescens</i>	IDF, ICH, MS, PPBG, SBS, ESSF	Dry to Fresh
Fireweed	<i>Epilobium angustifolium</i>	(IDF), ICH, MS, ESSF	Dry to Wet
Fern	<i>Athyrium filix-femina</i> <i>Dryopteris assimilis</i> <i>Pteridium aquilinum</i>	ICH	Fresh to Wet
Dry Alder	<i>Alnus viridis</i> ssp. <i>sinuata</i> <i>Epilobium angustifolium</i> <i>Calamagrostis rubescens</i>	ICH, MS, SBS	Dry to Fresh
Wet Alder	<i>Alnus viridis</i> ssp. <i>sinuata</i> <i>Alnus incana</i> ssp. <i>tenuifolia</i> <i>Rubus parviflorus</i> <i>Lonicera involucrata</i> <i>Calamagrostis canadensis</i> <i>Athyrium filix-femina</i> <i>Dryopteris assimilis</i>	ICH, (MS), ESSF	Wet
Mixed Shrub	<i>Rubus parviflorus</i> <i>Rubus ideaus</i> <i>Lonicera involucrata</i> <i>Acer glabrum</i> <i>Alnus viridis</i> ssp. <i>sinuata</i> <i>Paxistima myrsinites</i> <i>Sambucus racemosa</i> <i>Oplapanax horridus</i> <i>Menziesia ferruginea</i> <i>Salix</i> spp. <i>Cornus sericea</i> <i>Athyrium filix-femina</i> <i>Pteridium aquilinum</i>	ICH, ESSF	Fresh to Wet
Dry Shrub	<i>Ceanothus velutinus</i> <i>Paxistima myrsinites</i> <i>Shepherdia canadensis</i> <i>Symphoricarpos albus</i> <i>Holodiscus discolor</i> <i>Physocarpus malvaceus</i>	ICH, IDF	Dry to Fresh
Ericaceous Shrub	<i>Rhododendron albiflorum</i> <i>Menziesia ferruginea</i> <i>Vaccinium membranaceum</i>	ICH, IDF	Dry to Moist
Subalpine Herbs	<i>Valeriana sitchensis</i> <i>Luzula</i> spp. (<i>Epilobium angustifolium</i>)	ESSF	Fresh to Wet
Aspen	<i>Populus tremuloides</i>	IDF, ICH, MS	Various

APPENDIX 1. (continued)

Complex	Major species	Biogeoclimatic zone	Sites
Cottonwood-Shrub	<i>Populus balsamifera</i> <i>Lonicera involucrata</i> <i>Cornus sericea</i> <i>Sambucus racemosa</i> <i>Rubus parviflorus</i> <i>Alnus viridis</i> ssp. <i>sinuata</i> <i>Alnus incana</i> ssp. <i>tenuifolia</i> <i>Calamagrostis canadensis</i>	IDF, ICH	Floodplains
Mixed Hardwood-Shrub	<i>Populus tremuloides</i> <i>Populus balsamifera</i> <i>Betula papyrifera</i> <i>Salix</i> spp. <i>Alnus viridis</i> ssp. <i>sinuata</i> <i>Alnus incana</i> ssp. <i>tenuifolia</i>	IDF, ICH, MS	Various
Willow	<i>Salix</i> spp.	IDF, ICH, MS, ESSF	Moist to Wet
Introduced Grasses	Domestic grasses	(PPBG), IDF, (ICH), MS, (ESSF)	Various

APPENDIX 2. PROBE forms and coding instructions

PROBE Form 1: Project Description

Date:	Year-month-date of original data collection.
Project number:	Number assigned by the project co-ordinator.
Project co-ord.:	Name/position title.
Assessors:	Name/company.
Agency:	Ministry, licensee or other group responsible for the installation.
Region:	Code to designate forest region.
District:	Code to designate forest district.
Mapsheet/opening:	Forest Cover mapsheet and opening number.
Origin:	Target crop tree origin (P=planted, N=natural).
Location:	Brief description of where the installation is located in the geographic area and in the opening, and where the POC is located.
History symbol:	Harvesting, site preparation, planting, and stand tending. Record year and method.
Planting history:	Planting administration number, report number and unit. Date (year and month) trees were planted. Species, seedlot, nursery, stock type and stock size.
Site description:	BGC subzone and ecosystem association (site series). Azimuth (aspect), slope, slope position, elevation, site class. Soil texture, soil type, humus type.
Target species:	Maximum of two crop species. Maximum of five non-crop species.
Subjective evaluation:	Was the treatment effective at reducing competition? Was the treatment effective at improving crop tree performance? Comments.

PROBE Form 1: Project Description

Date:

Project Number:

Project Co-ordinator: Name:

Position Title:

Assessors:

Name:

Company:

Agency:

Region:

District

Map sheet, opening no.:

Origin:

Location:

History symbol:

Planting history:

Planting admin. no.:

Report no.:

Unit:

Date planted:

Species:

Seedlot:

Nursery:

Stock type:

Stock size:

Site description:

BGC subzone / ecosystem association:

Azimuth:

Slope:

Slope position:

Elevation:

Site class:

Soil texture:

Soil type:

Humus type:

Target Species:

Crop species:

Non-crop species:

Subjective evaluation: (Pre-treatment: Is treatment needed? Post-treatment: Was treatment effective?)

PROBE Form 2: Crop Tree Growth

Date:	Year-month-date of original data collection.
Project number:	Number assigned by the project co-ordinator.
Location:	Geographical area in which the installation is located.
Treatment:	T (Treatment) or C (Control).
Subplot:	Subplot number (1–36).
Species:	Species, identified by first four letters of the genus and first three letters of the species (e.g., <i>Pseudotsuga menziesii</i> is PSEUMEN)
Total height:	If terminal bud set, measure to bottom of bud. If not set, measure to base of top whorl. Measured to nearest 1cm on a metre stick or 10cm on a height pole.
Leader height:	Measured from base of terminal bud to base of lower node. If terminal bud set, measure current year's height increment. If not set, measure previous year's height increment. Measured to nearest 1cm on a metre stick, or nearest 10cm on a height pole.
Crown diameter:	Average of two perpendicular measurements, measured to nearest 1cm.
Crown length:	Measured from the base of the terminal bud (or top whorl if bud not set) to the bottom of the lowest laterals which support needles or leaves. Measured to nearest 1cm on a metre stick, or 10cm on a height pole.
Stem diameter:	Measured at root collar to nearest 0.1cm using calipers.
Relative height:	Height of crop tree relative to modal height of surrounding vegetation (%).
Competitive status:	See Appendix 3.
Condition/damage:	See Appendix 3. Only record damage cause if it is considered to be a threat to crop tree performance.
Vigour:	See Appendix 3.

Page __ of __

Treatment : 2082

PROBE Form 3: Target Non-crop Vegetation

Date:	Year-month-date of original data collection.
Project number:	Number assigned by the project co-ordinator.
Location:	Geographical area in which the installation is located.
Treatment:	T (Treatment) or C (Control).
Subplot:	Subplot number (1–36).
Species:	In all subplots, percent cover and height are measured for all potentially competitive vegetation, all hardwoods, all shrubs, and all herbs. They are also measured for up to five selected target species. The target species are identified by the first four letters of the genus and first three letters of the species (e.g., <i>Betula papyrifera</i> is BETUPAP).
Cover:	Estimated to nearest 5%.
Height:	Modal height is estimated to nearest 10cm. It is the most frequent height of individuals in the subplot.
Density:	Number of stems in the subplot for hardwoods (> 10cm tall) only. Hardwoods include <i>Betula papyrifera</i> , <i>Populus tremuloides</i> , and <i>Populus balsamifera</i> ssp. <i>trichocarpa</i> .

PROBE Form 3: Target Non-crop Vegetation

Date:

Page of

Project Number:

Location:

Treatment number:

Subplot	Species	Cover (%)	Height (cm)	Density (#/plot)
	A L L			
	H A R D W O S			
	S H R U B S			
	H E R B S			
	A L L			
	H A R D W O S			
	S H R U B S			
	H E R B S			
	A L L			
	H A R D W O S			
	S H R U B S			
	H E R B S			
	A L L			
	H A R D W O S			
	S H R U B S			
	H E R B S			

206/07

PROBE Form 4: Target Specimens

Date:	Year-month-date of original data collection.
Project number:	Number assigned by the project co-ordinator.
Location:	Geographical area in which the installation is located.
Treatment:	T (Treatment) or C (Control).
Subplot:	Subplot number (1–36).
Species:	Species with woody stems only. The species are identified by the first four letters of the genus and first three letters of the species (e.g., <i>Betula papyrifera</i> is BETUPAP).
Distance:	Distance from the main stem of the target crop tree to the main stem (or middle of clump) of the target specimen, measured to nearest dm.
Azimuth:	Azimuth from crop tree to middle of target specimen.
Total height:	Height to top of tree. Measured to nearest 1cm on a metre stick, or 10cm on a height pole.
Height increment:	Height increment can be difficult to measure because elongation may not yet be complete and last year's node may be hidden. Where elongation appears to be 80% complete, record the current year's leader height; otherwise record previous year's leader height. Measured to nearest 1cm on a metre stick, or 10cm on a height pole.
Sprout number:	Total number of adventitious shoots.
Sprout length:	Average sprout length to nearest 10cm.
Vigour:	See Appendix 3.
Condition/damage:	See Appendix 3. If two or more damaging agents occur on the same specimen, record the most damaging agent. Only record damage cause if it is considered to be a threat to target specimen performance.
Phenology:	See Appendix 3.
ECW rating:	Linear scale of 0–100, in which 0% is healthy and 100% is dead.

Page 1 of 1

Treatment number:

20640

PROBE Form 5: Photographic Record

Date: Year-month-date of original data collection.

Project number: Number assigned by the project co-ordinator.

Location: Geographical area in which the installation is located.

Treatment: T (Treatment) or C (Control).

Corner photos: Number of the photo taken at each of the four corners in the Treatment and Control plots.

Habitat plot photos

Subplot number: Number of subplot.

Photo number: Photo number (given on slide jacket).

Photo azimuth: Direction of photo.

Location: Location in which the photo was taken relative to the crop tree (distance and azimuth from camera to crop tree).

Probe Form 5: Photographic Record

Date: _____ Page: _____ of: _____

Project Number: _____ Location: _____ Treatment: _____

Corner Photos:

Corner	Photo number
SE	_____
SW	_____
NE	_____
NW	_____

Habitat Plot Photos:

Subplot number: _____ Photo number: _____ Photo azimuth: _____

Distance/azimuth from camera to crop tree: _____

Crop tree condition: _____

General habitat: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Subplot number: _____ Photo number: _____ Photo azimuth: _____

Distance/azimuth from camera to crop tree: _____

Crop tree condition: _____

General habitat: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Subplot number: _____ Photo number: _____ Photo azimuth: _____

Distance/azimuth from camera to crop tree: _____

Crop tree condition: _____

General habitat: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Subplot number: _____ Photo number: _____ Photo azimuth: _____

Distance/azimuth from camera to crop tree: _____

Crop tree condition: _____

General habitat: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

Target species: _____

APPENDIX 3. Codes for competitive status, vigour, condition, damage, phenology, and tree symbols

Competitive Status Codes (from Herring and Pollack 1985)

Competitive status is a qualitative assessment of the height of the target crop tree relative to the height of surrounding vegetation. The three codes are:

- 1 **Overtopped.** Leader of crop tree is currently overtopped by surrounding vegetation.
- 2 **Threatened.** Leader of crop tree is roughly equivalent to height of surrounding vegetation. It is likely to be overtopped within two growing seasons.
- 3 **Free-to-grow.** Leader of crop tree is well above surrounding vegetation and is unlikely to become threatened.

Vigour Codes (from Herring and Pollack 1985)

Crop tree vigour is a qualitative assessment which integrates shoot growth, caliper thickness, leaf area, needle length and foliage colour. The five codes are:

- 0 **Dead.**
- 1 **Moribund.** Near death; little or no visible shoot growth.
- 2 **Poor.** Little or possibly etiolated shoot growth; few and/or short needles; small caliper.
- 3 **Moderate.** Moderate shoot growth, leaf area, needle length, and caliper.
- 4 **Good.** Vigorous shoot growth; high leaf area; long needles; deep green colour; thick caliper.

Condition Codes (modified from Herring and Pollack 1985)

Foliage Condition

A	Defoliated
B	Browsed
D	Dead lateral buds
E	Semi-chlorotic
F	Decomposing
G	Galls
H	Healthy
L	Delignification
M	Mottled
N	Necrotic
O	Other
R	Reflushed
V	Stunted needles
Y	Chlorotic

Leader Condition

A	Missing
B	Browsed
C	Curled
E	Etiolated
F	Forked
H	Healthy
N	Not flushed
O	Other
R	Reflushed
S	Broken
T	Dead terminal bud
Z	Stunted needles on leader

Stem Condition

A	Girdled
B	Bent
C	Cut
D	Dead
F	Forked
G	Gall
H	Healthy
K	Crooked
L	Leaning
M	Piece missing (e.g., scar, lateral missing)
O	Other
P	Peeled
S	Smashed, crushed or trampled
U	Buried
Z	Blistered

Damage Cause Codes (from B.C. Ministry of Forests 1989)

Animal Damage

AB	Bear
AC	Cow
AD	Deer
AE	Elk
AH	Hare or rabbit
AM	Moose
AP	Porcupine
AS	Squirrel
AU	Unknown
AV	Vole
AW	Sheep
AX	Bird
AZ	Beaver

Insects

IA	Aphids
IB	Bark beetles
ID	Defoliators
IP	Pitch mass insects
IS	Shoot insects
IU	Unknown
IW	Weevils
M	Mites

Non-Biological

CV	Vegetation competition
NB	Fire
ND	Drought
NF	Flooding
NG	Frost
NH	Hail
NK	Fumekill
NL	Slash/debris
NR	Redbelt
NS	Slide
NU	Unknown
NW	Windthrow
NY	Snow or ice
TC	Chemical
TD	Treatment damage
TL	Logging
TP	Planting
TR	Pruning
TS	Skidroad
TT	Thinning/spacing

Disease

DA	Foliage diseases
DD	Stem rot
DF	Brooming
DL	Leader and branch dieback
DM	Dwarf mistletoe
DP	Bark diseases (superficial)
DR	Root diseases
DS	Stem diseases
DU	Unknown

Phenology Codes (from Dierschke 1972)

Vegetative

Deciduous tree or shrub

- 0 Closed bud
- 1 Buds with green tips
- 2 Green leaf out but not folded
- 3 Leaf unfolding up to 25%
- 4 Leaf unfolding up to 50%
- 5 Leaf unfolding up to 75%
- 6 Full leaf unfolding
- 7 First leaves turn yellow
- 8 Leaf yellowing up to 50%
- 9 Leaf yellowing over 50%
- 10 Bare

Conifer

- 0 Closed bud
- 1 Swollen bud
- 2 Split bud
- 3 Shoot capped
- 4 Shoot elongate
- 5 Shoot full length, lighter green
- 6 Shoot mature, equally green

Generative

Deciduous and conifer

- 0 Without blossom buds
- 1 Blossom buds recognizable
- 2 Blossom buds strongly swollen
- 3 Shortly before flowering
- 4 Beginning flowering
- 5 In bloom up to 25%
- 6 In bloom up to 50%
- 7 Full bloom
- 8 Fading
- 9 Completely faded
- 10 Bearing fruit

Tree Symbols (from Lloyd *et al.* 1990)

Scientific name	Common name	Symbol
<i>Abies amabilis</i>	amabilis fir	Ba
<i>Abies lasiocarpa</i>	subalpine fir	Bl
<i>Betula papyrifera</i>	paper birch	Ep
<i>Betula occidentalis</i>	water birch	Ew
<i>Chamaecyparis nootkatensis</i>	yellow-cedar	Yc
<i>Larix lyallii</i>	alpine larch	La
<i>Larix occidentalis</i>	western larch	Lw
<i>Picea engelmannii</i>	Englemann spruce	Se
<i>Picea engelmannii</i> x <i>glauca</i>	hybrid white spruce	Sxw
<i>Picea glauca</i>	white spruce	Sw
<i>Picea mariana</i>	black spruce	Sb
<i>Pinus albicaulis</i>	whitebark pine	Pa
<i>Pinus contorta</i>	lodgepole pine	Pl
<i>Pinus monticola</i>	western white pine	Pw
<i>Pinus ponderosa</i>	ponderosa pine	Py
<i>Populus tremuloides</i>	trembling aspen	At
<i>Populus balsamifera</i> ssp. <i>trichocarpa</i>	black cottonwood	Act
<i>Pseudotsuga menziesii</i>	Douglas-fir	Fd
<i>Thuja plicata</i>	western redcedar	Cw
<i>Tsuga heterophylla</i>	western hemlock	Hw
<i>Tsuga mertensiana</i>	mountain hemlock	Hm

APPENDIX 4. Silviculture survey forms and coding instructions (from B.C. Ministry of Forests 1989)

FS 657: Silviculture Survey

Only a subset of the information on this form must be completed. That information is listed below under the major section headings. Basic Site Information does not need to be recorded on this form because it is recorded on PROBE-1 and EM-2.

Admin.:

Region, District, location, mapsheet/opening no., administration no. (=PROBE Project Number), examiner, date and history symbol.

Site and Survey Factors:

For objectives of survey, enter PROBE next to "other."

Ecological Classification:

BGC zone, subzone, variant, site series, edatopic grid.

Acceptable Tree Characteristics:

List all conifer species occurring within the installation.

Survey Parameters:

For target only, determine from the regional stocking standards the stocking standard/ha, maximum/plot, inter-tree spacing, and minimum stocking standard.

FS 658: Silviculture Survey Plot

Plot no.:	Subplot number (1–36).
Layer:	Planted or natural.
Total trees:	Determine by adding total conifers and hardwood density, which is recorded on PROBE Form 3.
Total conifers:	Total conifer density.
Species:	Record total density of each species in the plot.
Total W:	Number of well-spaced trees.
Total FG:	Number of free-growing trees, as defined in Appendix 3.

FS 659: Silviculture Survey Summary

This form is not used because the data is summarized using PROBE computer programs.

SILVICULTURE
SURVEY

USE THIS FORM IN THE FIELD TO ESTABLISH SPECIFIC STANDARDS FOR THE SURVEY RECORD PLOTS ON THE FORM FS 400. SUMMARIZE PLOTS ON FORM FS 400. REFERENCE CARD FS 400 PROVIDES PROCEDURES.

REGION	DISTRICT	LOCATION	BAF PLOT TOPENING NO.		
ADMINISTRATION NO.	LICENCE NO.	LICENCEE	C.F. NO.	BLOCK	HISTORY SYMBOL
STRATUM	STRATUM AREA (ha)	AIR PHOTO NO.	EXAMINER	DATE	T M Y

OBJECTIVES OF SURVEY (CHECK APPLICABLE BOX)	☐ STOCKING	☐ PLANTABILITY	☐ FREE GROWING	☐ BRUSHING	☐ OTHER _____
---	-----------------------------------	---------------------------------------	---------------------------------------	-----------------------------------	--------------------------------------

SITE	ELEVATION: MINIMUM _____ MAXIMUM _____ AVERAGE _____	ASPECT _____	SLOPE POS. _____	SLOPE %: MINIMUM _____ MAXIMUM _____ AVERAGE _____	TERRAIN: ☐ EVEN ☐ ROLLING ☐ GULLED ☐ BROKEN	DESIGNED SITE CLASS: ☐ GOOD ☐ MEDIUM ☐ POOR ☐ LOW
-------------	---	---------------------	-------------------------	---	--	--

SOIL	LPH (cm) _____	HUMUS FORM _____	SOIL TEXTURE _____	ROOTING DEPTH (cm) _____	SOIL DEPTH (cm) _____	DRAINAGE: ☐ GOOD ☐ FAIR ☐ POOR
	☐ COARSE FRAGMENTS	☐ LOW (< 5%)	☐ MODERATE (5-10%)	☐ HIGH (> 10%)		

ECOLOGICAL CLASSIFICATION FOR MOST PREDOMINANT AREA OF STRATUM	BOG ZONE _____	SUBZONE _____	VARIANT _____	SITE SERIES _____	EDAPHIC SRD: Moisture _____ Nutrients _____
--	-----------------------	----------------------	----------------------	--------------------------	--

ACCEPTABLE TREE CHARACTERISTICS					
CHECK (+/-)	A	U	SPECIES	STORAGE	QUALITY (Location, Free-Growing Criteria)
A = ACCEPTABLE U = UNACCEPTABLE					

SURVEY PARAMETERS	STOCKING STANDARDS ₉₀	MINIMUM PLANT	WATER TREE SPACING	MINIMUM STOCKING STANDARD	DISTANCE BETWEEN
To assess well spacing (m) or (m ²)	TARGET		REAL		PLOTS (m) LINES (m)
	PLANTING				RADIUS (m) AREA (m ²)
	EXPECTED SURVIVAL %	PLANTING STOCKING STANDARD	TARGET STOCKING STANDARD	EXPECTED SURVIVAL RATE (%)	NATURAL FILL IN RATE (m ² /ha)
	MULTIPLIER				

FILL ORIGINAL IN DISTRICT OFFICE ON OPENING FILE

875. 4007 1-4338 180xw



BASIC SITE INFORMATION

NOT DRAWN FROM ESTIMATED DURING WW TWO,
WAS A THOUGHT OF STRATUM

DESCRIPTION OF ACCEPTABLE PLANTABLE SPOTS

SCREENING DEPTH _____ (m) SCALP SIZE _____ (m) ACCEPTABLE PLANTING MEDIUM _____

TREE SPECIES PLANTERS MUST RECOGNIZE _____ MINIMUM HEIGHT FOR TREES TO BE WORKED _____ (m) PLANTING DIFFICULTY: (see FS 680 and 702) ☐ EASY ☐ MODERATE ☐ DIFFICULT ☐ SEVERE

VISUAL ESTIMATE OF SITE CONDITIONS

ESTIMATED SLASH COVER _____ % AVERAGE SLASH HEIGHT _____ (m)

SLASH CLASS ☐ HIGH ☐ MEDIUM ☐ LOW ☐ NL

SLASH DISTRIBUTION ☐ CONTINUOUS ☐ PATCHY ☐ SCATTERED

ESTIMATED BRUSH COVER _____ % AVERAGE BRUSH HEIGHT _____ (m)

BRUSH SEVERITY ☐ HIGH ☐ MEDIUM ☐ LOW ☐ NL

BRUSH DISTRIBUTION ☐ CONTINUOUS ☐ PATCHY ☐ SCATTERED

SNAGS ESTIMATED DENSITY _____ /ha

WORK HAZARD: ☐ HIGH ☐ MEDIUM ☐ LOW ☐ NL

MACHINE TRAFFICABILITY: ☐ GOOD ☐ MEDIUM ☐ LOW ☐ NL

SOIL COMPACTION HAZARD: ☐ HIGH ☐ MEDIUM ☐ LOW ☐ NL

INDICATE THE PRESENCE OR ABSENCE OF THE FOLLOWING:

WATER BODIES ☐ YES ☐ NO TYPE _____

HIGH WATER TABLE ☐ YES ☐ NO DEPTH _____

FISHERIES VALUES ☐ YES ☐ NO TYPE _____

WILDLIFE VALUES ☐ YES ☐ NO TYPE _____

RANGE USE ☐ YES ☐ NO TYPE _____

PEST DAMAGE ☐ YES ☐ NO TYPE _____

RECREATION VALUES ☐ YES ☐ NO TYPE _____

OTHER RESOURCE VALUES ☐ YES ☐ NO TYPE _____

ACCESS

	LAND				AIR		WATER
	DRD	4WD	ATV	WALK	PILOT	HEL	
DISTANCE							
	(m)				(km)		(km)

TRAVEL TIME

_____ HOURS FROM _____ LOCATION

POTENTIAL ROAD CONSTRUCTION AND MAINTENANCE PROBLEMS (Show details on map)

SLIDE - Year _____ (m) CULTIVETS Buried Length Diameter

WASHOUT - Length _____ (m) _____

NEW ROAD - Length _____ (m) _____

ACCOMMODATION ☐ HOTEL ☐ CAMPSITE (NAME: _____) ☐ NONE ☐ OTHER _____

GENERAL DESCRIPTION OF MAJOR LIMITING SITE FACTORS, TREATMENT OBJECTIVES AND NEARBY PEST OR SITE PROBLEMS:

FIG. 6.63 (continued)

SILVICULTURE
SURVEY PLOT

Page _____ of _____

[illegible]

COMMENTS

(prearrangement, an apt,
watercourse, wildlife
stated damage, etc.)

DATA ON THE LEADER HEIGHT, TOTAL HEIGHT AND AGE OF WELL SPACED TREES, MAIN COMPETING VEGETATION AND INVENTORY COMPONENT SHOULD BE RECORDED ON THE FIRST PLOT ESTABLISHED IN THE STANLAW, AND EVERY FOURTH PLOT THEREAFTER. PEST TYPE, INCIDENCE AND SEVERITY SHOULD BE NOTED IN THE COMMENTS SECTION, WHEN OBSERVED.

F8-608 SIL 90/4

APPENDIX 5. Habitat monitoring forms (from Habitat Monitoring Committee 1990)



Province of British Columbia
Ministry of Forests
Ministry of Environment

PROJECT DESCRIPTION FORM (EM-1) (FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

PROJECT IDENTIFICATION

1. PROJECT ID NAME

2. PROJECT ID NUMBER

3. PROJECT COORDINATOR

4. STARTING DATE

5. AGENCIES

PROJECT LOCATION

6. NTS SHEET

NTS SHEET

OR

BCGS (Forest Cover) Map No.

BCGS (Forest Cover) Map No.

7. MoE Sub-region

Management Unit / Subunit

8. MoF District

Name

Code

Stock Range

Name

Code

Range Unit

Name

Code

Pasture

Name

Code

9. GEOGRAPHIC LOCATION

PROJECT GOALS AND OBJECTIVES

10. SITE HISTORY

11. PROJECT GOALS (enter letters by priority, followed by species codes; maximum of 4)

PRIORITY 1

PRIORITY 2

PRIORITY 3

PRIORITY 4

A. Improve suitability for wildlife

B. Improve suitability for livestock

C. Improve suitability for timber

D. Reduce fire hazard

E. Other (specify in Comments)

12. PROJECT OBJECTIVES (enter letters by priority; maximum of 4)

PRIORITY 1

PRIORITY 2

PRIORITY 3

PRIORITY 4

A. Increase forage quantity

B. Increase forage quality

C. Increase forage availability

D. Change plant species composition

E. Alter range condition

F. Remove decadent plant material

G. Stimulate sprouting/suckering

H. Increase or maintain snow interception cover

I. Increase thermal cover

J. Increase security cover

K. Reduce volatile fuels

L. Improve animal movement

M. Stimulate germination/seeds

N. Improve conifer release

O. Increase timber growth rate

P. Reduce tree stocking (stems/hectare)

Q. Open forest canopy

R. Assess forage utilization

S. Assess herbicide effects

T. Control erosion

U. Provide nesting habitat

V. Other (specify in Comments)

13. HABITAT TECHNIQUE UTILIZED (enter letters by priority; maximum of 4)

PRIORITY 1

PRIORITY 2

PRIORITY 3

PRIORITY 4

A. Prescribed burning

B. Livestock grazing

- season long

C. Short duration grazing

D. Rest rotation grazing

E. Deferred rotation grazing

F. Reverse rotation grazing

G. No livestock grazing

H. Slashing

I. Site Rehabilitation

J. Juvenile spacing

K. Precommercial thinning

L. Commercial thinning

M. Coppice

N. Improvement cut (sanitation)

O. Selection cut

P. Shelterwood cut

Q. Seed tree cut

R. Clearcut

S. Tillage and scarification

T. Seeding and planting

U. Fertilizer application

V. Herbicide application

W. Water level manipulation

X. Other (specify in Comments)

PROJECT SUMMARY

14. SIZE OF TREATMENT AREA

ha

15. TOTAL PROJECT COST

16. COMPLETION DATE

Y

M

D

17. SUMMARY / EVALUATION REPORT

Available

Date

Y

M

18. COMMENTS



SITE DESCRIPTION FORM (EM-2) (FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

SERIES FORM

PROJECT IDENTIFICATION AND SAMPLING

① DATE Y M D ① PROJECT ID Name Number

② SURVEYOR ④ SITE NO.

⑤ SAMPLING (circle appropriate technique, type and timing)

<u>Technique</u>	<u>Type</u>	<u>Timing</u> (measurement number)
A. Reconnaissance with single macroplot only	A. Control	0. Initial measurement pre-treatment
B. Reconnaissance with single macroplot and nested microplots	B. Treatment	1. First measurement post-treatment
C. Multiplot with macroplots only		2. Second measurement post-treatment
D. Multiplot with macroplots and nested microplots		3. Third measurement post-treatment

LOCATION OF TIE POINT AND TIE LINE

6. TIE POINT LOCATION: PHOTO NO. and COORDINATES

Flight Line Photo # X Y

7. TIE POINT COMMENTS

8. TIE LINE BEARING 9. TIE LINE DISTANCE m

SAMPLE LOCATION (MACROPLOT 1)

⑩ NTS SHEET ⑪ BCGS (Forest Cover) Map No. Polygon No. ⑫ MACROPLOT RADIUS m

⑬ UTM SYSTEM Easting Northing 14. MACROPLOT LOCATION: PHOTO NO. AND COORDINATES

Zone Flight Line Photo # X Y

MULTILOT DETAILS

15. STRIP BEARING ⑬ MACROPLOT INTERVAL (distance between plots) m 17. SAMPLE STRIP COMMENTS

⑮ NUMBER OF MICROPLOTS per MACROPLOT 19. MICROPLOT RADIUS m 20. PHOTOGRAPHIC RECORD

BIOPHYSICAL DESCRIPTION

		MACROPLOT # 1	MACROPLOT # 2	MACROPLOT # 3	MACROPLOT # 4	* ECOLOGICAL MOISTURE REGIME
⑮ ASPECT (°)						
⑮ SLOPE (%)						B. arid
⑮ ELEVATION (m)						C. subarid
⑮ ECOLOGICAL MOISTURE REGIME (see*)						D. submoist
25. SURFACE SUBSTRATE (% cover)	decaying wood					E. moist
	bedrock					F. subhygic
	cobbles & stones					G. hygro
	mineral soil					H. subhygro
	organic matter					I. hygro
	water					
	TOTAL	100%	100%	100%	100%	

26. SITE POSITION MACRO 27. SITE SURFACE SHAPE 28. TERRAIN CLASSIFICATION

- A. apex
B. face
C. upper slope
D. middle slope
E. lower slope
F. valley floor
G. plain

- A. concave
B. convex
C. straight

- A. mapped
B. field interpreted

Year

TEXTURE	SOIL	CLIM. DATA	SURFACE	MOIST.	QUAL.	CLASS.
1.						
2.						
3.						

29. BIOGEOCLIMATIC CLASSIFICATION

Zone Sub Var Ph

30. SITE ASSOCIATION

Assoc. S Type P

31. FOREST COVER MAP

Primary Species

Secondary Species

*Age Class

*Height Class

*Stocking Class

*Canopy Closure

*Site Class

32. ECOLOGICAL CLASSIFICATION

Ecoregion

Ecosection

Biophysical Habitat Class

Phase

33. SOIL CLASSIFICATION

Year

Soil

- A. mapped
B. field interpreted

Percent (x 10)

Classification

Texture

Drainage

34. WILDLIFE CAPABILITY

Capability Rating

Map Year

Winter Range

Capability Rating

Summer Range

Capability Rating

Environmental

Conditions

HERBACEOUS STANDING CROP

MACROPLOT(MA) NO.	MA 1				MA 2			
MICROPLOT(MI) NO.	M1.1	M1.2	M1.3	M1.4	M2.1	M2.2	M2.3	M2.4
35. PRODUCTION ESTIMATE (g)								
36. PRODUCTION CLIPPED (g)								
MACROPLOT(MA) NO.	MA 3				MA 4			
MICROPLOT(MI) NO.	M3.1	M3.2	M3.3	M3.4	M4.1	M4.2	M4.3	M4.4
35. PRODUCTION ESTIMATE (g)								
36. PRODUCTION CLIPPED (g)								

- *AGE CLASS
1. 1-20 years
2. 21-40
3. 41-60
4. 61-80
5. 81-100
6. 101-120
7. 121-140
8. 141-160
9. 161-180
10. 181-200
11. 201-220
12. 221-240
13. 241-260
14. 261-280
15. 281-300
16. 301-320
17. 321-340
18. 341-360
19. 361-380
20. 381-400
21. 401-420
22. 421-440
23. 441-460
24. 461-480
25. 481-500
26. 501-520
27. 521-540
28. 541-560
29. 561-580
30. 581-600
31. 601-620
32. 621-640
33. 641-660
34. 661-680
35. 681-700
36. 701-720
37. 721-740
38. 741-760
39. 761-780
40. 781-800
41. 801-820
42. 821-840
43. 841-860
44. 861-880
45. 881-900
46. 901-920
47. 921-940
48. 941-960
49. 961-980
50. 981-1000
- *HEIGHT CLASS
1. 0-10 ft
2. 10.1-15 ft
3. 15.1-20 ft
4. 20.1-25 ft
5. 25.1-30 ft
6. 30.1-35 ft
7. 35.1-40 ft
8. 40.1-45 ft
9. 45.1-50 ft
10. 50.1-55 ft
11. 55.1-60 ft
12. 60.1-65 ft
13. 65.1-70 ft
14. 70.1-75 ft
15. 75.1-80 ft
16. 80.1-85 ft
17. 85.1-90 ft
18. 90.1-95 ft
19. 95.1-100 ft
20. 100.1-105 ft
21. 105.1-110 ft
22. 110.1-115 ft
23. 115.1-120 ft
24. 120.1-125 ft
25. 125.1-130 ft
26. 130.1-135 ft
27. 135.1-140 ft
28. 140.1-145 ft
29. 145.1-150 ft
30. 150.1-155 ft
31. 155.1-160 ft
32. 160.1-165 ft
33. 165.1-170 ft
34. 170.1-175 ft
35. 175.1-180 ft
36. 180.1-185 ft
37. 185.1-190 ft
38. 190.1-195 ft
39. 195.1-200 ft
40. 200.1-205 ft
41. 205.1-210 ft
42. 210.1-215 ft
43. 215.1-220 ft
44. 220.1-225 ft
45. 225.1-230 ft
46. 230.1-235 ft
47. 235.1-240 ft
48. 240.1-245 ft
49. 245.1-250 ft
50. 250.1-255 ft
51. 255.1-260 ft
52. 260.1-265 ft
53. 265.1-270 ft
54. 270.1-275 ft
55. 275.1-280 ft
56. 280.1-285 ft
57. 285.1-290 ft
58. 290.1-295 ft
59. 295.1-300 ft
60. 300.1-305 ft
61. 305.1-310 ft
62. 310.1-315 ft
63. 315.1-320 ft
64. 320.1-325 ft
65. 325.1-330 ft
66. 330.1-335 ft
67. 335.1-340 ft
68. 340.1-345 ft
69. 345.1-350 ft
70. 350.1-355 ft
71. 355.1-360 ft
72. 360.1-365 ft
73. 365.1-370 ft
74. 370.1-375 ft
75. 375.1-380 ft
76. 380.1-385 ft
77. 385.1-390 ft
78. 390.1-395 ft
79. 395.1-400 ft
80. 400.1-405 ft
81. 405.1-410 ft
82. 410.1-415 ft
83. 415.1-420 ft
84. 420.1-425 ft
85. 425.1-430 ft
86. 430.1-435 ft
87. 435.1-440 ft
88. 440.1-445 ft
89. 445.1-450 ft
90. 450.1-455 ft
91. 455.1-460 ft
92. 460.1-465 ft
93. 465.1-470 ft
94. 470.1-475 ft
95. 475.1-480 ft
96. 480.1-485 ft
97. 485.1-490 ft
98. 490.1-495 ft
99. 495.1-500 ft
100. 500.1-505 ft
101. 505.1-510 ft
102. 510.1-515 ft
103. 515.1-520 ft
104. 520.1-525 ft
105. 525.1-530 ft
106. 530.1-535 ft
107. 535.1-540 ft
108. 540.1-545 ft
109. 545.1-550 ft
110. 550.1-555 ft
111. 555.1-560 ft
112. 560.1-565 ft
113. 565.1-570 ft
114. 570.1-575 ft
115. 575.1-580 ft
116. 580.1-585 ft
117. 585.1-590 ft
118. 590.1-595 ft
119. 595.1-600 ft
120. 600.1-605 ft
121. 605.1-610 ft
122. 610.1-615 ft
123. 615.1-620 ft
124. 620.1-625 ft
125. 625.1-630 ft
126. 630.1-635 ft
127. 635.1-640 ft
128. 640.1-645 ft
129. 645.1-650 ft
130. 650.1-655 ft
131. 655.1-660 ft
132. 660.1-665 ft
133. 665.1-670 ft
134. 670.1-675 ft
135. 675.1-680 ft
136. 680.1-685 ft
137. 685.1-690 ft
138. 690.1-695 ft
139. 695.1-700 ft
140. 700.1-705 ft
141. 705.1-710 ft
142. 710.1-715 ft
143. 715.1-720 ft
144. 720.1-725 ft
145. 725.1-730 ft
146. 730.1-735 ft
147. 735.1-740 ft
148. 740.1-745 ft
149. 745.1-750 ft
150. 750.1-755 ft
151. 755.1-760 ft
152. 760.1-765 ft
153. 765.1-770 ft
154. 770.1-775 ft
155. 775.1-780 ft
156. 780.1-785 ft
157. 785.1-790 ft
158. 790.1-795 ft
159. 795.1-800 ft
160. 800.1-805 ft
161. 805.1-810 ft
162. 810.1-815 ft
163. 815.1-820 ft
164. 820.1-825 ft
165. 825.1-830 ft
166. 830.1-835 ft
167. 835.1-840 ft
168. 840.1-845 ft
169. 845.1-850 ft
170. 850.1-855 ft
171. 855.1-860 ft
172. 860.1-865 ft
173. 865.1-870 ft
174. 870.1-875 ft
175. 875.1-880 ft
176. 880.1-885 ft
177. 885.1-890 ft
178. 890.1-895 ft
179. 895.1-900 ft
180. 900.1-905 ft
181. 905.1-910 ft
182. 910.1-915 ft
183. 915.1-920 ft
184. 920.1-925 ft
185. 925.1-930 ft
186. 930.1-935 ft
187. 935.1-940 ft
188. 940.1-945 ft
189. 945.1-950 ft
190. 950.1-955 ft
191. 955.1-960 ft
192. 960.1-965 ft
193. 965.1-970 ft
194. 970.1-975 ft
195. 975.1-980 ft
196. 980.1-985 ft
197. 985.1-990 ft
198. 990.1-995 ft
199. 995.1-1000 ft
200. 1000.1-1005 ft
201. 1005.1-1010 ft
202. 1010.1-1015 ft
203. 1015.1-1020 ft
204. 1020.1-1025 ft
205. 1025.1-1030 ft
206. 1030.1-1035 ft
207. 1035.1-1040 ft
208. 1040.1-1045 ft
209. 1045.1-1050 ft
210. 1050.1-1055 ft
211. 1055.1-1060 ft
212. 1060.1-1065 ft
213. 1065.1-1070 ft
214. 1070.1-1075 ft
215. 1075.1-1080 ft
216. 1080.1-1085 ft
217. 1085.1-1090 ft
218. 1090.1-1095 ft
219. 1095.1-1100 ft
220. 1100.1-1105 ft
221. 1105.1-1110 ft
222. 1110.1-1115 ft
223. 1115.1-1120 ft
224. 1120.1-1125 ft
225. 1125.1-1130 ft
226. 1130.1-1135 ft
227. 1135.1-1140 ft
228. 1140.1-1145 ft
229. 1145.1-1150 ft
230. 1150.1-1155 ft
231. 1155.1-1160 ft
232. 1160.1-1165 ft
233. 1165.1-1170 ft
234. 1170.1-1175 ft
235. 1175.1-1180 ft
236. 1180.1-1185 ft
237. 1185.1-1190 ft
238. 1190.1-1195 ft
239. 1195.1-1200 ft
240. 1200.1-1205 ft
241. 1205.1-1210 ft
242. 1210.1-1215 ft
243. 1215.1-1220 ft
244. 1220.1-1225 ft
245. 1225.1-1230 ft
246. 1230.1-1235 ft
247. 1235.1-1240 ft
248. 1240.1-1245 ft
249. 1245.1-1250 ft
250. 1250.1-1255 ft
251. 1255.1-1260 ft
252. 1260.1-1265 ft
253. 1265.1-1270 ft
254. 1270.1-1275 ft
255. 1275.1-1280 ft
256. 1280.1-1285 ft
257. 1285.1-1290 ft
258. 1290.1-1295 ft
259. 1295.1-1300 ft
260. 1300.1-1305 ft
261. 1305.1-1310 ft
262. 1310.1-1315 ft
263. 1315.1-1320 ft
264. 1320.1-1325 ft
265. 1325.1-1330 ft
266. 1330.1-1335 ft
267. 1335.1-1340 ft
268. 1340.1-1345 ft
269. 1345.1-1350 ft
270. 1350.1-1355 ft
271. 1355.1-1360 ft
272. 1360.1-1365 ft
273. 1365.1-1370 ft
274. 1370.1-1375 ft
275. 1375.1-1380 ft
276. 1380.1-1385 ft
277. 1385.1-1390 ft
278. 1390.1-1395 ft
279. 1395.1-1400 ft
280. 1400.1-1405 ft
281. 1405.1-1410 ft
282. 1410.1-1415 ft
283. 1415.1-1420 ft
284. 1420.1-1425 ft
285. 1425.1-1430 ft
286. 1430.1-1435 ft
287. 1435.1-1440 ft
288. 1440.1-1445 ft
289. 1445.1-1450 ft
290. 1450.1-1455 ft
291. 1455.1-1460 ft
292. 1460.1-1465 ft
293. 1465.1-1470 ft
294. 1470.1-1475 ft
295. 1475.1-1480 ft
296. 1480.1-1485 ft
297. 1485.1-1490 ft
298. 1490.1-1495 ft
299. 1495.1-1500 ft
300. 1500.1-1505 ft
301. 1505.1-1510 ft
302. 1510.1-1515 ft
303. 1515.1-1520 ft
304. 1520.1-1525 ft
305. 1525.1-1530 ft
306. 1530.1-1535 ft
307. 1535.1-1540 ft
308. 1540.1-1545 ft
309. 1545.1-1550 ft
310. 1550.1-1555 ft
311. 1555.1-1560 ft
312. 1560.1-1565 ft
313. 1565.1-1570 ft
314. 1570.1-1575 ft
315. 1575.1-1580 ft
316. 1580.1-1585 ft
317. 1585.1-1590 ft
318. 1590.1-1595 ft
319. 1595.1-1600 ft
320. 1600.1-1605 ft
321. 1605.1-1610 ft
322. 1610.1-1615 ft
323. 1615.1-1620 ft
324. 1620.1-1625 ft
325. 1625.1-1630 ft
326. 1630.1-1635 ft
327. 1635.1-1640 ft
328. 1640.1-1645 ft
329. 1645.1-1650 ft
330. 1650.1-1655 ft
331. 1655.1-1660 ft
332. 1660.1-1665 ft
333. 1665.1-1670 ft
334. 1670.1-1675 ft
335. 1675.1-1680 ft
336. 1680.1-1685 ft
337. 1685.1-1690 ft
338. 1690.1-1695 ft
339. 1695.1-1700 ft
340. 1700.1-1705 ft
341. 1705.1-1710 ft
342. 1710.1-1715 ft
343. 1715.1-1720 ft
344. 1720.1-1725 ft
345. 1725.1-1730 ft
346. 1730.1-1735 ft
347. 1735.1-1740 ft
348. 1740.1-1745 ft
349. 1745.1-1750 ft
350. 1750.1-1755 ft
351. 1755.1-1760 ft
352. 1760.1-1765 ft
353. 1765.1-1770 ft
354. 1770.1-1775 ft
355. 1775.1-1780 ft
356. 1780.1-1785 ft
357. 1785.1-1790 ft
358. 1790.1-1795 ft
359. 1795.1-1800 ft
360. 1800.1-1805 ft
361. 1805.1-1810 ft
362. 1810.1-1815 ft
363. 1815.1-1820 ft
364. 1820.1-1825 ft
365. 1825.1-1830 ft
366. 1830.1-1835 ft
367. 1835.1-1840 ft
368. 1840.1-1845 ft
369. 1845.1-1850 ft
370. 1850.1-1855 ft
371. 1855.1-1860 ft
372. 1860.1-1865 ft
373. 1865.1-1870 ft
374. 1870.1-1875 ft
375. 1875.1-1880 ft
376. 1880.1-1885 ft
377. 1885.1-1890 ft
378. 1890.1-1895 ft
379. 1895.1-1900 ft
380. 1900.1-1905 ft
381. 1905.1-1910 ft
382. 1910.1-1915 ft
383. 1915.1-1920 ft
384. 1920.1-1925 ft
385. 1925.1-1930 ft
386. 1930.1-1935 ft
387. 1935.1-1940 ft
388. 1940.1-1945 ft
389. 1945.1-1950 ft
390. 1950.1-1955 ft
391. 1955.1-1960 ft
392. 1960.1-1965 ft
393. 1965.1-1970 ft
394. 1970.1-1975 ft
395. 1975.1-1980 ft
396. 1980.1-1985 ft
397. 1985.1-1990 ft
398. 1990.1-1995 ft
399. 1995.1-2000 ft
400. 2000.1-2005 ft
401. 2005.1-2010 ft
402. 2010.1-2015 ft
403. 2015.1-2020 ft
404. 2020.1-2025 ft
405. 2025.1-2030 ft
406. 2030.1-2035 ft
407. 2035.1-2040 ft
408. 2040.1-2045 ft
409. 2045.1-2050 ft
410. 2050.1-2055 ft
411. 2055.1-2060 ft
412. 2060.1-2065 ft
413. 2065.1-2070 ft
414. 2070.1-2075 ft
415. 2075.1-2080 ft
416. 2080.1-2085 ft
417. 2085.1-2090 ft
418. 2090.1-2095 ft
419. 2095.1-2100 ft
420. 2100.1-2105 ft
421. 2105.1-2110 ft
422. 2110.1-2115 ft
423. 2115.1-2120 ft
424. 2120.1-2125 ft
425. 2125.1-2130 ft
426. 2130.1-2135 ft
427. 2135.1-2140 ft
428. 2140.1-2145 ft
429. 2145.1-2150 ft
430. 2150.1-2155 ft
431. 2155.1-2160 ft
432. 2160.1-2165 ft
433. 2165.1-2170 ft
434. 2170.1-2175 ft
435. 2175.1-2180 ft
436. 2180.1-2185 ft
437. 2185.1-2190 ft
438. 2190.1-2195 ft
439. 2195.1-2200 ft
440. 2200.1-2205 ft
441. 2205.1-2210 ft
442. 2210.1-2215 ft
443. 2215.1-2220 ft
444. 2220.1-2225 ft
445. 2225.1-2230 ft
446. 2230.1-2235 ft
447. 2235.1-2240 ft
448. 2240.1-2245 ft
449. 2245.1-2250 ft
450. 2250.1-2255 ft
451. 2255.1-2260 ft
452. 2260.1-2265 ft
453. 2265.1-2270 ft
454. 2270.1-2275 ft
455. 2275.1-2280 ft
456. 2280.1-2285 ft
457. 2285.1-2290 ft
458. 2290.1-2295 ft
459. 2295.1-2300 ft
460. 2300.1-2305 ft
461. 2305.1-2310 ft
462. 2310.1-2315 ft
463. 2315.1-2320 ft
464. 2320.1-2325 ft
465. 2325.1-2330 ft
466. 2330.1-2335 ft
467. 2335.1-2340 ft
468. 2340.1-2345 ft
469. 2345.1-2350 ft
470. 2350.1-2355 ft
471. 2355.1-2360 ft
472. 2360.1-2365 ft
473. 2365.1-2370 ft
474. 2370.1-2375 ft
475. 2375.1-2380 ft
476. 2380.1-2385 ft
477. 2385.1-2390 ft
478. 2390.1-2395 ft
479. 2395.1-2400 ft
480. 2400.1-2405 ft
481. 2405.1-2410 ft
482. 2410.1-2415 ft
483. 2415.1-2420 ft
484. 2420.1-2425 ft
485. 2425.1-2430 ft
486. 2430.1-2435 ft
487. 2435.1-2440 ft
488. 2440.1-2445 ft
489. 2445.1-2450 ft
490. 2450.1-2455 ft
491. 2455.1-2460 ft
492. 2460.1-2465 ft
493. 2465.1-2470 ft
494. 2470.1-2475 ft
495. 2475.1-2480 ft
496. 2480.1-2485 ft
497. 2485.1-2490 ft
498. 2490.1-2495 ft
499. 2495.1-2500 ft
500. 2500.1-2505 ft
501. 2505.1-2510 ft
502. 2510.1-2515 ft
503. 2515.1-2520 ft
504. 2520.1-2525 ft
505. 2525.1-2530 ft
5



(FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

SERIES FORM

⑦ SPECIES LIST (circle one)

A. vascular and non-vascular plants (complete)

B. vascular only (complete)

C. partial list

① DATE	Y	M	D

② PROJECT ID:

Answer

Wissenschaft

3. SURVEYOR

④ SITE NO.

⑤ (Tributyl) (Mann, Fig. 1)

④MACROPLLOT NO.

[illegible]F81171 H88 BL1
EINVOCH

Ⓢ = mandatory for data systems



HERBICIDE / FERTILIZER FORM (EM-5A)

(FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

PROJECT IDENTIFICATION AND SAMPLING

① DATE Y M D ② PROJECT ID Name Number

③ SURVEYOR ④ SITE NO. ⑤ TREATMENT APPLIED
A. herbicide
B. fertilizer

⑥ KEY PLANT SPECIES Plants Tagged
Primary
Secondary

⑦ CONTROL AREA DESIGNATED

8. PROJECT COMMENTS

HERBICIDE OR FERTILIZER USED

⑨ CONTRACTOR Trade Name Composition

CHEMICAL COMPOSITION

⑩ ACTIVE INGREDIENT (a.i.)

11. CARRIER
A. water
B. oil
C. other (specify in Comments)

12. OTHER ADDITIVES ⑬ ACTUAL TREATMENT AREA ha

APPLICATION DETAILS (complete 14, 15 OR 16)

14. SPOT HERBICIDE APPLICATION
Method D. cut stump
A. foliar E. spot-gun
B. ground F. other (specify in Comments)
C. stem injection
Equipment Make/Model
Mix Concentration: Active Ingredient g/l or % Grid Size m
Additive g/l or %
Spray Mix Coverage: ml / spot Prescribed Application Rate: g of a.i. / spot

15. BROADCAST HERBICIDE APPLICATION
Method Equipment Make/Model
A. helicopter
B. fixed wing
C. tanker truck
D. other (specify in Comments)
Mix Concentration: Active Ingredient g/l or %
Additive g/l or %
Application Ground Speed km / h Flying Height m Swath Width m
Sprayer Output l / min Spray Mix Coverage l / ha Application Rate kg of a.i. / ha

16. FERTILIZER APPLICATION
Method Equipment Make/Model
A. helicopter
B. fixed wing
C. tanker truck
D. manual
E. other (specify in Comments)
Prescribed Application Rate kg / ha Actual Application Rate kg / ha

⑰ APPLICATION COVERAGE: Missed Areas
A. none missed
B. slightly missed
C. significant portions missed
Overlap Areas
A. no overlap
B. slight overlap
C. significant overlap

18. APPLICATION COMMENTS

ENVIRONMENTAL CONDITIONS DURING APPLICATION

19. TEMPERATURE

at start °C
at finish °C

20. HUMIDITY

at start %
at finish %

21. WIND Speed km / h

Direction °

22. PRECIPITATION DURING PREVIOUS WEEK

Type: A. none
B. rain
C. snow

Amount mm

Date(s) M D, M D

23. PRECIPITATION ON DATE OF APPLICATION

Type: A. none
B. rain
C. snow

Amount mm

24. PRECIPITATION DURING FOLLOWING WEEK

Type: A. none
B. rain
C. snow

Amount mm

Date(s) M D, M D

25. FOLIAGE MOISTURE CONDITION:

A. dry
B. slightly wet
C. wet

26. SOIL SURFACE CONDITION:

A. dry
B. wet
C. frozen
D. snow-covered



Province of British Columbia
Ministry of Forests
Ministry of Environment

SERIES FORM

HERBICIDE IMPACT ASSESSMENT (EM-5B)

(FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

(27) DATE Y M D (28) PROJECT ID Name Number

(29) SURVEYOR

(30) SITE NO.

(31) TIMING (Measurement No.)

MACRO-PLOT	(32) ENTRIES			(33) IMPACT CATEGORY										34. MACROPLOT COMMENTS		
	GENUS	SPECIES	C.S.	0		1		2		3		4			5	
				DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.		DOT COUNT	NO.
PLOT 1																
PLOT 2																
PLOT 3																
PLOT 4																
	35. SAMPLE STRIP SUMMARY														TOTAL STEMS NO.	36. SAMPLE STRIP COMMENTS
	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.		
SPECIES TOTALS AND PER- CENT																

FS 11798 HRI 8/11 - web
ENV 20652

☐ = mandatory for data system



Province of British Columbia
Ministry of Forests
Ministry of Environment

HERBICIDE RECOVERY ASSESSMENT (EM-5C) (FOR USE WITH THE "PROCEDURES FOR HABITAT MONITORING MANUAL")

SERIES FORM

(37) DATE Y M D (38) PROJECT ID Name Number
(39) SURVEYOR (40) SITE NO. (41) TIMING (Measurement No.)

MACRO-PLOT	(42) ENTRIES			(43) RECOVERY CATEGORY																44. MACROPLOT COMMENTS		
				STEM CATEGORY								BASAL CATEGORY										
	GENUS	SPECIES	C.S.	0		1		2		3		0		1		2		3				
				DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.	DOT COUNT	NO.			
PLOT 1																						
PLOT 2																						
PLOT 3																						
PLOT 4																						
SPECIES TOTALS AND PER-CENT				45. SAMPLE STRIP SUMMARY																TOTAL STEM	TOTAL BASAL	
				%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	NO.	NO.	
46. SAMPLE STRIP COMMENTS																						

FS 1179C HR 9/11 - wsl
ENV 2065C

☐ = mandatory for data system

APPENDIX 6. Fuel Assessment form and tally rules (from Trowbridge *et al.* 1986)

Fuel Assessment Form

This form is completed according to procedures described by Trowbridge *et al.* (1986), with the following modifications:

- Unique no:** PROBE installation number
- Plot:** T (Treatment).
- Line:** Lines A, B, and C correspond to gridlines 1, 2, 3, respectively. Another copy of the form is required for gridlines 4, 5, and 6.
- Depths:** Pre-burn depth of mosses and litter (cm), depth of burn (cm), and post-burn duff depth (cm) are measured at 24 depth-of-burn pins located along each line (i.e., 6 gridpoints $\times$ 4 pins per gridpoint). Assume that 5 m, 10 m, 15 m, etc. correspond with gridpoint 1, 2, 3, etc., respectively. Enter depths for North/South pins under “in” and for East/West pins under “out.”
- Plot diagram:** Ignore this diagram.
- Slope:** Measured at the beginning of each gridline.

Tally Rules (summarized from Trowbridge *et al.* 1986).

Tally rules that apply to all size classes are described in detail by Trowbridge *et al.* (1986). In brief, the eight rules are:

1. All pieces must be *downed, dead* woody material.
2. Where the gridline intersects the end of a piece, count the piece only if the line crosses its central axis.
3. Where the gridline *exactly* intersects the central axis of the end of a piece, tally every other such piece.
4. Where the central axis of a piece coincides *exactly* with the gridline, do **not** tally the piece.
5. Where the gridline intersects a curved or angular piece more than once, tally each intersection.
6. Where a piece has been shattered or is rotten, estimate its original diameter.
7. Tally a stump or root only if it is uprooted and not encased in soil.
8. Tally only those pieces whose central axis is above the F and H layers at the point of gridline intersection.

Unique No. Plot Recorded by Pre-fire Post-fire
 FUEL ASSESSMENTS Date Pre-fire Post-fire
 (Sections 4.4 and 6.1 in Handbook)

Line	Species and diameters of slash pieces larger than 7 cm in diameter (Indicate location of 5, 10, 15, 20 and 25 m marks)																			
A	Pre	sp																		
		dia																		
	Post	dia																		
B	Pre	sp																		
		dia																		
	Post	dia																		
C	Pre	sp																		
		dia																		
	Post	dia																		

Intersections by pieces smaller than 7 cm					
class	0-0.5	0.6-1.0	1.1-3.0	3.1-5.0	5.1-7.0
length	5 m	10 m	15 m	20 m	25 m
A	Pre				
	Post				
B	Pre				
	Post				
C	Pre				
	Post				

Species composition of slash < 7 cm					
Species					
Line A					
Line B					
Line C					


Material soil exposure			
pre-burn	A	B	C
post-burn			

Notes

Preburn depth of mosses and litter (cm)										
Line	5 m		10 m		15 m		20 m		25 m	
	in	out	in	out	in	out	in	out	in	out
A										
B										
C										

Depth of burn (cm)									
A									
B									
C									

Post-burn duff depth (cm)										
Line	5 m		10 m		15 m		20 m		25 m	
	in	out	in	out	in	out	in	out	in	out
A										
B										
C										



APPENDIX 7. MSP Soil Disturbance Survey forms and Soil Disturbance Categories (from Curran and Thompson 1991, unpubl.)

1-MSP: Gridpoint-Intercept Survey

This form is completed according to procedures described by Curran and Thompson (1991, unpubl.), with the following modifications:

- Treatment Unit (TU):** T (Treatment) or C (Control).
- TU size:** Size of T and C plots (e.g., 0.80 ha)
- Grid spacing:** Ten metres between gridpoints, 1m from gridpoint to point observation.
- Gridpoint:** Thirty-six gridpoints are evaluated in each T and C.
- Slope:** Measure slope at the beginning of each gridline. Enter the value at the appropriate gridpoint (e.g., 1, 7, 13, etc.).
- Point observations:** Four point observations of soil disturbance are made at each gridpoint. The points are located 1m in each cardinal direction from the gridpoint. Provide comments, if necessary. Refer to "Soil Disturbance Categories for MSP Areas" in this Appendix.

Soil Disturbance Categories for MSP areas (Curran and Thompson 1991, unpubl.)

Object	Code	Description		
<i>Most severe disturbance types</i>				
Traversed	T	Traverse method disturbance: disturbance that would be measured by the traverse method (i.e., all types of haul roads regardless of whether they are being traversed or not, landings, fire guards, and backspur trails).		
Skidroad	R	Skidroad cutbank or running surface.		
	B	Skidroad berm/sidecast (counted to points where it is continuously 5cm deep in mineral soil; 15cm in woody debris).		
<hr/>				
<i>Heavy Machine Traffic</i>		Soil type	Depth criteria	Disturbance type count
<hr/>				
			(into mineral soil)	
	1	Cohesive	5–10cm	High impact machine traffic, Wheel/track impressions and rutting
	2	Cohesive	>10cm	High impact machine traffic, Wheel/track impressions and rutting
	3	Non-cohesive sandy ^a	>15cm	High impact machine traffic displacement
<hr/>				
			(into organic layer)	
	4	Organic	>20cm	High impact machine traffic rutting

APPENDIX 7. (continued)

Object	Code	Description
MSP^b	W	Wide Gouge: ≥5cm deep gouge into mineral soil (over 80% of its area) that is >2 m in diameter in any orientation from the point.
	G	Deep Gouge: ≥30cm deep gouge into the mineral soil at the point, regardless of length or width.
	S ^c	Wide Scalp: a 2.0m diameter patch (or larger), where the forest floor has been removed from over 80% of the area in any orientation from the point.
	C ^c	Continuous Gouge: ≥5cm deep gouge (at any place in the gouge's cross-section where the point lies) into the mineral soil that is >1.5m along one of its axes , length or width in any orientation from the point.
Other	O ^d	Other types of disturbance. Minor scalps and gouges which do not fit into any of the above categories. Light skidding disturbance should be classified as MSP disturbance or "L."
<hr/>		
Light Machine Traffic	Soil type	Depth criteria Disturbance type count
	L ^d All	< above criteria Light machine traffic disturbance not meeting any other criteria in this table (i.e., wheel/track impressions with intact forest floor and depth less than criteria for "1" or "2").
<hr/>		
Less Severe and Possibly Beneficial Disturbance Types		
	P ^e	Piles of slash debris that are not associated with landings (which are Traversed disturbance) and are 3.0×3.0 m or larger in size, greater than 50cm deep, and less than 33% mineral soil. For example, piles and windrows form rough-bunching or windrowing.
	D ^e	Deposits: deposits and mixtures of forest floor and mineral soil that are not the sidecast of bladed structures such as haul roads, landings, and skidroads. Piles that are greater than 33% mineral soil are counted as Deposits.
Undisturbed	"-"	A forest floor with no mineral soil exposed or any impression from machine travel. Also includes surfaces beneath stumps or large coarse fragments, or natural bedrock surfaces. If an assessment beneath slash or large logs is possible, then it is made. Otherwise, these surfaces are recorded as undisturbed.

^a "Non-cohesive sandy soils" have a very weak moist cast which is non-sticky and will not form a moist worm ≤4 mm (i.e., S, LS, and some SL textures).

^b Classify disturbance based on the soil disturbance at the point. For MSP disturbance types, when the point lands on a scalp or gouge, assess the overall soil condition to see if it meets the spatial categories "W," "S," or "C."

^c Disturbance categories counted as detrimental on more sensitive sites.

^d The "Other" and "L" categories for MSP are different from "Other" and "L" categories in Field Guide Insert No. 5, "Measuring Soil Disturbance Following Timber Harvest."

^e Disturbance categories not counted in MSP guidelines.

- Used for undisturbed so that these can appear as a horizontal line on the survey form.

Note: When a point can be classified into more than one category, place it in the most severe. (This table is ranked in order of severity from top to bottom.)

3A-MSP: Calculation Card Gridpoint-Intercept

This form is completed according to procedures described by Curran and Thompson (1991, unpubl.), with the following modifications:

Treatment unit: T (Treatment) or C (Control).

Transect: Corresponds with gridline, of which there are six in a Treatment unit.

Category tallies: For each gridline (transect), tally the number of point observations in each category. There are 24 point observations (6 gridpoints $\times$ 4 observations/gridpoint) in each gridline (6 gridpoints/gridline).

Calculation procedure:

1. Enter the number of point observations (not gridpoints).
2. Calculate the mean number of points for each category ($n=6$).
3. Calculate the standard deviation ($n=6$).
4. Calculate the Mean% as: $(\text{Mean}/24) \times 100$.

Surveyor	Date	BCGS opening	
License/license #	CP	Block	Treatment unit

Category Tallies																	
Transect	R	B	1	2	3	4	L	O	T	W	G	C	S	P	D	-	
1																	
2																	
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	
11																	
12																	
13																	
14																	
15																	
16																	
17																	
18																	
19																	
20																	
21																	
22																	
23																	
24																	
25																	
26																	
27																	
28																	
29																	
30																	

MEAN																
STD																
MEAN %																

Calculation Procedure

1. Enter the number of gridpoints.
2. Enter the MEAN number of points calculated for the tally sheet.
3. Enter the STD calculated on the tally sheet.
4. Calculate MEAN %.

APPENDIX 8. Analysis of Variance table

ANOVA for a randomized complete block design with three replications

Source	df		SS	MS	F
Total	$pr-1$	5	SSTot		
Block	$r-1$	2	SSR	MSR	FR
Treatment	$p-1$	1	SSTr	MSTr	FTr
Error	$(r-1)(p-1)$	2	SSE	MSE	FE

$r = \text{block} = 3$

$p = \text{treatment} = 2$

REFERENCES

- Alpha Software Corporation. 1991. Alpha Four, Version 2. Alpha Software Corp., Burlington, Mass.
- Angove, K. and B. Bancroft. 1983. A guide to some common plants of the southern interior of British Columbia. B.C. Min. For., Victoria, B.C. Land Manage. Handb. No. 7.
- Braumandl, T.F. and M.P. Curran. (editors). 1992. A field guide for site identification and interpretation for the Nelson Forest Region. B.C. Min. For., Victoria, B.C. Land Manage. Handb. No. 20.
- British Columbia Ministry of Forests. 1989. Silviculture manual. Vol. 1. Silv. Branch, Victoria, B.C.
- . 1990. First approximation of correlated guidelines for free growing stocking standards for the ecosystems of British Columbia. Silv. Branch, Victoria, B.C.
- Canada Soil Survey Committee, Subcommittee on Soil Classification. 1978. The Canadian system of soil classification. Ottawa, Ont. Can. Dep. Agric. Publ. 1646.
- Comeau, P. and T. Braumandl. 1991. Light attenuation and competition by overtopping vegetation in the Interior Cedar Hemlock zone — Project No. 3.48. B.C. Min. For. & For. Can., Victoria, B.C. FRDA Res. Memo No. 159.
- Dierschke, H. 1972. On the recording and presentation of phenological phenomena in plant communities. *In* 1970 International Symp. for Vegetation Science. Dr. W. Junk (editor). N.V. The Hague, Netherlands.
- Goldberg, D.E. and P.A. Werner. 1983. Equivalence of competitors in plant communities: a null hypothesis and field experimental approach. *Am. J. Bot.* 70:1098–1104.
- Habitat Monitoring Committee. 1990. Procedures for environmental monitoring in range and wildlife habitat management. Draft edition version 4.1. B.C. Min. Environ. and B.C. Min. For., Victoria, B.C. 196p.
- Herring, L. and J. Pollack. 1985. Experimental design protocol for forest vegetation management research: level A trials. First approx. B.C. Min. For., Victoria, B.C. Res. Rep. RR84012-HQ.
- Hurlbert, S.H. 1984. Pseudo-replication and the design of ecological field experiments. *Ecolog. Monogr.*, 54:187–211.
- Kimmins, J.P. and P.G. Comeau. 1990. Forest vegetation management research strategy for the southern interior, British Columbia. B.C. Min. For. & For. Can., Victoria, B.C. FRDA Rep. No. 132.
- Klinka, K., R.N. Green, R.L. Trowbridge, and L.E. Lowe. 1981. Taxonomic classification of humus forms in ecosystems of British Columbia. B.C. Min. For., Victoria, B.C. Land Manage. Rep. No. 8.
- Lloyd, D., K. Angove, G. Hope, and C. Thompson. 1990. A guide to site identification and interpretation for the Kamloops Forest Region. B.C. Min. For., Victoria, B.C. Land Manage. Handb. No. 23.
- Luttmerding, H.A., D.A. Demarchi, C.A. Lea, D.V. Meidinger, and T. Vold. (editors). 1990. Describing ecosystems in the field. 2nd ed. B.C. Min. Environ. and B.C. Min. For., Victoria, B.C. B.C. Min. Environ. Manual 11.
- SAS Institute Inc. 1985. SAS procedures guide for personal computers, Version 6. SAS Institute Inc., Cary, N.C.
- Trowbridge, R., B. Hawkes, A. Macadam, and J. Parminter. 1986. Field handbook for prescribed fire assessments in British Columbia: logging slash fuels. B.C. Min. For., Victoria, B.C. Land Manage. Handb. No. 11.
- Wagner, R.G. and S.R. Radosevich. 1991. Neighborhood predictors of interspecific competition in young Douglas-fir plantations. *Can. J. For. Res.* 21:821–828.
- Weiner, J. 1984. Neighborhood interference amongst *Pinus rigida* individuals. *J. Ecol.* 72:183–195.