


Importance of small streams



Lisa Nordin, MSc., RPBio.







FIGURE 13.1 A stream reach showing many of the elements and processes that link streams and riparian areas (E. Leinberger, University of British Columbia).



Site Level / Watershed Level



- Highly productive at site level
- Transport water, nutrients and organic matter including invertebrates to downstream reaches
- Lower gradient reaches important to juvenile fish
- Higher gradient reaches support many unique species that do not co-exist well with fish



Travel and forage corridors



BC Adventure Network



Pinterest- Whistler and BC Wildlife



Roger Patterson 1967




Okanagan Similkameen Stewardship Society



Fish BC

Lastly.....



- Most Abundant
- Most sensitive
- Least buffered
- Consequently, most at risk

References

- Burns, J.E. 1970. The importance of streamside vegetation to trout and salmon in British Columbia. Vancouver Island Region, Fish and Wildlife Branch, Department of Recreation and Conservation. Nanaimo, B.C. 14 pp.
- Rosenfeld, J.S., MacDonald, S., Foster, D., Amrhein, S., Bales, B., Williams, T., Rance, F., and T. Livingstone. 2002. Importance of small streams as rearing habitat for coastal cutthroat trout. *North American Journal of Fisheries Management* 22:177-187.
- Saunders, D.L., Meeuwig, J.J. and A.C.J.Vincent. 2002. Freshwater Protected Areas; Strategies for Conservation. *Conservation Biology* 16 (1): 30-41
- Wallace, J.B., Eggert, S.L. Meyer, J.L. and J.R. Webster. 1997. Multiple Trophic Levels of a Forest Stream Linked to Terrestrial Litter Inputs. *Science*. 277 (5322); 102-104
- Wipfli & Gregovich, 2002. Export of invertebrates and detritus from fishless headwater streams in southeastern Alaska: implications for downstream salmonid production. *Freshwater Biology*. 47:957-969