

Uneven-aged Stocking Standards* -- Single-tree selection only

Col. 1	Col. 2	Col. 3	Col. 4	Col. 5	Col. 1	Col. 2	Col. 3	Col. 4	Col. 5
Target from Table A standards	Layer**	Stocking***			Target from Table A standards	Layer**	Stocking***		
		Target pa	MIN pa	MIN p			Target pa	MIN pa	MIN p
(stems/ha)		(well-spaced/ha)			(stems/ha)		(well-spaced/ha)		
1200	1	600	300	250	800	1	300	150	150
	2	800	400	300		2	400	200	200
	3	1000	500	400		3	600	300	300
	4	1200	700	600		4	800	400	400
1000	1	400	200	200	600	1	300	150	150
	2	600	300	250		2	400	200	200
	3	800	400	300		3	500	300	300
	4	1000	500	400		4	600	400	400
900	1	400	200	200	400	1	200	100	100
	2	500	300	250		2	300	125	125
	3	700	400	300		3	300	150	150
	4	900	500	400		4	400	200	200

MIN - minimum

* Maximum regeneration delay is seven years. For a seven-year regeneration delay, the early free growing is 12 years and the late free growing is 15 years. Regeneration delay can be met immediately following harvest if the residual stand has no significant damage or pest problems and meets minimum stocking standards. If regeneration is achieved immediately following harvest, earliest free growing date is 12 months after completion of harvest and the latest date is 24 months after completion of harvest.

**Stand Layer Definition

Layer 1	Mature	trees $\geq$ 12.5 cm dbh
Layer 2	Pole	trees 7.5 cm to 12.4 cm dbh
Layer 3	Sapling	trees $\geq$ 1.3 m height to 7.4 cm dbh
Layer 4	Regeneration	trees < 1.3 m height

*** pa - preferred and acceptable species p - preferred species

Preferred and acceptable species and "Target from Table A standards" are as specified in Table A by biogeoclimatic ecosystem classification (BEC) site series.