


Ecology and Earth Sciences Section

An overview of recent accomplishments


Evelyn Hamilton, MSc, RPBio
Manager, Ecology and Earth Science Section,
Research Branch
BC Ministry of Forests and Range

Outline

- ▶ Overview of Research Branch
- ▶ Ecology and Earth Sciences Section
- ▶ EES Program Highlights
 - Ecology
 - Watershed Science
 - Soils
 - Climatology and Carbon
 - Range Ecology
 - Landscape Ecology
 - Wildlife Habitat
- ▶ Summary




Overview of Research Branch




Ecology and Earth Science Section

- ▶ Organization chart
- ▶ Main activities
- ▶ Key drivers
- ▶ Key clients and collaborators




Description of EES Section

- ▶ Organization chart
- ▶ Locations – Victoria, Smithers, Kamloops
- ▶ Allocation of effort
- ▶ Strengths



EVELYN HAMILTON
Manager
(2086) ML6



Main activities

- ▶ Provincial research program coordination.
- ▶ Research on long and short term projects
- ▶ Providing expert advice to decision makers
- ▶ Developing assessment procedures
- ▶ Analysis of implications of management actions
- ▶ Using existing info to address new issues (e.g., implications of climate change)
- ▶ Extension and consulting


Approx. 40% of time spent on research

Approx. 60% of time supporting clients through consulting and other activities



Key drivers

- ▶ MFR Act (S. 4) – Manage, protect, conserve and plan the use of forest and range resources, encourage maximum productivity and competitive forest and range industries
- ▶ Economics/industry competitiveness – green energy; carbon economy; supporting communities
- ▶ Balancing competition for forest resources
- ▶ Climate change and carbon economy
- ▶ First Nations partners in the new economy
- ▶ Maintaining public trust in resource management




Key clients and collaborators

Key clients

- ▶ Forest Stewardship Division
- ▶ Operations Division
- ▶ Tenure and Revenue Division
- ▶ BCTS
- ▶ Other agencies – MOE, MEMFR, DFO


Key collaborators

- ▶ Regional MFR scientists, UBC, UNBC, SFU, TRU, CFS, FP Innovations, BVC, FORREX



Ecology Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions





*Shirley Mah
A/ Team Leader, Ecology Program and
Interpretations Ecologist
Victoria*

Biogeoclimatic Ecosystem Classification

& Climate Change




*Will Mackenzie
Research Ecologist– BEC system
Smithers*





*Elizabeth Campbell
Research Ecologist –
Ecosystem Processes
Victoria*



*Ben Heemskerk
Assistant Ecologist
Smithers*



TECHNICAL REPORT 051

Ecosystem and Tree Attributes Affecting the Presence of Functional Wildlife Tree Types

2009



Ministry of Forests and Range
Forest Science Program

051


TECHNICAL REPORT 046

A Framework for Documenting the Effects of the Mountain Pine Beetle Outbreak in Sub-boreal Forests of Northern British Columbia (E.P. 1369): Establishment Report

28



Ministry of Forests and Range
Forest Science Program



*John Parminter
Fire Ecologist
(Emeritus scientist)
North Road Lab*

Watershed Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions





*Peter Tschaplinski
Manager Watershed Research Program, Victoria office*

*Dan Hogan
Research Scientist ,Geomorphology , UBC office*



*Robin Pike
Watershed Ecosystem Specialist
Victoria office*

Peer-reviewed Synthesis Article

Hydrologic Models for Forest Management Applications: Part 2: Incorporating the Effects of Climate Change

Jos Beckers, Robin Pike, Arifia Z. Warner, Todd Hedding, Brian Smardon, and Axel Anderson

Introduction

In Alberta and British Columbia, several detailed studies of climate trends, future climate predictions, and potential effects on hydrology have been conducted (e.g., Rodenhuis et al. 2007; Pike et al. 2008a, 2008b; Sauchyn and Kulshreshtha 2008; Walker and Sydeman 2009). These studies indicate that a changing climate will alter watershed many aspects of short- and long-term watershed management. From an operational perspective, watershed-scale hydrologic models could be

specific qualities required in a hydrologic model for climate change applications in a forest management context, reviews the suitability of several currently available models, and discusses suggested improvements for climate change and forest management applications.

Background

Our climate change review focuses on the nine short-listed models in Part 1 that were identified as suitable for address.

Several detailed studies of climate trends, future climate predictions, and potential effects

1. increased atmospheric evaporative demand;
2. altered vegetation composition affecting evaporation and precipitation interception;
3. decreased snow accumulation and earlier melt;
4. accelerated melting of permafrost, lake ice, and river ice;
5. glacier mass balance adjustments;
6. increased stream and lake temperatures;
7. increased frequency or magnitude of disturbances; and
8. altered streamflow.

Associated with each of these are specific processes (e.g., evapotranspiration), watershed outputs (e.g., timing and magnitude of peak/low flows, stream temperature), and other factors (e.g., growing conditions for trees, wildfire risk) that could be affected by anticipated shifts in meteorological and hydrological conditions. These processes, associated model inputs, watershed outputs, and other factors must therefore be present in hydrologic models to enable the investigation of climate change

86

Extension Note

MARCH 2008

Lisa Nordin, Hydrology
Research Associate

Ministry Contacts:
John Rex and David Maloney
B.C. Ministry of Forests
and Range
Northern Interior Region
5th Floor, 1011 4th Avenue
Prince George, B.C. V2L 3H9

87

Extension Note

MARCH 2008

R.G. Pike, D.L. Spittlehouse,
K.E. Bennett, V.N. Egginton,
P.J. Tschaplinski,
T.G. Morduck, and
A.T. Werner

88

Extension Note

Mar 2008

Rita Winkler
Southern Interior Region,
B.C. Ministry of Forests and
Range, Kamloops, BC

John Rex
Northern Interior Region,
B.C. Ministry of Forests and
Range, Prince George, BC

Pat Tei
Southern Interior Region,
B.C. Ministry of Forests and
Range, Kamloops, BC

David Maloney
Northern Interior Region,
B.C. Ministry of Forests and
Range, Prince George, BC

Todd Redding
FORREC Forest Research
Extension Partnership,
Nanaimo, BC

90

Extension Note

MARCH 2009

John Rex
Northern Interior Region
B.C. Ministry of Forests and Range
Prince George, BC

Philip Krauskopf
Northern Interior Region
B.C. Ministry of Forests and Range
Prince George, BC

David Maloney
Southern Interior Region
B.C. Ministry of Forests and Range
Kamloops, BC

Peter Tschaplinski
Research Branch
B.C. Ministry of Forests and Range
Victoria, BC

Mountain Pine Beetle and Salvage Harvesting: Small Stream and Riparian Zone Response in the Sub-Boreal Spruce Zone

Introduction

The mountain pine beetle (MPB) infestation and its related salvage harvesting activities adjacent to streams have the potential to influence small stream (< 2 m bankfull width) and riparian zone function by altering and removing riparian vegetation and causing other disturbances. These alterations are a significant management issue because small streams are the most predominant channel type on the landscape, comprising upwards of 80% of a watershed's total channel length (Shreve 1969). Accordingly, understanding the influence of these changes on small streams and their riparian zones is important to forest management.

The MPB infestation of riparian pine stands and/or subsequent riparian harvesting can alter riparian structure by changing microclimate


influence of MPB and salvage harvesting on riparian zones and small streams. The initial studies described here were implemented in the Vanderhoof Forest District because it has been heavily affected by MPB and has correspondingly seen increased levels of salvage harvesting to recover some economic value from the forest. These studies aim to provide short-term answers to questions about the influence of the MPB and salvage harvesting on small streams, namely:

- What is the riparian stand structure of small streams in MPB-affected areas of the SBS within the Vanderhoof Forest District?
- How do the MPB infestation (grey attack stage) and salvage harvesting affect riparian zones and small streams, including shade?

Study Design...

Creek

Carnation Creek – The World's Longest Continuous Study of Fish-Forestry Interactions



How do Pacific coastal watersheds function? How do forestry practices affect these functions? Scientists have analyzed these complex questions for 35 years at Carnation Creek, a small watershed on Vancouver Island. The Carnation Creek research project is the longest-running study of the effects of forestry practices on watershed processes and is responsible for much of our present knowledge of fish-forestry interactions. In the late 1960s, conflicts over approaches to fisheries and forest resource management highlighted our lack of understanding of Pacific coastal watersheds. To fill this gap, Forest Sciences of Canada and the B.C. Ministry of Forests and Range, in partnership with MacMillan Bloedel, initiated the Carnation Creek project. The project's partners started to involve many scientists and researchers from the B.C. Ministry of Forests and Range, as well as other agencies, to conduct research at Carnation Creek. Research results from the project have provided the scientific basis for improved forest practices in British Columbia and have outlined some major findings.


Forest Sciences of Canada
B.C. Ministry of Forests and Range

Soils Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions



Shannon Berch
Team Leader, Soils and
Research Scientist
North Road Lab, Victoria



Commercially Important Wild Mushrooms
and Fungi of British Columbia
What the Buyers Are Buying

91

Extension Note

84

Extension Note

Fertilization on Fine Roots,
Soil Mesofauna in Young Lodgepole
Pines in Central British Columbia

Best Practices for Soil
in Mountain Pine Bee
Infestations

Prince George,
Prince Rupert
and Cariboo
Forest Regions

March 2009
Page 84/84-03



LTSPS RESEARCH NOTE

**Soil Fauna in the Sub-Boreal Spruce (SBS)
Installations of the Long-Term Soil Productivity
(LTSP) Study of Central British Columbia: One-
Year Results for Soil Mesofauna and Macrofauna**

Jeff Battipelli and Shannon Berch

Introduction

The diversity of animal species in forest soils rivals that of a coral reef, with hundreds of species represented by thousands or millions of individuals occupying a single square meter of soil. Soil fauna come in a variety of shapes and sizes and play a variety of roles in soil development and the maintenance of soil fertility. Soil mesofauna are defined as organisms bigger than 2 mm in body width and are the most conspicuous animals within the soil ecosystem. Included within this group are ants, beetles, spiders, centipedes, millipedes, and earthworms. Although generally less numerous than soil mesofauna, macrofauna can represent a significant proportion of the animal biomass in the soil and can play an important role in soil ecosystem function. For example, millipedes and earthworms break up organic matter, increasing its surface area and thereby enhancing

microbial activity and nutrient cycling. Soil macrofauna mix and redistribute mineral and organic material and microorganisms within the soil profile. Soil mesofauna range from 0.1 to 2 mm wide and include mites, springtails, Diplura, Symphyla, and Tardigrada, as well as smaller forms of spiders, pseudoscorpions, and insect larvae. Via grazing, they control bacterial and fungal biomass, thus liberating mineral nutrients and stimulating further fungal growth. Furthermore, soil mesofauna transport microbe propagules and spores into new substrates and contribute to the development of soil structure and humus formation through the deposition of fecal pellets.

Timber harvesting and site preparation can modify physical and chemical properties of the soil, consequently affecting soil porosity, composition, and organic matter content. Forest floor and mineral soil temperatures and soil moisture. Organic matter supplies nutrients in soil, influences soil structure, and provides habitats in soil, influences soil structure, and provides habitats in soil. The alteration of soil porosity through compaction or displacement can directly influence biological activity, reducing the diversity and density of soil fauna and altering the structure and function of the soil fauna community.

In 1990, in conjunction with a similar research program in the United States, the B.C. Ministry of Forests established the Long-Term Soil Productivity (LTSP) study to determine the effects of three levels of organic matter removal and soil compaction on soil

soil following intensive fertilization of lodgepole pine. In contrast, the density of mites in the soils under intensive repeated fertilization. It is not possible to ascribe the different responses of soil fine roots, ectomycorrhizae, and soil mesofauna found at these two sites to a specific cause, nor is it possible to state with certainty that the previously reported differences in aboveground tree growth responses of pine and spruce at these two sites (small and large, respectively) are directly linked with the belowground changes. Root and soil biota characteristics at other lodgepole pine and spruce "maximum productivity" sites are currently being examined to further understand the possible relationship between tree species and above- and belowground changes following repeated fertilization and the functional relevance of these effects.

Introduction

Nutrient deficiencies are widespread throughout the British Columbia interior, and the growth benefits of fertilizing with nitrogen (N) and other nutrients have been well documented for several conifer species. Although a single fertilizer application typically produces only a temporary increase

and how these soil seasonal windows of opportunity. Accordingly, this additional guidance on timing and operation to conserve soil fertility and summarize the factors affecting the harvest of live and provide species to avoid or minimize the disturbance to affected stands.

Background

The LTSP study in British Columbia first the landscape with pine loss dramatic change when (et al. 2008), or on foliar of dead tree ground the Dead and less than 1 processes and with have also intercept

FIGURE 1. Mean density of mesofauna (mostly mites and springtails) in the central forest control of the SBS LTSP sites.



Ministry of Forests and Range
Forest Science Program

Compatible Management of Timber and Pine Mushrooms

Ministry of Forests and
Range, Research Branch

The Centre for Non-
Timber Resources
Royal Roads University

March 2009



Shannon M. Berch and J. Marty Kranabetter
BC Ministry of Forests and Range
Research Branch, Victoria



*Chuck Bulmer
Research Soil Scientist
Kalamalka Lab
Vernon*



Climatology and Carbon Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions



*Dave Spittlehouse
Team Leader, Forest Climatology
and Carbon Research
Victoria office*

PCIC Regional Analysis Tool (BETA): [Help](#)


Data Options

Experiment: CLIMATEBC HIST-run1 (HIST) Variable: Mean Temperature
Timeslice: 1998 Time of Year: Annual

Display Options

Window: BC Albers Region: --Entire Window--
Update

Plot Options



Map Range

☐ Auto
☒ Fixed Range


-5 to 15


Options

☐ Show Grid

Map Size


800x600





*Caren Dymond
Forest carbon and
climate change researcher
Victoria office and PFC*



[illegible][illegible][illegible]


Landscape Ecology Program


- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions






*Adrian Walton
Team Leader,
Landscape Ecology
Victoria office*










Huapeng Chen
Landscape Modelling
Biologist
Victoria office





Range Ecology Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions





*Reg Newman, Team Leader, Range Ecology Research
Brian Wallace, Range Ecologist – Soils
Kamloops office*


Linking Range Health Assessment Methodology with Science

Reg Newman¹, Rick Tucker², Francis Njenga², Graeme Hope³, Maja Krzic⁴ and Sheryl Wurtz¹

The plant community approach to assessing range conditions came into question in the 1980's. Range health assessment has responded to this challenge by the traditional plant-based approach to determine the state of rangelands, and include assessment of ecological functions in addition to plant community information.

Range health has been defined as:

- The ability of rangeland to perform certain key functions... where all parts make up the whole, are present and working together (Jodanis et al. 2005)
- The degree to which the integrity of the soil and the ecological processes of rangeland ecosystems are sustained (Rusby 1986).



Grassland cover in rangelands contains some of BC's most threatened habitats. Thanks to improved scientific studies and better management, rangelands are becoming more resilient.

Relationships among Soil/Vegetation Properties and Rangeland Health Indicators

Sarah Lamagna¹, Maja Krzic¹, and Reg Newman²

¹University of British Columbia, Faculty of Land & Food Systems, Vancouver, BC; ²BC Ministry of Forests & Range, Kamloops, BC

Rationale and Study Objective

- * Rangeland health assessment methods rely on qualitative, visual estimates and concerns have been raised regarding their reliability. Studies on development and application of criteria and indicators for forests and rangelands are often lacking, or have been done on a limited number of sites with relatively narrow ranges of climate and soil type.


- * Development of unambiguous indicators of rangeland health should start with identification of vegetation and soil properties that integrate important ecological processes, followed by examination of linkages among these selected properties and rangeland health assessment indicators.



Regression Analysis

- * Soil bulk density had a significant ($P < 0.001$), positive relationship with percentage cover of Junegrass. Bulk density increased with increasing Junegrass cover until 15% cover was reached (Fig. 3).

- * Junegrass occurs at earlier seral stages and increases with grazing in the rough fescue grasslands of BC.



Understory Succession following Ecosystem Restoration Treatments in Ingrown Dry Forests

Reg Newman¹, Hillary Page², John Parminter³, and Brian Wallace¹

Dry forests of ponderosa pine (*Pinus ponderosa*) and interior Douglas-fir (*Pseudotsuga menziesii* var. *glauca*) occurring in the valleys at low- to mid-elevations within the southern Interior of British Columbia.


Fire suppression, overgrazing, and selective logging in these forests are believed to have caused forest encroachment on grasslands and ingrowth within open forests. Encroachment has established in previously treeless openings. Ingrowth is associated with recruitment, primarily by shade-tolerant species, such as interior Douglas-fir, within low-density, open forests.




Open forest - pine stand near Creston, BC.



Tree forest - pine stand near Creston, BC.



Premier Lake - 1954



Premier Lake - 1994

The photo pair above illustrates forest cover changes during a 40 year period near Premier Lake. This has been estimated that 200 to 3000 ha of open forest and grassland are lost.



Ingrowth often results in the retrogression of understory plant communities, the loss of habitat for sensitive wildlife species, and an increased risk of catastrophic wildfires. Changes in forest structure with ingrowth forests reduces forage availability for wildlife and livestock. For


A photograph of a snowy mountain landscape under a clear blue sky. In the center, a helicopter is landing on a snow-covered slope, kicking up a cloud of white snow. Two people in orange gear are visible near the helicopter. In the foreground, a herd of caribou is partially visible, partially obscured by the dark, silhouetted branches of evergreen trees. The background shows rolling snow-covered hills with scattered evergreen trees.

Wildlife Habitat Program

- ▶ Staff and locations
- ▶ Accomplishments
- ▶ What's new
- ▶ Future directions



*Don Morgan
Team Leader,
Wildlife Habitat
Smithers office*



Habitat Supply Research Network

[Home](#)[Research Areas](#)[Species](#)[News](#)[Links](#)[About the Network](#)[Network Members](#)[+Join the Network](#)

Select a
research
area...




Select a species...



The Habitat Supply Research Network is a member-maintained network of habitat researchers

SCENARIO MATRIX






*Bruce McLellan
Wildlife Habitat Biologist
Kamloops*

*Fred Hovey
Wildlife Habitat Modeller
Kamloops*

Summary

- ▶ Ecology, Watershed, Soils, Climate, Carbon, Landscape, Range and Wildlife Habitat
- ▶ Support clients across MFR and other agencies thru research, assessments and expert advice to develop policy and practices
- ▶ Climate change and carbon, resiliency, vulnerability assessments, bioenergy, provincial soils map, HSMs, watershed science synthesis, TSSDST, MPB, ClimateWNA, GHC, offset protocols and scenarios



Contact information

Evelyn Hamilton

Manager, Ecology and Earth Science Section

Ministry of Forests and Range, Research Branch,
4th floor, 727 Fisgard Street, Victoria, V8W 9C1

Phone (o) 250 387-3650 (c) 250 213 2808

email Evelyn.Hamilton@gov.bc.ca

website www.for.gov.bc.ca/hre

