

An aerial photograph of a vast forest landscape. In the center, a dark, calm lake is surrounded by dense evergreen forests. The terrain is hilly, with varying shades of green and brown indicating different forest types or perhaps the effects of a beetle outbreak. In the far distance, a range of mountains is visible under a cloudy sky. The overall scene is expansive and natural.

**Mountain pine beetle dispersal in
British Columbia, Canada:
spatiotemporal patterns
and
role in the spread and expansion of
the present outbreak**

Huapeng Chen and Adrian Walton

Outline


2

- Introduction
- Objectives
- Study area
- Data and method
- Results
- Discussion
- Conclusions

Introduction

3


- BC Mountain Pine Beetle (MPB) infestations
 - ▣ As of 2009, 47% or 630 million m³ of merchantable pines



Introduction

4

□ Life cycle of MPB



Introduction

5

- Dispersal is a key stage in the life circle of MPB
- Dispersal is a key population process in MPB
- A better understanding of dispersal:
 - ▣ Valuable insights into MPB outbreak dynamics
 - ▣ Help to minimize outbreak impacts, achieving sustainable forest management goals
- **However, dispersal is least understood in MPB**

Objectives

6

- What are the spatiotemporal patterns of short-distance dispersal (SDD) and long-distance dispersal (LDD)?
- What are the roles of SDD and especially LDD in the spread and expansion of the current MPB outbreak?

Objectives


7

- Hypotheses to be tested:
 - ▣ SDD is a dominant factor in the spread and expansion of the MPB outbreak
 - ▣ LDD is a key factor to initiate new infestations in remote areas
 - ▣ The occurrence of LDD is random, and if not, its occurrence is positively correlated to severity and patchiness of the MPB infestations

Study area

8

- Defined as the cumulative annual MPB infestation areas from 1999 to 2008



Data

9

- BC Provincial-Level MPB Model output raster datasets (1999 – 2008)
 - ▣ Produced using the BC aerial overview survey data
 - ▣ 400x400 meter cell size

Data


10

- Aerial overview (sketch mapping) survey data
 - ▣ Spatial extent (coverage): province
 - ▣ Temporal scale: 1914 up to present
 - ▣ Spatial scale: 1: 100, 000 to 1: 250, 000
 - ▣ Severity estimation: yes, based on a representative polygon drawn on a topographic map

Data

11

□ Aerial overview survey data



Severity classes

- T (Trace, < 1%)
- L (Light, 1 – 10%)
- M (Moderate, 11 – 30%)
- S (Severe, 31 – 50%)
- V (Very severe, 51 – 100%)

Data


12

□ Data processing

- ▣ Scaled the raster datasets to 1200 meters because a grid of the same size was used in the correction procedures
- ▣ Combined to create a master grid
- ▣ Severity of a grid cell were calculated from 9 cells of the raster datasets using an area weighted average
- ▣ Severities of the grid cells were re-grouped into two general classes: light infestation (L), combining T, L, and M, and severe infestation (S), combining S and V
- ▣ Total number of master grid cells: 165,731


Data

13



Raster datasets

Up scale




Master grid

Method

14

□ Conceptual model of MPB infestations



Method





15

□ Conceptual model of MPB infestations

	SINK, NL and NS	Beetles only fly into the cell
	SOURCE, LN and SN	Beetles only fly out of the cell
	TRANSFORMER, LL and SS, LS, and SL	Beetles fly into and out of the cell, being a SINK and SOURCE

NL	NL	NS	NS	LL
NL	NS	NS	LL	LL
SS	SS	LN	LN	SN
SS	SS	LN	SN	SN
LS	LS	SL	SL	SL



 NL	 NL	 NS	 NS	 LL
 NL	 NS	 NS	 LL	 LL
 SS	 SS	 LN	 LN	 SN
 SS	 SS	 LN	 SN	 SN
 LS	 LS	 SL	 SL	 SL

Method


16

- Key assumptions for the conceptual model of MPB infestations
 - ▣ The beetles fly from a source to a sink
 - ▣ The infestation severity is a good proxy of beetle population abundance
 - ▣ It takes a year for infested trees to fade from green to red attack they have been attacked
 - ▣ The infested trees always produce a brood of beetles in the next year

Method

17


- Characterize dispersal patterns
 - Grid cells were grouped into five types: (1) sink NL and NS; (2) source LN and SN; (3) transformer LL and SS; (4) transformer LS; and (5) transformer SL.



Method

18

- Characterize dispersal patterns
 - ▣ Four types of potential sinks



- ▣ Four types of potential sources



TRANSFORMERS




Method

19

□ Characterize dispersal patterns

- Measured the distance from the edge of each individual sink patch to the edge of the nearest individual source patch



Method

20

- Characterize dispersal patterns
 - ▣ Percentage of three modes of dispersal used to characterize dispersal pattern
 - ▣ Results only reported for:
 - Sink type NL and NS, dispersal to non-infested areas, ie., new infestations
 - Sink type LL and SS, dispersal to infested areas, ie., old infestations

Method

21


□ Statistical analyses

- ▣ The chi-square goodness fit was used to test if the occurrences of three modes of dispersal to both non-infested and infested areas are random
- ▣ If the test was rejected for LDD, the Spearman rank correlation coefficients between the occurrences of LDD and the total number of infested grid cells as well as landscape aggregation indices

Results

22



□ Dynamics of the mountain pine beetle outbreak



Results

23


□ Dispersal at the regional (provincial) scale



Results

24


□ LDD distance distribution












Results





25



- Dispersal at the local (district) scale
 - ▣ The occurrences of three modes of mountain pine beetle dispersal to non-infested areas
















Results

32

- Dispersal at the local (district) scale
 - ▣ The occurrences of three modes of mountain pine beetle dispersal to infested areas




Results

34

- Dispersal at the local (district) scale
 - ▣ Spearman rank coefficients of occurrences of long-distance dispersal (LDD) with total number of infested grid cells and landscape aggregation indices (AI) of the infestations
 - LDD to non-infested areas have stronger negative rank correlations between its occurrences and total number of infested grid cells, and aggregation indices of the infestations than LDD to infested areas for all districts and for most individual districts


Infested cell number

ALL (-0.6058)*



Landscape AI


ALL (-0.5080)*



Results

37

- Dispersal at the local (district) scale
 - ▣ Means of landscape aggregation indices of the infestation patches



Discussion

38

- LDD plays a more important role in the early stage of infestations
 - ▣ As the infestations intensify and expand (assumed populations reaching epidemic levels), rapid depletion of the host resources may largely decrease the possibility of successful colonization for the pioneer beetles from LDD because LDD may be more costly in energy and the pioneer beetles from LDD may have a higher expectation and requirement in terms of quantity and quality of the hosts

Conclusions

39

- The conceptual framework of mountain pine beetle infestations proposed in this study provides a practical and analytical approach to quantify dispersal and detect its spatiotemporal patterns at the landscape scale. This approach can be applied to other bark beetle species with long-term and sufficient monitoring data.
- The conceptual framework of MPB could be used to determine stages of infestation dynamics


Conclusions

40

- LDD is a key factor in the initiation and early stage of the infestations in new remote areas and may play a critical role in expansion of the outbreak through facilitating the shifts of the beetle populations from incipient-epidemic levels to epidemic levels.

2007

- Long-distance dispersal sinks
- Long-distance dispersal sources



Conclusions

42

- The spatiotemporal patterns of LDD can be used to detect and map potential conduits of LDD at the landscape scale, assisting to identify and prioritize the mitigation areas in strategic and tactical planning.
- ▣ Implications to MPB management
 - LDD patterns should be considered in MPB provincial management zonations
 - LDD patterns should be considered in making suppression strategies for the BMUs
 - LDD should be considered in MPB risk rating or susceptibility systems

Conclusions

43

□ Future research

▣ Mechanisms driving LDD

- Link topographic modeling and meso-scale meteorological modeling together
- Explore how topographic features interact with meso-scale meteorological events, affecting spatial patterns of LDD at the landscape scale

Question??

44

