

SAR

Species at Risk

Key Topographic or Distinguishing Features

Report Criteria

TSA:	Golden	Prov Stat:	
BEC:	MS dk	Sec7	

Species	Prov Status	Key Stand Features	Crown Closure	Critical Aspect 1	Slope Range Comment	Age Class	Topographic or Distinguishing Feature	Max Elevation	Lead Species	Additional Management Considerations	Likelihood
 Barn Swallow (Hirundo rustico)	Blue-listed	Riparian habitats and wetlands	Not Available	Not Available	Not Relevant	Not Available	Wetland, riparian, aquatic systems, Open forest or grasslands	Not available	Not Available	Likely use clearcuts since prefer open habitats; also, may use vets since nest in buildings, barns, etc. Retain herbaceous vegetation near wetlands/riparian	Not available
 Bighorn Sheep (Ovis canadensis)	Blue-listed	Snow-interception, open, steep south aspect slopes; talus; mineral licks; watering features	Not Available	S	27°<Escape terrain<85°	1-9	Steep rocky slopes	3000m	Se, Pl, Fd	Avoid introducing invasive species (non-native plants, weeds); retain habitat features in seasonal habitats (old-growth in lower elevation winter habitats). Consider intensive silviculture or habitat enhancement (spacing/thinning) outside Apr-Jul, Oct-Nov.	Moderate
 Fisher (Martes pennanti)	Blue-listed	large, elevated CWD (>65cm DBH), decay classes 2-6 (logs), largest coniferous and deciduous trees available, particularly with heart rot and/or cavities, trees with brooms, wildlife tree classes 2-3 preferred, 2-6 acceptable		Not Available	Slope likely not applicable, critical habitat component: stand structure	>4	River/stream, wetlands	2500	Not Applicable	Old forest structure more important than forest type. Large diameter Act critical for breeding, largest trees within a landscape should be retained; highest habitat suitability are characteristics of old growth SBS, SWB, and BWBS zones, particularly riparian/wetland; retain trees with heart rot and >54cm bole diameter at 5m above ground; maintain connectivity between upland and riparian.	Moderate-High
Great Blue Heron (Ardea herodias herodias)	Blue-listed	Riparian forest, including Act and mixed conif-decid, trees capable of supporting nest platform 50-100cm diameter	Not Available	Not Relevant	Not Relevant	>2	valley bottoms	1100	Act, Cw, Fd, Hw, Sx	Avoid disturbance mid-February-August 31 (breeding period) within 500m of breeding colonies; no harvest core (nest/colony), partial cut 300m buffer, clearcut beyond	Low
 Grizzly Bear (Ursus arctos)	Blue-listed	Riparian habitats and wetlands, avalanche tracks and run out zones, Hedysarum (pea family/vetch) and glacier lily complexes, Sub-alpine parkland meadows, berry-producing sites	Not Available	N (denning)	generally steep, north aspects (denning)	1-9	Mid-slope (for denning), Slides (avalanche chutes), talus slopes, alpine meadows, wetlands	alpine	Not Applicable	Grizzly bear mgmt should follow a broad landscape approach; listed here are stand and watershed requirements only; see high level plans (ie: FSP results and strategies). At cutblock level, key is to remove bear attractants, and at watershed level implement aggressive access mgmt plan that deactivates roads asap. Across landscape and over time, early seral stages that produce herbs and berries are critical for foraging.	Moderate
 Lewis's Woodpecker (Melanerpes lewis)	Red-listed	Preferred tree size >44cm, in absence of large trees retain >29cm (acceptable)	<25%	west side of valley	Not Relevant	5-9	Not Applicable	1160	Act, At, Fd	Nests found >1000m elevation were at burn edges; used shrub, herbaceous openings to forage for insects.	Very Low

Species		Prov Status	Key Stand Features	Crown Closure	Critical Aspect 1	Slope Range Comment	Age Class	Topographic or Distinguishing Feature	Max Elevation	Lead Species	Additional Management Considerations	Likelihood
	Northern Leopard Frog (<i>Rana pipiens</i>)	Red-listed	Not Applicable, frogs avoid wooded areas; but will travel overland in 15-30cm tall veg.	Not Applicable	Not Available	Not Relevant	Not Applicable	Wetland, riparian, aquatic systems	1400	Not applicable ; avoid road construction	Frogs breed in shallow ponds or wetlands, may disperse through connecting riparian areas to deeper ponds that do not freeze to the bottom, where they remain overwinter. This core area (1x2km) should be protected from traffic and livestock, with a 50m buffer (management zone). Winter area =1.7km-800m from breeding and adjacent terrestrial foraging habitat.	Very Low
	Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Blue-listed	forest openings, Snags, riparian forest, tall live conifers, standing dead trees taller than surrounding forest	Not Available	Not Available	May not be relevant if wetland/riparian/bog dependent	Not Available	Wetland, riparian, aquatic systems, marsh, bog/fen; ponds	Not available	Not Available	Variable height snags appear to be critical: used for perching, foraging (insect capture). Live coniferous trees may be most important for nest placement (2-15m above ground). Harvest systems should mimic natural disturbance regime: fire, insect, flood, where applicable (ie: openings with standing dead trees some taller than surrounding forest).	Moderate
	Sandhill Crane (<i>Grus canadensis</i>)	Blue-listed	Riparian forest concealing characteristics, wetlands	Not Available	Not Available	Flat terrain	0-9	Wetland, riparian, aquatic systems	1220	Not Available	Maintain shallow freshwater wetlands/aquatic systems, retain adjacent riparian forests; avoid harvesting within 800m of aquatic systems used for breeding Apr-Aug (breeding season)	Low-Moderate
	Wolverine (<i>Gulo gulo</i>)	Blue-listed	Mature - old-growth structure	Not Applicable	Not applicable	Not Relevant	>4	valley bottoms, Riparian corridors	Not available	Not Applicable	Landscape planning is very important for wide-ranging wolverines: adjacency of early and late seral structural stages is important to provide habitat features across all BEC (elevations); avoid development on both sides of a watershed at the same time and provide old forest connectivity between watersheds as travel corridors.	High
	Yellow Rail (<i>Coturnicops noveboracensis</i>)	Red-listed	wetlands	Not Available	Not Available	Likely flat is most applicable, since nest and forage for insects at/near water.	Not Available	wetlands	Not available	Not available; may not be relevant: existing shore/water's edge trees	Very little information is available for this species and it is documented only in the MS and BWBS zones in the Columbia & Peace FDs, respectively. May be prudent to buffer all wetlands with min 10m no harvest/disturbance, including ephemeral sites - apply riparian regulations regardless of wetland size to ensure existing riparian forest, if any, and/or surrounding features, are retained.	Unknown
	American Bittern (<i>Botaurus lentiginosus</i>)	Blue-listed	wetlands	Not Applicable	Not Available	Likely flat is most applicable, since nest and forage for in/near water.	Not Applicable	Wetland, riparian, aquatic systems	1300m	Not Applicable	Large wetlands, and aquatic and wetland complexes, should be avoided for any resource use, including timber harvest. For harvesting: maintain a buffer of existing stand, if present, surrounding wetland to avoid impact or disturbance; if grassland surrounding wetland, maintain existing grassland buffer with no roads, etc, sufficient to avoid impact. Dense emergent vegetation is a seral stage and wetland management (MoE?) should ensure sustainability by monitoring and maintaining water levels. Riparian Area Mgmt Guidebook may be used to determine buffer size.	